

The Relationship Between Sea Surface Temperature and Ocean Acidification in Shanghai during 1990 and 2010

[Speaker
Zoom
video]

Bactrosaurus_Lavani_Energico




Climatematch
Academy

Study Area

◆ Shanghai

East China Sea (ECS) - Shanghai's coastal region






We take the city or social community of reference to the metropolis of Shanghai and its coastal region.

It is a semi-enclosed marginal sea spanning an area of 744,000 km², and its boundaries are approximately 25° to 40° North and 120° to 130° East



Scientific background

- The sea surface temperature (SST) of Shanghai coastal region had experienced a drastic upward trend in the last 100 years, increasing by around 1.5 degree Celsius.
- In contrast, the pH level of this region decreased rapidly from 8.13 in 1900 to 8.06 in 2010, indicating the occurrence of ocean acidification.
- The ECS is generally considered a CO₂ source in autumn and a CO₂ sink in other seasons, based on limited field observation.



Source: Liu & Zhang, (2013); Liu et al. (2022); Liu et al. (2023)



Knowledge gap



Our hypothesis

How does the sea surface temperature (SST) influences the ocean pH and pCO₂ value in Shanghai's coastal and marine region in different seasons (during 1990 to 2010)?

- We expect that two variables we focus on are inversely proportional (i.e., increase sea surface temperature means decreased pH).
- In same way, we expect that two variables are inversely proportional (i.e., increase sea surface temperature means decreased pCO₂).
- In the warm season (i.e., spring and summer), increase in dissolved inorganic carbon will result in decreased pH level.



Datasets

CO2 concentration

Trends in Atmospheric -
Ocean Carbon Dioxide

Source: NOAA Global Monitoring
Laboratory

Units: u-atm

Version: 2023-07

pH level

Global surface ocean
acidification indicators from
1750 to 2100

Source: NOAA National Centers
for Environmental Information

Abbreviation: pHT

pH scale: Total scale (T)

Sea Surface Temperature (SST)


Source: NOAA Global Monitoring
Laboratory and Coupled Model
Intercomparison Project (CMIP)

Abbreviation: tos

Units: Celsius degrees




Results - pCO₂ trend



In the selected time interval, the partial pressure of carbon dioxide (pCO₂) in the waters of the East China Sea (ECS) near the coast of Shanghai increased by about 20 uatm from 1990 to 2010. Likewise, the concentration takes values higher than the average concentrations of CO₂ in the world; which could indicate a greater susceptibility to capture atmospheric CO₂.




Results - pCO₂ trend



In this graph we can see the evolution at the rate of decades with respect to pCO₂, with a marked tendency to increase the concentration of CO₂, specifically the regions closest to the Shanghai coast (circled in red).




Results - Sea Surface Temperature trend



Regarding the surface temperature of the East China Sea, it can be observed that there are insignificant changes throughout the interval 1990 and 2010, where temperature fluctuated between the range of -1.5 to 1.5 °C, however a slight tendency of increase can be seen here.



Results - pH trend




Along the same lines, it can be seen that the pH of the waters of the East China Sea (ECS) decreased to values very close to 8 between the years 1990 and 2010, which is a sign of ocean acidification off the coast of Shanghai.

The perceived magnitude of that alteration may not seem substantial at first glance, but due to pH being measured on a logarithmic scale, it corresponds to an approximately 30% rise in acidity. As a result, this change has notable repercussions on seawater chemistry and the dependent ecosystems.




Results - pH trend (annual)



In this graph we can see the decrease in the pH level, specifically the regions closest to the coast of Shanghai (circled in red), which indicates greater acidity conditions for the ecosystems coastal marine. This would be attributed to the increased CO₂ concentration, the incoming flows of nutrients and other chemical substances from Chajiang River, or sea surface temperature (SST).



Results - pH trend (seasonal)



Here we can see the marked changes that the pH level has in the months of the year, showing the influence of seasonality. Reaching conditions of greater acidity in the warm months compared to the months of cold conditions of the sea.




Significance and societal impact

The region's rugged coastlines and well-developed continental shelf makes it suitable for marine fisheries and coastal aquaculture of finfish, shellfish, and seaweed (Kang 2006).



The ECS is characterised by high biodiversity (Liu 2013a), with up to 12,933 tropical and subtropical species. Importantly, almost half (48%) of ECS species are endemic (Ding et al. 2008).

Clams Distribution in ECS



Based on this information, we can see that ocean acidification would have impacts on marine biodiversity and risk the stability of fishing and aquaculture activities.

Conclusions

- The pH level and sea surface temperature in Shanghai coastal region are inversely proportional (i.e., increased SST means decreased pH).
- We can conclude that SST has a seasonal effect on pH, since during hot seasons in Shanghai, the pH level decreases significantly, contrast to the cold seasons.
- Societal impacts have not been made clear in any form of public communication; however, we should be aware of the potential influences of ocean acidification.

