

Fall 2025 ECEN 629 Homework Set 3

Due Oct. 21 12:45pm online (or 2pm in class)

Prob. 1 Suppose $f : \mathbf{R}^n \rightarrow \mathbf{R}$ is convex, and define

$$g(x, t) = f(x/t), \quad \text{dom } g = \{(x, t) | x/t \in \text{dom } f, t > 0\}.$$

Show that g is quasiconvex, and then write down the optimization procedure similarly to that on page 17 of the slides of Chapter 7, specialized to this problem.

Hint: consider the function $t \cdot f(x/t)$.

Prob. 2 Some non-convex problems can also be easy to solve. Find the optimal solution for the following problems in closed form.

(a) For $x \in \mathbf{R}^n$

$$\begin{aligned} &\text{minimize: } \|x\|_2 \\ &\text{subject to: } 10 \geq \|x\|_1 \geq 1. \end{aligned}$$

(b) Let $x = (x_1, x_2) \in \mathbf{R}^2$

$$\begin{aligned} &\text{minimize: } x_1 + x_2 \\ &\text{subject to: } x_1 + 3x_2 = 5; \\ &\quad (x_1 - 3)^2 + (x_2 - 4)^2 \geq 25. \end{aligned}$$

Prob. 3 Huber's penalty function is defined as

$$\phi(u) = \begin{cases} u^2 & |u| \leq M \\ M(2|u| - M) & |u| > M \end{cases}.$$

(a) Is the function $\phi(u)$ convex in general?

(b) Show that the minimization problem

$$\text{minimize: } \sum_{i=1}^m \phi(a_i^T x - b_i)$$

is equivalent to the following problem

$$\begin{aligned} &\text{minimize: } \sum_{i=1}^m (u_i^2 + 2Mv_i) \\ &\text{subject to: } -u - v \preceq Ax - b \preceq u + v \\ &\quad 0 \preceq u \preceq M\mathbf{1} \\ &\quad v \succeq 0. \end{aligned}$$

(c) A general form of Huber's penalty function with parameters (α, β, γ) is

$$\phi(u) = \begin{cases} \alpha \cdot u^2 & |u| \leq M \\ \beta M(\gamma|u| - M) & |u| > M \end{cases}.$$

For any fixed value of $\alpha > 0$, find the valid assignment of (β, γ) in terms of α such that the penalty function is convex.

Prob. 4 Write the following problems as equivalent LPs:

(a)

$$\begin{aligned} &\text{minimize: } \|x\|_\infty \\ &\text{subject to: } \|Ax - b\|_1 \leq 1 \end{aligned}$$

where $A \in \mathbb{R}^{m \times n}$ and $b \in \mathbb{R}^m$.

(b)

$$\begin{aligned} &\text{minimize: } \|x\|_\infty \\ &\text{subject to: } \|Ax - b\|_\infty \leq 1 \end{aligned}$$

where $A \in \mathbb{R}^{m \times n}$ and $b \in \mathbb{R}^m$.

(c)

$$\begin{aligned} &\text{minimize: } \|x\|_1 \\ &\text{subject to: } \|Ax - b\|_\infty \leq 1 \end{aligned}$$

where $A \in \mathbb{R}^{m \times n}$ and $b \in \mathbb{R}^m$.

(d)

$$\begin{aligned} &\text{minimize: } \|x\|_1 \\ &\text{subject to: } \|Ax - b\|_2 \leq 1 \end{aligned}$$

where $A \in \mathbb{R}^{m \times n}$ and $b \in \mathbb{R}^m$.

For the problems above, what if we exchange the objective function and the constraint, can you still write down the problems as LPs?

Prob. 5 Consider the optimization problem in $\mathbb{R}^2$

$$\begin{aligned} &\text{minimize: } f_0(x_1, x_2) \\ &\text{subject to: } (x_1 - 1)^2 + (x_2 - 2)^2 \leq 2 \\ &\quad 2x_1 + x_2 \geq 3 \\ &\quad x_1 \geq 0, x_2 \geq 0. \end{aligned}$$

First make a sketch of the feasible set. For each of the following objective functions, give the optimal set and the optimal value, using two methods: geometrically and the first-order optimality condition. Note that you should not use KKT directly here, but the first-order condition in Chapter 4.

- (a) $f_0(x_1, x_2) = x_1 + x_2$;
- (b) $f_0(x_1, x_2) = x_1^2 + x_2^2$.

Prob. 6 Consider the optimization problem

$$\begin{aligned} & \text{minimize: } f_0(x_1, x_2) \\ & \text{subject to: } x_1 + x_2 \geq 2 \\ & \quad 2x_1 + x_2 \geq 3 \\ & \quad x_1 \geq 0, x_2 \geq 0. \end{aligned}$$

First make a sketch of the feasible set. For each of the following objective functions, give the optimal set and the optimal value, using two methods: geometrically and the first-order optimality condition.

- (a) $f_0(x_1, x_2) = 3x_1 + 2x_2$;
- (b) $f_0(x_1, x_2) = \max\{x_1, x_2\}$.

Prob. 7 Give explicit solutions for the following LP problems over $\mathbf{R}^n$

(a)

$$\begin{aligned} & \text{minimize: } c^T x \\ & \text{subject to: } a^T x \leq b, \end{aligned}$$

where $a \neq 0$.

(b)

$$\begin{aligned} & \text{minimize: } c^T x \\ & \text{subject to: } \mathbf{1}^T x \leq \alpha \\ & \quad 0 \cdot \mathbf{1} \preceq l \preceq x \preceq u. \end{aligned}$$

Prob. 8. Find the dual optimization problems for the following optimization problems

(a)

$$\begin{aligned} & \text{minimize: } c^T x \\ & \text{subject to: } Ax = b \\ & \quad \ell \preceq x \preceq u, \end{aligned}$$

where $x \in \mathbf{R}^n$ is the variable, and $\ell, u \in \mathbf{R}^n$.

(b) Recall last time you wrote the following problem as an equivalent LP:

$$\begin{aligned} & \text{minimize: } \|x\|_\infty \\ & \text{subject to: } \|Ax - b\|_\infty \leq 1 \end{aligned}$$

where $A \in \mathbb{R}^{m \times n}$ and $b \in \mathbb{R}^m$. Find the dual problem for the equivalent LP.

(c)

$$\begin{aligned} & \text{minimize: } \sum_{i=1}^n x_i \log x_i \\ & \text{subject to: } Ax \succeq b \\ & \quad c^T x = d, \end{aligned}$$

whose domain is R_+^n , i.e., $x_i > 0$ for all $i = 1, 2, \dots, n$. Assume $c \succeq 0$.

(d)

$$\begin{aligned} & \text{minimize: } x^T Q x \\ & \text{subject to: } Ax = b, \end{aligned}$$

where Q is a positive definite matrix.