

Maximum Smoothness Algorithm for Building Forward Curves

Jake C. Fowler

June 2019

1 Introduction

This document describes the maximum-smoothness algorithm used to increase the granularity of constructed forward curves. The rest of this document will refer to forward prices and curves, even though the same methodology can be applied to futures and swap instruments as well.

The methodology is an adaption of that described in (Author). This makes reference to Benth et al. (2007) and Lim and Xiao (2002), the later reference which uses maximum smoothness in the context of building interest rate forward curves.

2 Deriving the Algorithm

2.1 Functional Form

The base of the algorithm is a spline, which by definition is made up of piecewise polynomial functions.

$$p(t) = \begin{cases} p_1(t) & \text{for } t \in [t_0, t_1) \\ p_2(t) & \text{for } t \in [t_1, t_2) \\ \vdots & \\ p_{n-1}(t) & \text{for } t \in [t_{n-2}, t_{n-1}) \\ p_n(t) & \text{for } t \in [t_{n-1}, t_n] \end{cases} \quad (1)$$

Where $t_0 < t_1 < \dots < t_{n-1} < t_n$ are the boundary points between the polynomials which make up the spline. In the context of building a forward curve, the variable t is defined as the time until start of delivery of a forward contract.

The boundary points are chosen to be start of the input forward prices. It is also assumed that the input forward prices are not for delivery periods

which overlap with any other input. Gaps between input forward contracts are permitted, in which case a boundary point will exit for the start of the gap.

The individual polynomial functions p_n themselves are of order 4, hence take the form:

$$p_i(t) = a_i + b_i t + c_i t^2 + d_i t^3 + e_i t^4 \quad (2)$$

The curve fitting algorithm essentially involves solving for the polynomial parameters a_i , b_i , c_i , d_i , and e_i for $i = 1 \dots n$.

In many cases the spline described above is not sufficient to derive a forward curve which shows strong price seasonality, especially when this seasonality cannot be directly observed in the traded forward prices. An example of this is the day-of-week seasonality for gas and power prices, which generally are lower at the weekend when demand is lower. As such the function form is as follows:

$$f(t) = (p(t) + S_{add}(t))S_{mult}(t) \quad (3)$$

Where the forward price for the period starting delivery at time t is given by $f(t)$, which consists of $p(t)$ adjusted by two arbitrary seasonal adjustment functions $S_{add}(t)$ an additive adjustment, and $S_{mult}(t)$ a multiplicative adjustment.

2.2 Constraints

2.2.1 Polynomial Boundary Point Constraints

As usual with splines, constraints are put in place that adjacent polynomials have equal value, first derivative, and second derivatives at the boundary points. These three constraints can be respectively expressed as:

$$a_i + b_i t + c_i t^2 + d_i t^3 + e_i t^4 - a_{i+1} - b_{i+1} t - c_{i+1} t^2 - d_{i+1} t^3 - e_{i+1} t^4 = 0 \quad (4)$$

$$b_i + 2c_i t + 3d_i t^2 + 4e_i t^3 - b_{i+1} - 2c_{i+1} t - 3d_{i+1} t^2 - 4e_{i+1} t^3 = 0 \quad (5)$$

$$2c_i + 6d_i t + 12e_i t^2 - 2c_{i+1} - 6d_{i+1} t - 12e_{i+1} t^2 = 0 \quad (6)$$

The above three equations should hold for the boundary points $t \in \{t_1, t_2, \dots, t_{n-2}, t_{n-1}\}$.

2.2.2 Forward Price Constraint

The most important constraints is that the derived forward curve averages back to the input traded forward prices. The market inputs to the forward curve model are traded forward prices F_i . Setting this equal to the average of the derived smooth curve:

$$F_i = \frac{\sum_{t \in T_i} (p(t) + S_{add}(t)) S_{mult}(t) w(t)}{\sum_{t \in T_i} w(t)} \quad (7)$$

Where $w(t)$ is a weighting function and T_i is the set of all delivery start times for the delivery periods at the granularity of the curve being built. The weighting function has two meanings from a business perspective.

- The discount factor, because the no-arbitrage price of a forward contract is equal to the discount factor weighted average of its components. However, in a low interest rate environment, the discount factor can be approximated as 1.0 for all maturities, in which case the discounting component can be ignored.
- The volume of commodity delivered in each period. For example, an off-peak power forward contract in the UK delivers over 12 hours in on weekdays, and 24 hours on weekends, hence $w(t)$ would equal double for t representing weekends compared to $w(t)$ when t represents a weekday delivery. Clock changes can also cause the total volume delivered over a day in a fixed time zone to vary due to hours lost or gained. Hence $w(t)$ can be used to account for this.
- For swaps which only fix on certain days (usually business days) $w(t)$ can be used to account for this by returning the number of fixing days in the period starting at t . For example if deriving a monthly curve $w(t)$ would evaluate to the number of fixing days in the month starting at t .

Equation 7 can be transformed into an equation linear on the parameters of the piecewise polynomial by substituting in the polynomial representation of $p(t)$:

$$\sum_{t \in T_i} (p_i(t) + S_{add}(t)) S_{mult}(t) w(t) = F_i \sum_{t \in T_i} w(t) \quad (8)$$

Rearranging:

$$\sum_{t \in T_i} (a_i + b_i t + c_i t^2 + d_i t^3 + e_i t^4 + S_{add}(t)) S_{mult}(t) w(t) = F_i \sum_{t \in T_i} w(t) \quad (9)$$

Rearranging again gives a form linear with respect to the unknown polynomial coefficients a_i , b_i , c_i and d_i .

$$\begin{aligned} a_i \sum_{t \in T_i} S_{mult}(t) w(t) + b_i \sum_{t \in T_i} S_{mult}(t) w(t) t + c_i \sum_{t \in T_i} S_{mult}(t) w(t) t^2 + \\ d_i \sum_{t \in T_i} S_{mult}(t) w(t) t^3 + e_i \sum_{t \in T_i} S_{mult}(t) w(t) t^4 = \\ F_i \sum_{t \in T_i} w(t) - \sum_{t \in T_i} S_{add}(t) S_{mult}(t) w(t) \end{aligned} \quad (10)$$

In the case where there are gaps between the input forward contract delivery periods, the forward price constraint is simply omitted.

2.2.3 Matrix Form of Constraints

Equations 4, 5, 6 and 10 can be expressed as the linear system $\mathbf{Ax} = \mathbf{b}$ where:

$$\mathbf{x} = \begin{bmatrix} \mathbf{x}_1 \\ \mathbf{x}_2 \\ \vdots \\ \mathbf{x}_{n-1} \\ \mathbf{x}_n \end{bmatrix}$$

$$\mathbf{b} = \begin{bmatrix} \mathbf{b}_1 \\ \mathbf{b}_2 \\ \vdots \\ \mathbf{b}_{n-1} \\ \mathbf{b}_n \end{bmatrix}$$

$$\mathbf{A} = \begin{bmatrix} \mathbf{A}_1 & \mathbf{0} & \dots & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{A}_2 & \dots & \mathbf{0} & \mathbf{0} \\ \vdots & & \ddots & & \vdots \\ \mathbf{0} & \mathbf{0} & & \mathbf{A}_{n-1} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \dots & \mathbf{0} & \mathbf{A}_n \end{bmatrix}$$

And:

$$\mathbf{x}_i = \begin{bmatrix} a_i \\ b_i \\ c_i \\ d_i \\ e_i \end{bmatrix}$$

$$\mathbf{b}_i = \begin{bmatrix} 0 \\ 0 \\ 0 \\ F_i \sum_{t \in T_i} w(t) - \sum_{t \in T_i} S_{add}(t) S_{mult}(t) w(t) \end{bmatrix}$$

$$\mathbf{A}_i = \begin{bmatrix} 1 & t_i & t_i^2 & t_i^3 & t_i^4 & -1 & -t_i & -t_i^2 & -t_i^3 & -t_i^4 \\ 0 & 1 & 2t_i & 3t_i^2 & 4t_i^3 & 0 & -1 & -2t_i & -3t_i^2 & -4t_i^3 \\ 0 & 0 & 2 & 6t & 12t^2 & 0 & 0 & -2 & -6t & -12t^2 \\ f_i^1 & f_i^2 & f_i^3 & f_i^4 & f_i^5 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Where the components in the last row are defined as:

$$\begin{aligned}
f_i^1 &= \sum_{t \in T_i} S_{mult}(t)w(t) \\
f_i^2 &= \sum_{t \in T_i} S_{mult}(t)w(t)t \\
f_i^3 &= \sum_{t \in T_i} S_{mult}(t)w(t)t^2 \\
f_i^4 &= \sum_{t \in T_i} S_{mult}(t)w(t)t^3 \\
f_i^5 &= \sum_{t \in T_i} S_{mult}(t)w(t)t^4
\end{aligned}$$

2.3 Smoothness Criteria

Maximum smoothness is obtained by finding the spline parameters which minimise the integral of the second derivative.

$$\begin{aligned}
\min \int_{t_0}^{t_n} p''(t)^2 dt &= \sum_{i=1}^n \int_{t_{i-1}}^{t_i} p_i''(t)^2 dt \\
&= \sum_{i=1}^n \int_{t_{i-1}}^{t_i} (2c_i + 6d_i t + 12e_i t^2)^2 dt \\
&= \sum_{i=1}^n \int_{t_{i-1}}^{t_i} (4c_i^2 + 24c_i d_i t + 48c_i e_i t^2 + 36d_i^2 t^2 + 144d_i e_i t^3 + 144e_i^2 t^4) dt \\
&= \sum_{i=1}^n 4c_i^2 \Delta_i^1 + 12c_i d_i \Delta_i^2 + 16c_i e_i \Delta_i^3 + 12d_i^2 \Delta_i^3 + 36d_i e_i \Delta_i^4 + \frac{144}{5} e_i^2 \Delta_i^5 \quad (11)
\end{aligned}$$

Where Δ_i^j is defined as the difference between t^j at the polynomial boundary points, i.e. $\Delta_i^j = t_i^j - t_{i-1}^j$. Recognising 11 as a quadratic form it can be reformulating in the following matrix form:

$$\sum_{i=1}^n \mathbf{x}_i^T \mathbf{H}_i \mathbf{x}_i \quad (12)$$

Where:

$$\mathbf{H}_i = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 4\Delta_i^1 & 6\Delta_i^2 & 8\Delta_i^3 \\ 0 & 0 & 6\Delta_i^2 & 12\Delta_i^3 & 18\Delta_i^4 \\ 0 & 0 & 8\Delta_i^3 & 18\Delta_i^4 & \frac{144}{5}\Delta_i^5 \end{bmatrix}$$

The objective function 12 can be arranged into a single matrix quadratic form without the summation as:

$$\mathbf{x}^T \mathbf{H} \mathbf{x} \quad (13)$$

Where:

$$\mathbf{x} = \begin{bmatrix} \mathbf{x}_1 \\ \mathbf{x}_2 \\ \vdots \\ \mathbf{x}_{n-1} \\ \mathbf{x}_n \end{bmatrix}$$

$$\mathbf{H} = \begin{bmatrix} \mathbf{H}_1 & \mathbf{0} & \dots & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{H}_2 & \dots & \mathbf{0} & \mathbf{0} \\ \vdots & & \ddots & & \vdots \\ \mathbf{0} & \mathbf{0} & & \mathbf{H}_{n-1} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \dots & \mathbf{0} & \mathbf{H}_n \end{bmatrix}$$

2.4 Minimisation Problem

The sections above show that finding the maximum smoothness curve comes down to finding the polynomial coefficients, vector \mathbf{x} , which minimises $\mathbf{x}^T \mathbf{H} \mathbf{x}$, subject to the linear constraints $\mathbf{A} \mathbf{x} = \mathbf{b}$. This problem is well suited to the method of Lagrange multipliers for which we first define the vector λ .

$$\lambda = \begin{bmatrix} \lambda_1 \\ \lambda_2 \\ \vdots \\ \lambda_{4n-2} \\ \lambda_{4n-1} \end{bmatrix} \quad (14)$$

$$\mathcal{L}(\mathbf{x}, \lambda) = \mathbf{x}^T \mathbf{H} \mathbf{x} + \lambda^T (\mathbf{A} \mathbf{x} - \mathbf{b}) \quad (15)$$

The minima $\min_{\mathbf{x}, \lambda} \mathcal{L}(\mathbf{x}, \lambda)$ is found as the solution where the partial derivatives of $\mathcal{L}(\mathbf{x}, \lambda)$ with respect to \mathbf{x} and λ are zero.

$$\frac{\partial \mathcal{L}(\mathbf{x}, \lambda)}{\partial \mathbf{x}} = 2\mathbf{H} \mathbf{x} + \mathbf{A}^T \lambda = 0 \quad (16)$$

$$\frac{\partial \mathcal{L}(\mathbf{x}, \lambda)}{\partial \lambda} = \mathbf{A} \mathbf{x} - \mathbf{b} = 0 \quad (17)$$

These can be arranged into a single linear system:

$$\begin{bmatrix} 2\mathbf{H} & \mathbf{A}^T \\ \mathbf{A} & \mathbf{0} \end{bmatrix} \begin{bmatrix} \mathbf{x}_{\min} \\ \lambda_{\min} \end{bmatrix} = \begin{bmatrix} \mathbf{0} \\ \mathbf{b} \end{bmatrix} \quad (18)$$

Hence the vector of spline polynomial coefficients for the maximum smoothness curve, \mathbf{x}_{\min} , can be found by solving this system.

$$\begin{bmatrix} \mathbf{x}_{\min} \\ \lambda_{\min} \end{bmatrix} = \begin{bmatrix} \mathbf{2H} & \mathbf{A}^T \\ \mathbf{A} & \mathbf{0} \end{bmatrix}^{-1} \begin{bmatrix} \mathbf{0} \\ \mathbf{b} \end{bmatrix} \quad (19)$$

Once 19 is solved, the spline parameters are taken from \mathbf{x}_{\min} and the derived forward prices are calculated by evaluating 3.

References

- Unknown Author. Constructing smooth forward curves in electricity markets. URL <https://www.maths.ox.ac.uk/system/files/attachments/M8%2013%20333715%20anonymised%20for%20pub.pdf>.
- F. E. Benth, S. Koekebakker, and F. Ollmar. Extracting and applying smooth forward curves from average-based commodity contracts with seasonal variation. *The Journal of Derivatives*, 15:52–66, 2007.
- K. G. Lim and Q. Xiao. Computing maximum smoothness forward rate curves. *Statistics and Computing*, 12:275–279, 2002.