

WIKIPEDIA

Neural oscillation

Neural oscillations, or **brainwaves**, are rhythmic or repetitive patterns of neural activity in the central nervous system. Neural tissue can generate oscillatory activity in many ways, driven either by mechanisms within individual neurons or by interactions between neurons. In individual neurons, oscillations can appear either as oscillations in membrane potential or as rhythmic patterns of action potentials, which then produce oscillatory activation of post-synaptic neurons. At the level of neural ensembles, synchronized activity of large numbers of neurons can give rise to macroscopic oscillations, which can be observed in an electroencephalogram. Oscillatory activity in groups of neurons generally arises from feedback connections between the neurons that result in the synchronization of their firing patterns. The interaction between neurons can give rise to oscillations at a different frequency than the firing frequency of individual neurons. A well-known example of macroscopic neural oscillations is alpha activity.

Neural oscillations in humans were observed by researchers as early as 1924 (by Hans Berger). More than 50 years later, intrinsic oscillatory behavior was encountered in vertebrate neurons, but its functional role is still not fully understood.^[1] The possible roles of neural oscillations include feature binding, information transfer mechanisms and the generation of rhythmic motor output. Over the last decades more insight has been gained, especially with advances in brain imaging. A major area of research in neuroscience involves determining how oscillations are generated and what their roles are. Oscillatory activity in the brain is widely observed at different levels of organization and is thought to play a key role in processing neural information. Numerous experimental studies support a functional role of neural oscillations; a unified interpretation, however, is still lacking.

Contents

History

Overview

Physiology

- Microscopic

- Mesososcopic

- Macroscopic

Mechanisms

- Neuronal properties

- Network properties

- Neuromodulation

Mathematical description

- Single neuron model

- Spiking model

Neural mass model

Kuramoto model

Activity patterns

Ongoing activity

Frequency response

Amplitude response

Phase resetting

Function

Pacemaker

Central pattern generator

Information processing

Perception

Motor coordination

Memory

Sleep and consciousness

Development

Pathology

Tremor

Epilepsy

Thalamocortical dysrhythmia

Applications

Clinical endpoints

Brain-computer interface


Examples

See also

References

Further reading

External links



Simulation of neural oscillations at 10 Hz. Upper panel shows spiking of individual neurons (with each dot representing an individual action potential within the population of neurons), and the lower panel the local field potential reflecting their summed activity. Figure illustrates how synchronized patterns of action potentials may result in macroscopic oscillations that can be measured outside the scalp.

History


Richard Caton discovered electrical activity in the cerebral hemispheres of rabbits and monkeys and presented his findings in 1875.^[2] Adolf Beck published in 1890 his observations of spontaneous electrical activity of the brain of rabbits and dogs that included rhythmic oscillations altered by light detected with electrodes directly placed on the surface of the brain.^[3] Before Hans Berger, Vladimir Vladimirovich Pravdich-Neminsky published the first animal EEG and the evoked potential of a dog.^[4]

Overview

Neural oscillations are observed throughout the central nervous system at all levels, and include spike trains, local field potentials and large-scale oscillations

which can be measured by electroencephalography (EEG). In general, oscillations can be characterized by their frequency, amplitude and phase. These signal properties can be extracted from neural recordings using time-frequency analysis. In large-scale oscillations, amplitude changes are considered to result from changes in synchronization within a neural ensemble, also referred to as local synchronization. In addition to local synchronization, oscillatory activity of distant neural structures (single neurons or neural ensembles) can synchronize. Neural oscillations and synchronization have been linked to many cognitive functions such as information transfer, perception, motor control and memory.^{[5][6][7][8]}

Neural oscillations have been most widely studied in neural activity generated by large groups of neurons. Large-scale activity can be measured by techniques such as EEG. In general, EEG signals have a broad spectral content similar to pink noise, but also reveal oscillatory activity in specific frequency bands. The first discovered and best-known frequency band is alpha activity (8–12 Hz)^{[9][10][11]} that can be detected from the occipital lobe during relaxed wakefulness and which increases when the eyes are closed.^[12] Other frequency bands are: delta (1–4 Hz), theta (4–8 Hz), beta (13–30 Hz), low gamma (30–70 Hz),^[13] and high gamma (70–150 Hz) frequency bands, where faster rhythms such as gamma activity have been linked to cognitive processing. Indeed, EEG signals change dramatically during sleep and show a transition from faster frequencies to increasingly slower frequencies such as alpha waves. In fact, different sleep stages are commonly characterized by their spectral content.^[14] Consequently, neural oscillations have been linked to cognitive states, such as awareness and consciousness.^{[15][16][13][11]}



Five frequency bands of neural oscillation, as seen in ten seconds of EEG.

Although neural oscillations in human brain activity are mostly investigated using EEG recordings, they are also observed using more invasive recording techniques such as single-unit recordings. Neurons can generate rhythmic patterns of action potentials or spikes. Some types of neurons have the tendency to fire at particular frequencies, so-called *resonators*.^[17] Bursting is another form of rhythmic spiking. Spiking patterns are considered fundamental for information coding in the brain. Oscillatory activity can also be observed in the form of subthreshold membrane potential oscillations (i.e. in the absence of action potentials).^[18] If numerous neurons spike in synchrony, they can give rise to oscillations in local field potentials. Quantitative models can estimate the strength of neural oscillations in recorded data.^[19]

Neural oscillations are commonly studied from a mathematical framework and belong to the field of "neurodynamics", an area of research in the cognitive sciences that places a strong focus upon the dynamic character of neural activity in describing brain function.^[20] It considers the brain a dynamical system and uses differential equations to describe how neural activity evolves over time. In particular, it aims to relate dynamic patterns of brain activity to cognitive functions such as perception and memory. In very abstract form, neural oscillations can be analyzed analytically. When studied in a more physiologically realistic setting, oscillatory activity is generally studied using computer simulations of a computational model.


The functions of neural oscillations are wide-ranging and vary for different types of oscillatory activity. Examples are the generation of rhythmic activity such as a heartbeat and the neural binding of sensory features in perception, such as the shape and color of an object. Neural oscillations also play an important role in many neurological disorders, such as excessive synchronization during seizure activity in epilepsy or tremor in patients with Parkinson's disease. Oscillatory activity can also be used to control external devices such as a brain-computer interface.^[21]

Physiology

Oscillatory activity is observed throughout the central nervous system at all levels of organization. Three different levels have been widely recognized: the micro-scale (activity of a single neuron), the meso-scale (activity of a local group of neurons) and the macro-scale (activity of different brain regions).^[22]

Microscopic

Neurons generate action potentials resulting from changes in the electric membrane potential. Neurons can generate multiple action potentials in sequence forming so-called spike trains. These spike trains are the basis for neural coding and information transfer in the brain. Spike trains can form all kinds of patterns, such as rhythmic spiking and bursting, and often display oscillatory activity.^[23] Oscillatory activity in single neurons can also be observed in sub-threshold fluctuations in membrane potential. These rhythmic changes in membrane potential do not reach the critical threshold and therefore do not result in an action potential. They can result from postsynaptic potentials from synchronous inputs or from intrinsic properties of neurons.



Tonic firing pattern of single neuron showing rhythmic spiking activity

Neuronal spiking can be classified by their activity patterns. The excitability of neurons can be subdivided in Class I and II. Class I neurons can generate action potentials with arbitrarily low frequency depending on the input strength, whereas Class II neurons generate action potentials in a certain frequency band, which is relatively insensitive to changes in input strength.^[17] Class II neurons are also more prone to display sub-threshold oscillations in membrane potential.

Mesosopic

A group of neurons can also generate oscillatory activity. Through synaptic interactions, the firing patterns of different neurons may become synchronized and the rhythmic changes in electric potential caused by their action potentials will add up (constructive interference). That is, synchronized firing patterns result in synchronized input into other cortical areas, which gives rise to large-amplitude oscillations of the local field potential. These large-scale oscillations can also be measured outside the scalp using electroencephalography (EEG) and magnetoencephalography (MEG). The electric potentials generated by single neurons are far too small to be picked up outside the scalp, and EEG or MEG

activity always reflects the summation of the synchronous activity of thousands or millions of neurons that have similar spatial orientation.^[24] Neurons in a neural ensemble rarely all fire at exactly the same moment, i.e. fully synchronized. Instead, the probability of firing is rhythmically modulated such that neurons are more likely to fire at the same time, which gives rise to oscillations in their mean activity (see figure at top of page). As such, the frequency of large-scale oscillations does not need to match the firing pattern of individual neurons. Isolated cortical neurons fire regularly under certain conditions, but in the intact brain cortical cells are bombarded by highly fluctuating synaptic inputs and typically fire seemingly at random. However, if the probability of a large group of neurons is rhythmically modulated at a common frequency, they will generate oscillations in the mean field (see also figure at top of page).^[23] Neural ensembles can generate oscillatory activity endogenously through local interactions between excitatory and inhibitory neurons. In particular, inhibitory interneurons play an important role in producing neural ensemble synchrony by generating a narrow window for effective excitation and rhythmically modulating the firing rate of excitatory neurons.^[25]

Macroscopic

Neural oscillation can also arise from interactions between different brain areas coupled through the structural connectome. Time delays play an important role here. Because all brain areas are bidirectionally coupled, these connections between brain areas form feedback loops. Positive feedback loops tend to cause oscillatory activity where frequency is inversely related to the delay time. An example of such a feedback loop is the connections between the thalamus and cortex – the thalamocortical radiations. This thalamocortical network is able to generate oscillatory activity known as recurrent thalamo-cortical resonance.^[26] The thalamocortical network plays an important role in the generation of alpha activity.^{[27][28]} In a whole-brain network model with realistic anatomical connectivity and propagation delays between brain areas, oscillations in the beta frequency range emerge from the partial synchronisation of subsets of brain areas oscillating in the gamma-band (generated at the mesoscopic level).^[29]

Mechanisms

Neuronal properties

Scientists have identified some intrinsic neuronal properties that play an important role in generating membrane potential oscillations. In particular, voltage-gated ion channels are critical in the generation of action potentials. The dynamics of these ion channels have been captured in the well-established Hodgkin-Huxley model that describes how action potentials are initiated and propagated by means of a set of differential equations. Using bifurcation analysis, different oscillatory varieties of these neuronal models can be determined, allowing for the classification of types of neuronal responses. The oscillatory dynamics of neuronal spiking as identified in the Hodgkin-Huxley model closely agree with empirical findings. In addition to periodic spiking, subthreshold membrane potential oscillations, i.e. resonance behavior that does not result in action potentials, may also contribute to oscillatory activity by facilitating synchronous activity of

neighboring neurons.^{[30][31]} Like pacemaker neurons in central pattern generators, subtypes of cortical cells fire bursts of spikes (brief clusters of spikes) rhythmically at preferred frequencies. Bursting neurons have the potential to serve as pacemakers for synchronous network oscillations, and bursts of spikes may underlie or enhance neuronal resonance.^[23]

Network properties

Apart from intrinsic properties of neurons, biological neural network properties are also an important source of oscillatory activity. Neurons communicate with one another via synapses and affect the timing of spike trains in the post-synaptic neurons. Depending on the properties of the connection, such as the coupling strength, time delay and whether coupling is excitatory or inhibitory, the spike trains of the interacting neurons may become synchronized.^[32] Neurons are locally connected, forming small clusters that are called neural ensembles. Certain network structures promote oscillatory activity at specific frequencies. For example, neuronal activity generated by two populations of interconnected *inhibitory* and *excitatory* cells can show spontaneous oscillations that are described by the Wilson-Cowan model.

If a group of neurons engages in synchronized oscillatory activity, the neural ensemble can be mathematically represented as a single oscillator.^[22] Different neural ensembles are coupled through long-range connections and form a network of weakly coupled oscillators at the next spatial scale. Weakly coupled oscillators can generate a range of dynamics including oscillatory activity.^[33] Long-range connections between different brain structures, such as the thalamus and the cortex (see thalamocortical oscillation), involve time-delays due to the finite conduction velocity of axons. Because most connections are reciprocal, they form feed-back loops that support oscillatory activity. Oscillations recorded from multiple cortical areas can become synchronized to form large scale brain networks, whose dynamics and functional connectivity can be studied by means of spectral analysis and Granger causality measures.^[34] Coherent activity of large-scale brain activity may form dynamic links between brain areas required for the integration of distributed information.^[16]

Neuromodulation

In addition to fast direct synaptic interactions between neurons forming a network, oscillatory activity is regulated by neuromodulators on a much slower time scale. That is, the concentration levels of certain neurotransmitters are known to regulate the amount of oscillatory activity. For instance, GABA concentration has been shown to be positively correlated with frequency of oscillations in induced stimuli.^[35] A number of nuclei in the brainstem have diffuse projections throughout the brain influencing concentration levels of neurotransmitters such as norepinephrine, acetylcholine and serotonin. These neurotransmitter systems affect the physiological state, e.g., wakefulness or arousal, and have a pronounced effect on amplitude of different brain waves, such as alpha activity.^[36]

Mathematical description


Oscillations can often be described and analyzed using mathematics. Mathematicians have identified several dynamical mechanisms that generate rhythmicity. Among the most important are harmonic (linear) oscillators, limit cycle oscillators, and delayed-feedback oscillators.^[37] Harmonic oscillations appear very frequently in nature—examples are sound waves, the motion of a pendulum, and vibrations of every sort. They generally arise when a physical system is perturbed by a small degree from a minimum-energy state, and are well understood mathematically. Noise-driven harmonic oscillators realistically simulate alpha rhythm in the waking EEG as well as slow waves and spindles in the sleep EEG. Successful EEG analysis algorithms were based on such models. Several other EEG components are better described by limit-cycle or delayed-feedback oscillations. Limit-cycle oscillations arise from physical systems that show large deviations from equilibrium, whereas delayed-feedback oscillations arise when components of a system affect each other after significant time delays. Limit-cycle oscillations can be complex but there are powerful mathematical tools for analyzing them; the mathematics of delayed-feedback oscillations is primitive in comparison. Linear oscillators and limit-cycle oscillators qualitatively differ in terms of how they respond to fluctuations in input. In a linear oscillator, the frequency is more or less constant but the amplitude can vary greatly. In a limit-cycle oscillator, the amplitude tends to be more or less constant but the frequency can vary greatly. A heartbeat is an example of a limit-cycle oscillation in that the frequency of beats varies widely, while each individual beat continues to pump about the same amount of blood.

Computational models adopt a variety of abstractions in order to describe complex oscillatory dynamics observed in brain activity. Many models are used in the field, each defined at a different level of abstraction and trying to model different aspects of neural systems. They range from models of the short-term behaviour of individual neurons, through models of how the dynamics of neural circuitry arise from interactions between individual neurons, to models of how behaviour can arise from abstract neural modules that represent complete subsystems.

Single neuron model

A model of a biological neuron is a mathematical description of the properties of nerve cells, or neurons, that is designed to accurately describe and predict its biological processes. The most successful and widely used model of neurons, the Hodgkin-Huxley model, is based on data from the squid giant axon. It is a set of nonlinear ordinary differential equations that approximates the electrical characteristics of a neuron, in particular the generation and propagation of action potentials. The model is very accurate and detailed and Hodgkin and Huxley received the 1963 Nobel Prize in physiology or medicine for this work.

The mathematics of the Hodgkin-Huxley model are quite complicated and several simplifications have been proposed, such as the FitzHugh-Nagumo model, the Hindmarsh-Rose model or the capacitor-switch model^[38] as an extension of the integrate-and-fire model. Such models only



Simulation of a Hindmarsh-Rose neuron showing typical bursting behavior: a fast rhythm generated by individual spikes and a slower rhythm generated by the bursts.


capture the basic neuronal dynamics, such as rhythmic spiking and bursting, but are more computationally efficient. This allows the simulation of a large number of interconnected neurons that form a neural network.

Spiking model

A neural network model describes a population of physically interconnected neurons or a group of disparate neurons whose inputs or signalling targets define a recognizable circuit. These models aim to describe how the dynamics of neural circuitry arise from interactions between individual neurons. Local interactions between neurons can result in the synchronization of spiking activity and form the basis of oscillatory activity. In particular, models of interacting pyramidal cells and inhibitory interneurons have been shown to generate brain rhythms such as gamma activity.^[39] Similarly, it was shown that simulations of neural networks with a phenomenological model for neuronal response failures can predict spontaneous broadband neural oscillations.^[40]

Neural mass model

Neural field models are another important tool in studying neural oscillations and are a mathematical framework describing evolution of variables such as mean firing rate in space and time. In modeling the activity of large numbers of neurons, the central idea is to take the density of neurons to the continuum limit, resulting in spatially continuous neural networks. Instead of modelling individual neurons, this approach approximates a group of neurons by its average properties and interactions. It is based on the mean field approach, an area of statistical physics that deals with large-scale systems. Models based on these principles have been used to provide mathematical descriptions of neural oscillations and EEG rhythms. They have for instance been used to investigate visual hallucinations.^[42]



Simulation of a neural mass model showing network spiking during the onset of a seizure.^[41] As the gain A is increased the network starts to oscillate at 3Hz.

Kuramoto model

The Kuramoto model of coupled phase oscillators^[43] is one of the most abstract and fundamental models used to investigate neural oscillations and synchronization. It captures the activity of a local system (e.g., a single neuron or neural ensemble) by its circular phase alone and hence ignores the amplitude of oscillations (amplitude is constant).^[44] Interactions amongst these oscillators are introduced by a simple algebraic form (such as a sine function) and collectively generate a dynamical pattern at the global scale. The Kuramoto model is widely used to study oscillatory brain activity


Simulation of Kuramoto model showing neural synchronization and oscillations in the mean field

and several extensions have been proposed that increase its neurobiological plausibility, for instance by incorporating topological properties of local cortical connectivity.^[45] In particular, it describes how the activity of a group of interacting neurons can become synchronized and generate large-scale oscillations. Simulations using the Kuramoto model with realistic long-range cortical connectivity and time-delayed interactions reveal the emergence of slow patterned fluctuations that reproduce resting-state BOLD functional maps, which can be measured using fMRI.^[46]


Activity patterns

Both single neurons and groups of neurons can generate oscillatory activity spontaneously. In addition, they may show oscillatory responses to perceptual input or motor output. Some types of neurons will fire rhythmically in the absence of any synaptic input. Likewise, brain-wide activity reveals oscillatory activity while subjects do not engage in any activity, so-called resting-state activity. These ongoing rhythms can change in different ways in response to perceptual input or motor output. Oscillatory activity may respond by increases or decreases in frequency and amplitude or show a temporary interruption, which is referred to as phase resetting. In addition, external activity may not interact with ongoing activity at all, resulting in an additive response.

Oscillatory responses



The frequency of ongoing oscillatory activity is increased between t1 and t2. The amplitude of ongoing oscillatory activity is increased between t1 and t2.



The phase of ongoing oscillatory activity is reset at t_1 .

Activity is linearly added to ongoing oscillatory activity between t_1 and t_2 .

Ongoing activity

Spontaneous activity is brain activity in the absence of an explicit task, such as sensory input or motor output, and hence also referred to as resting-state activity. It is opposed to induced activity, i.e. brain activity that is induced by sensory stimuli or motor responses. The term *ongoing brain activity* is used in electroencephalography and magnetoencephalography for those signal components that are not associated with the processing of a stimulus or the occurrence of specific other events, such as moving a body part, i.e. events that do not form evoked potentials/evoked fields, or induced activity. Spontaneous activity is usually considered to be noise if one is interested in stimulus processing; however, spontaneous activity is considered to play a crucial role during brain development, such as in network formation and synaptogenesis. Spontaneous activity may be informative regarding the current mental state of the person (e.g. wakefulness, alertness) and is often used in sleep research. Certain types of oscillatory activity, such as alpha waves, are part of spontaneous activity. Statistical analysis of power fluctuations of alpha activity reveals a bimodal distribution, i.e. a high- and low-amplitude mode, and hence shows that resting-state activity does not just reflect a noise process.^[47] In case of fMRI, spontaneous fluctuations in the blood-oxygen-level dependent (BOLD) signal reveal correlation patterns that are linked to resting states networks, such as the default network.^[48] The temporal evolution of resting state networks is correlated with fluctuations of oscillatory EEG activity in different frequency bands.^[49]

Ongoing brain activity may also have an important role in perception, as it may interact with activity related to incoming stimuli. Indeed, EEG studies suggest that visual perception is dependent on both the phase and amplitude of cortical oscillations. For instance, the amplitude and phase of alpha activity at the moment of visual stimulation predicts whether a weak stimulus will be perceived by the subject.^{[50][51][52]}

Frequency response

In response to input, a neuron or neuronal ensemble may change the frequency at

which it oscillates, thus changing the rate at which it spikes. Often, a neuron's firing rate depends on the summed activity it receives. Frequency changes are also commonly observed in central pattern generators and directly relate to the speed of motor activities, such as step frequency in walking. However, changes in *relative* oscillation frequency between different brain areas is not so common because the frequency of oscillatory activity is often related to the time delays between brain areas.

Amplitude response

Next to evoked activity, neural activity related to stimulus processing may result in induced activity. Induced activity refers to modulation in ongoing brain activity induced by processing of stimuli or movement preparation. Hence, they reflect an indirect response in contrast to evoked responses. A well-studied type of induced activity is amplitude change in oscillatory activity. For instance, gamma activity often increases during increased mental activity such as during object representation.^[53] Because induced responses may have different phases across measurements and therefore would cancel out during averaging, they can only be obtained using time-frequency analysis. Induced activity generally reflects the activity of numerous neurons: amplitude changes in oscillatory activity are thought to arise from the synchronization of neural activity, for instance by synchronization of spike timing or membrane potential fluctuations of individual neurons. Increases in oscillatory activity are therefore often referred to as event-related synchronization, while decreases are referred to as event-related desynchronization.^[54]

Phase resetting

Phase resetting occurs when input to a neuron or neuronal ensemble resets the phase of ongoing oscillations.^[55] It is very common in single neurons where spike timing is adjusted to neuronal input (a neuron may spike at a fixed delay in response to periodic input, which is referred to as phase locking^[17]) and may also occur in neuronal ensembles when the phases of their neurons are adjusted simultaneously. Phase resetting is fundamental for the synchronization of different neurons or different brain regions^{[16][33]} because the timing of spikes can become phase locked to the activity of other neurons.

Phase resetting also permits the study of evoked activity, a term used in electroencephalography and magnetoencephalography for responses in brain activity that are directly related to stimulus-related activity. Evoked potentials and event-related potentials are obtained from an electroencephalogram by stimulus-locked averaging, i.e. averaging different trials at fixed latencies around the presentation of a stimulus. As a consequence, those signal components that are the same in each single measurement are conserved and all others, i.e. ongoing or spontaneous activity, are averaged out. That is, event-related potentials only reflect oscillations in brain activity that are phase-locked to the stimulus or event. Evoked activity is often considered to be independent from ongoing brain activity, although this is an ongoing debate.^{[56][57]}

Asymmetric amplitude modulation

It has recently been proposed that even if phases are not aligned across trials, induced activity may still cause event-related potentials because ongoing brain oscillations may not be symmetric and thus amplitude modulations may result in a baseline shift that does not average out.^{[58][59]} This model implies that slow event-related responses, such as asymmetric alpha activity, could result from asymmetric brain oscillation amplitude modulations, such as an asymmetry of the intracellular currents that propagate forward and backward down the dendrites.^[60] Under this assumption, asymmetries in the dendritic current would cause asymmetries in oscillatory activity measured by EEG and MEG, since dendritic currents in pyramidal cells are generally thought to generate EEG and MEG signals that can be measured at the scalp.^[61]

Function

Neural synchronization can be modulated by task constraints, such as attention, and is thought to play a role in feature binding,^[62] neuronal communication,^[5] and motor coordination.^[7] Neuronal oscillations became a hot topic in neuroscience in the 1990s when the studies of the visual system of the brain by Gray, Singer and others appeared to support the neural binding hypothesis.^[63] According to this idea, synchronous oscillations in neuronal ensembles bind neurons representing different features of an object. For example, when a person looks at a tree, visual cortex neurons representing the tree trunk and those representing the branches of the same tree would oscillate in synchrony to form a single representation of the tree. This phenomenon is best seen in local field potentials which reflect the synchronous activity of local groups of neurons, but has also been shown in EEG and MEG recordings providing increasing evidence for a close relation between synchronous oscillatory activity and a variety of cognitive functions such as perceptual grouping^[62] and attentional top-down control.^{[13][11][10]}

Pacemaker

Cells in the sinoatrial node, located in the right atrium of the heart, spontaneously depolarize approximately 100 times per minute. Although all of the heart's cells have the ability to generate action potentials that trigger cardiac contraction, the sinoatrial node normally initiates it, simply because it generates impulses slightly faster than the other areas. Hence, these cells generate the normal sinus rhythm and are called pacemaker cells as they directly control the heart rate. In the absence of extrinsic neural and hormonal control, cells in the SA node will rhythmically discharge. The sinoatrial node is richly innervated by the autonomic nervous system, which up or down regulates the spontaneous firing frequency of the pacemaker cells.

Central pattern generator

Synchronized firing of neurons also forms the basis of periodic motor commands for rhythmic movements. These rhythmic outputs are produced by a group of interacting neurons that form a network, called a central pattern generator. Central pattern generators are neuronal circuits that—when activated—can produce rhythmic motor patterns in the absence of sensory or descending inputs that carry specific timing information. Examples are walking, breathing, and

swimming,^[64] Most evidence for central pattern generators comes from lower animals, such as the lamprey, but there is also evidence for spinal central pattern generators in humans.^{[65][66]}

Information processing

Neuronal spiking is generally considered the basis for information transfer in the brain. For such a transfer, information needs to be coded in a spiking pattern. Different types of coding schemes have been proposed, such as rate coding and temporal coding. Neural oscillations could create periodic time windows in which input spikes have larger effect on neurons, thereby providing a mechanism for decoding temporal codes.^[67]

Perception

Synchronization of neuronal firing may serve as a means to group spatially segregated neurons that respond to the same stimulus in order to bind these responses for further joint processing, i.e. to exploit temporal synchrony to encode relations. Purely theoretical formulations of the binding-by-synchrony hypothesis were proposed first,^[68] but subsequently extensive experimental evidence has been reported supporting the potential role of synchrony as a relational code.^[69]

The functional role of synchronized oscillatory activity in the brain was mainly established in experiments performed on awake kittens with multiple electrodes implanted in the visual cortex. These experiments showed that groups of spatially segregated neurons engage in synchronous oscillatory activity when activated by visual stimuli. The frequency of these oscillations was in the range of 40 Hz and differed from the periodic activation induced by the grating, suggesting that the oscillations and their synchronization were due to internal neuronal interactions.^[69] Similar findings were shown in parallel by the group of Eckhorn, providing further evidence for the functional role of neural synchronization in feature binding.^[70] Since then, numerous studies have replicated these findings and extended them to different modalities such as EEG, providing extensive evidence of the functional role of gamma oscillations in visual perception.

Gilles Laurent and colleagues showed that oscillatory synchronization has an important functional role in odor perception. Perceiving different odors leads to different subsets of neurons firing on different sets of oscillatory cycles.^[71] These oscillations can be disrupted by GABA blocker picrotoxin,^[72] and the disruption of the oscillatory synchronization leads to impairment of behavioral discrimination of chemically similar odorants in bees^[73] and to more similar responses across odors in downstream β -lobe neurons.^[74] Recent follow-up of this work has shown that oscillations create periodic integration windows for Kenyon cells in the insect mushroom body, such that incoming spikes from the antennal lobe are more effective in activating Kenyon cells only at specific phases of the oscillatory cycle.^[67]

Neural oscillations are also thought to be involved in the sense of time^[75] and in somatosensory perception.^[76] However, recent findings argue against a clock-like function of cortical gamma oscillations.^[77]

Motor coordination

Oscillations have been commonly reported in the motor system. Pfurtscheller and colleagues found a reduction in alpha (8–12 Hz) and beta (13–30 Hz) oscillations in EEG activity when subjects made a movement.^{[54][78]} Using intra-cortical recordings, similar changes in oscillatory activity were found in the motor cortex when the monkeys performed motor acts that required significant attention.^{[79][80]} In addition, oscillations at spinal level become synchronised to beta oscillations in the motor cortex during constant muscle activation, as determined by cortico-muscular coherence.^{[81][82][83]} Likewise, muscle activity of different muscles reveals inter-muscular coherence at multiple distinct frequencies reflecting the underlying neural circuitry involved in motor coordination.^{[84][85]}

Recently it was found that cortical oscillations propagate as travelling waves across the surface of the motor cortex along dominant spatial axes characteristic of the local circuitry of the motor cortex.^[86] It has been proposed that motor commands in the form of travelling waves can be spatially filtered by the descending fibres to selectively control muscle force.^[87] Simulations have shown that ongoing wave activity in cortex can elicit steady muscle force with physiological levels of EEG-EMG coherence.^[88]

Oscillatory rhythms at 10 Hz have been recorded in a brain area called the inferior olive, which is associated with the cerebellum.^[18] These oscillations are also observed in motor output of physiological tremor^[89] and when performing slow finger movements.^[90] These findings may indicate that the human brain controls continuous movements intermittently. In support, it was shown that these movement discontinuities are directly correlated to oscillatory activity in a cerebello-thalamo-cortical loop, which may represent a neural mechanism for the intermittent motor control.^[91]

Memory

Neural oscillations, in particular theta activity, are extensively linked to memory function. Theta rhythms are very strong in rodent hippocampi and entorhinal cortex during learning and memory retrieval, and they are believed to be vital to the induction of long-term potentiation, a potential cellular mechanism for learning and memory. Coupling between theta and gamma activity is thought to be vital for memory functions, including episodic memory.^{[92][93]} Tight coordination of single-neuron spikes with local theta oscillations is linked to successful memory formation in humans, as more stereotyped spiking predicts better memory.^[94]

Sleep and consciousness

Sleep is a naturally recurring state characterized by reduced or absent consciousness and proceeds in cycles of rapid eye movement (REM) and non-rapid eye movement (NREM) sleep. Sleep stages are characterized by spectral content of EEG: for instance, stage N1 refers to the transition of the brain from alpha waves (common in the awake state) to theta waves, whereas stage N3 (deep or slow-wave sleep) is characterized by the presence of delta waves.^[95] The normal order of sleep stages is N1 → N2 → N3 → N2 → REM.

Development

Neural oscillations may play a role in neural development. For example, retinal waves are thought to have properties that define early connectivity of circuits and synapses between cells in the retina.^[96]

Pathology

Specific types of neural oscillations may also appear in pathological situations, such as Parkinson's disease or epilepsy. These pathological oscillations often consist of an aberrant version of a normal oscillation. For example, one of the best known types is the spike and wave oscillation, which is typical of generalized or absence epileptic seizures, and which resembles normal sleep spindle oscillations.

Catherine Metzger
13 Octobre 1869

Handwriting of a person affected by Parkinson's disease showing rhythmic tremor activity in the strokes

Tremor

A tremor is an involuntary, somewhat rhythmic, muscle contraction and relaxation involving to-and-fro movements of one or more body parts. It is the most common of all involuntary movements and can affect the hands, arms, eyes, face, head, vocal cords, trunk, and legs. Most tremors occur in the hands. In some people, tremor is a symptom of another neurological disorder. Many different forms of tremor have been identified, such as essential tremor or Parkinsonian tremor. It is argued that tremors are likely to be multifactorial in origin, with contributions from neural oscillations in the central nervous systems, but also from peripheral mechanisms such as reflex loop resonances.^[97]



Generalized 3 Hz spike and wave discharges reflecting seizure activity

Epilepsy

Epilepsy is a common chronic neurological disorder characterized by seizures. These seizures are transient signs and/or symptoms of abnormal, excessive or hypersynchronous neuronal activity in the brain.^[98]

Thalamocortical dysrhythmia

In thalamocortical dysrhythmia (TCD), normal thalamocortical resonance is disrupted. The thalamic loss of input allows the frequency of the thalamo-cortical column to slow into the theta or delta band as identified by MEG and EEG by machine learning.^[99] TCD can be treated with neurosurgical methods like thalamotomy.

Applications

Clinical endpoints

Neural oscillations are sensitive to several drugs influencing brain activity; accordingly, biomarkers based on neural oscillations are emerging as secondary endpoints in clinical trials and in quantifying effects in pre-clinical studies. These biomarkers are often named "EEG biomarkers" or "Neurophysiological Biomarkers" and are quantified using quantitative electroencephalography (qEEG). EEG biomarkers can be extracted from the EEG using the open-source Neurophysiological Biomarker Toolbox.

Brain-computer interface

Neural oscillation has been applied as a control signal in various brain-computer interfaces (BCIs).^[100] For example, a non-invasive BCI can be created by placing electrodes on the scalp and then measuring the weak electric signals. Although individual neuron activities cannot be recorded through non-invasive BCI because the skull damps and blurs the electromagnetic signals, oscillatory activity can still be reliably detected. The BCI was introduced by Vidal in 1973^[101] as challenge of using EEG signals to control objects outside human body.

After the BCI challenge, in 1988, alpha rhythm was used in a brain rhythm based BCI for control of a physical object, a robot.^{[102][103]} Alpha rhythm based BCI was the first BCI for control of a robot.^{[104][105]} In particular, some forms of BCI allow users to control a device by measuring the amplitude of oscillatory activity in specific frequency bands, including mu and beta rhythms.

Examples

A non-inclusive list of types of oscillatory activity found in the central nervous system:

- Alpha wave
- Beta wave
- Bursting
- Cardiac cycle
- Delta wave
- Epileptic seizure
- Gamma wave
- Mathematical modeling of electrophysiological activity in epilepsy
- Mu wave
- PGO waves
- Thalamocortical oscillations
- Sharp wave-ripple complexes
- Sleep spindle
- Subthreshold membrane potential oscillations

- [Theta wave](#)

See also

- [Cybernetics](#)
- [Dynamical systems theory](#)
- [EEG analysis](#)
- [Neurocybernetics](#)
- [Oscillatory neural network](#)
- [Systems neuroscience](#)
- [ThetaHealing](#)
- [Phase resetting in neurons](#)

References

1. Llinás RR (2014). "Intrinsic electrical properties of mammalian neurons and CNS function: a historical perspective" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4219458>). *Frontiers in Cellular Neuroscience*. **8**: 320. doi:10.3389/fncel.2014.00320 (<https://doi.org/10.3389%2Ffncel.2014.00320>). PMC 4219458 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4219458>). PMID 25408634 (<https://pubmed.ncbi.nlm.nih.gov/25408634>).
2. "Caton, Richard - The electric currents of the brain" (<http://echo.mpiwg-berlin.mpg.de/ECHODOCUVIEW?url=/permanent/vlp/lit27690/index.meta&ww=0.7143&wh=0.7143&wx=0.2632>). *echo.mpiwg-berlin.mpg.de*. Retrieved 2018-12-21.
3. Coenen A, Fine E, Zayachkivska O (2014). "Adolf Beck: a forgotten pioneer in electroencephalography". *Journal of the History of the Neurosciences*. **23** (3): 276–86. doi:10.1080/0964704x.2013.867600 (<https://doi.org/10.1080%2F0964704x.2013.867600>). PMID 24735457 (<https://pubmed.ncbi.nlm.nih.gov/24735457>) . S2CID 205664545 (<https://api.semanticscholar.org/CorpusID:205664545>).
4. Pravdich-Neminsky VV (1913). "Ein Versuch der Registrierung der elektrischen Gehirnerscheinungen". *Zentralblatt für Physiologie*. **27**: 951–60.
5. Fries P (October 2005). "A mechanism for cognitive dynamics: neuronal communication through neuronal coherence". *Trends in Cognitive Sciences*. **9** (10): 474–80. doi:10.1016/j.tics.2005.08.011 (<https://doi.org/10.1016%2Fj.tics.2005.08.011>). PMID 16150631 (<https://pubmed.ncbi.nlm.nih.gov/16150631>). S2CID 6275292 (<https://api.semanticscholar.org/CorpusID:6275292>).
6. Fell J, Axmacher N (February 2011). "The role of phase synchronization in memory processes". *Nature Reviews. Neuroscience*. **12** (2): 105–18. doi:10.1038/nrn2979 (<https://doi.org/10.1038%2Fnrn2979>). PMID 21248789 (<https://pubmed.ncbi.nlm.nih.gov/21248789>). S2CID 7422401 (<https://api.semanticscholar.org/CorpusID:7422401>).
7. Schnitzler A, Gross J (April 2005). "Normal and pathological oscillatory communication in the brain". *Nature Reviews. Neuroscience*. **6** (4): 285–96. doi:10.1038/nrn1650 (<https://doi.org/10.1038%2Fnrn1650>). PMID 15803160 (<https://pubmed.ncbi.nlm.nih.gov/15803160>). S2CID 2749709 (<https://api.semanticscholar.org/CorpusID:2749709>).

8. Baldauf D, Desimone R (April 2014). "Neural mechanisms of object-based attention" (<https://www.science.org/lookup/doi/10.1126/science.1247003>). *Science*. **344** (6182): 424–7. Bibcode:2014Sci...344..424B (<https://ui.adsabs.harvard.edu/abs/2014Sci...344..424B>). doi:10.1126/science.1247003 (<https://doi.org/10.1126%2Fscience.1247003>). PMID 24763592 (<https://pubmed.ncbi.nlm.nih.gov/24763592>). S2CID 34728448 (<https://api.semanticscholar.org/CorpusID:34728448>).
9. Foster JJ, Sutterer DW, Serences JT, Vogel EK, Awh E (July 2017). "Alpha-Band Oscillations Enable Spatially and Temporally Resolved Tracking of Covert Spatial Attention" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5675530>). *Psychological Science*. **28** (7): 929–941. doi:10.1177/0956797617699167 (<https://doi.org/10.1177%2F0956797617699167>). PMC 5675530 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5675530>). PMID 28537480 (<https://pubmed.ncbi.nlm.nih.gov/28537480>).
10. de Vries IE, Marinato G, Baldauf D (August 2021). "Decoding object-based auditory attention from source-reconstructed MEG alpha oscillations" (<https://www.jneurosci.org/content/early/2021/08/24/JNEUROSCI.0583-21.2021>). *The Journal of Neuroscience*. **41** (41): 8603–8617. doi:10.1523/JNEUROSCI.0583-21.2021 (<https://doi.org/10.1523%2FJNEUROSCI.0583-21.2021>). PMC 8513695 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8513695>). PMID 34429378 (<https://pubmed.ncbi.nlm.nih.gov/34429378>).
11. Bagherzadeh Y, Baldauf D, Pantazis D, Desimone R (February 2020). "Alpha Synchrony and the Neurofeedback Control of Spatial Attention" (<https://doi.org/10.1016/j.neuron.2019.11.001>). *Neuron*. **105** (3): 577–587.e5. doi:10.1016/j.neuron.2019.11.001 (<https://doi.org/10.1016%2Fj.neuron.2019.11.001>). PMID 31812515 (<https://pubmed.ncbi.nlm.nih.gov/31812515>). S2CID 208614924 (<https://api.semanticscholar.org/CorpusID:208614924>).
12. Berger H, Gray CM (1929). "Über das Elektroenkephalogramm des Menschen". *Arch Psychiat Nervenkr*. **87**: 527–570. doi:10.1007/BF01797193 (<https://doi.org/10.1007%2FBF01797193>). hdl:11858/00-001M-0000-002A-5DE0-7 (<https://hdl.handle.net/11858%2F00-001M-0000-002A-5DE0-7>). S2CID 10835361 (<https://api.semanticscholar.org/CorpusID:10835361>).
13. Baldauf D, Desimone R (April 2014). "Neural mechanisms of object-based attention" (<https://www.science.org/doi/10.1126/science.1247003>). *Science*. **344** (6182): 424–7. Bibcode:2014Sci...344..424B (<https://ui.adsabs.harvard.edu/abs/2014Sci...344..424B>). doi:10.1126/science.1247003 (<https://doi.org/10.1126%2Fscience.1247003>). PMID 24763592 (<https://pubmed.ncbi.nlm.nih.gov/24763592>). S2CID 34728448 (<https://api.semanticscholar.org/CorpusID:34728448>).
14. Dement W, Kleitman N (November 1957). "Cyclic variations in EEG during sleep and their relation to eye movements, body motility, and dreaming". *Electroencephalography and Clinical Neurophysiology*. **9** (4): 673–90. doi:10.1016/0013-4694(57)90088-3 (<https://doi.org/10.1016%2F0013-4694%2857%2990088-3>). PMID 13480240 (<https://pubmed.ncbi.nlm.nih.gov/13480240>).
15. Engel AK, Singer W (January 2001). "Temporal binding and the neural correlates of sensory awareness". *Trends in Cognitive Sciences*. **5** (1): 16–25. doi:10.1016/S1364-6613(00)01568-0 (<https://doi.org/10.1016%2FS1364-6613%2800%2901568-0>). PMID 11164732 (<https://pubmed.ncbi.nlm.nih.gov/11164732>). S2CID 11922975 (<https://api.semanticscholar.org/CorpusID:11922975>).
16. Varela F, Lachaux JP, Rodriguez E, Martinerie J (April 2001). "The brainweb: phase synchronization and large-scale integration". *Nature Reviews. Neuroscience*. **2** (4): 229–39. doi:10.1038/35067550 (<https://doi.org/10.1038%2F35067550>). PMID 11283746 (<https://pubmed.ncbi.nlm.nih.gov/11283746>). S2CID 18651043 (<https://api.semanticscholar.org/CorpusID:18651043>).

17. Izhikevich EM (2007). *Dynamical systems in neuroscience*. Cambridge, Massachusetts: The MIT Press.
18. Llinás R, Yarom Y (July 1986). "Oscillatory properties of guinea-pig inferior olivary neurones and their pharmacological modulation: an in vitro study" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1182792>). *The Journal of Physiology*. **376**: 163–82. doi:10.1113/jphysiol.1986.sp016147 (<https://doi.org/10.1113%2Fjphysiol.1986.sp016147>). PMC 1182792 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1182792>). PMID 3795074 (<https://pubmed.ncbi.nlm.nih.gov/3795074>).
19. Mureşan RC, Jurjuţ OF, Moca VV, Singer W, Nikolić D (March 2008). "The oscillation score: an efficient method for estimating oscillation strength in neuronal activity". *Journal of Neurophysiology*. **99** (3): 1333–53. doi:10.1152/jn.00772.2007 (<https://doi.org/10.1152%2Fjn.00772.2007>). PMID 18160427 (<https://pubmed.ncbi.nlm.nih.gov/18160427>).
20. Burrow T (1943). "The neurodynamics of behavior. A phylobiological foreword". *Philosophy of Science*. **10** (4): 271–288. doi:10.1086/286819 (<https://doi.org/10.1086%2F286819>). S2CID 121438105 (<https://api.semanticscholar.org/CorpusID:121438105>).
21. Vansteensel MJ, Pels EG, Bleichner MG, Branco MP, Denison T, Freudenburg ZV, et al. (November 2016). "Fully Implanted Brain-Computer Interface in a Locked-In Patient with ALS" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5326682>). *The New England Journal of Medicine*. **375** (21): 2060–2066. doi:10.1056/NEJMoa1608085 (<https://doi.org/10.1056%2FNEJMoa1608085>). hdl:1874/344360 (<https://hdl.handle.net/1874%2F344360>). PMC 5326682 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5326682>). PMID 27959736 (<https://pubmed.ncbi.nlm.nih.gov/27959736>).
22. Haken H (1996). *Principles of brain functioning*. Springer. ISBN 978-3-540-58967-9.
23. Wang XJ (July 2010). "Neurophysiological and computational principles of cortical rhythms in cognition" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2923921>). *Physiological Reviews*. **90** (3): 1195–268. doi:10.1152/physrev.00035.2008 (<https://doi.org/10.1152%2Fphysrev.00035.2008>). PMC 2923921 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2923921>). PMID 20664082 (<https://pubmed.ncbi.nlm.nih.gov/20664082>).
24. Nunez PL, Srinivasan R (1981). *Electric fields of the brain: The neurophysics of EEG* (<https://books.google.com/books?id=gu5qAAAAMAAJ>). Oxford University Press. ISBN 9780195027969.
25. Cardin JA, Carlén M, Meletis K, Knoblich U, Zhang F, Deisseroth K, et al. (June 2009). "Driving fast-spiking cells induces gamma rhythm and controls sensory responses" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3655711>). *Nature*. **459** (7247): 663–7. Bibcode:2009Natur.459..663C (<https://ui.adsabs.harvard.edu/abs/2009Natur.459..663C>). doi:10.1038/nature08002 (<https://doi.org/10.1038%2Fnature08002>). PMC 3655711 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3655711>). PMID 19396156 (<https://pubmed.ncbi.nlm.nih.gov/19396156>).
26. Llinás R, Ribary U, Contreras D, Pedroarena C (November 1998). "The neuronal basis for consciousness" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1692417>). *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*. **353** (1377): 1841–9. doi:10.1098/rstb.1998.0336 (<https://doi.org/10.1098%2Frstb.1998.0336>). PMC 1692417 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1692417>). PMID 9854256 (<https://pubmed.ncbi.nlm.nih.gov/9854256>).

27. Bollimunta A, Mo J, Schroeder CE, Ding M (March 2011). "Neuronal mechanisms and attentional modulation of corticothalamic α oscillations" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3505610>). *The Journal of Neuroscience*. **31** (13): 4935–43. doi:10.1523/JNEUROSCI.5580-10.2011 (<https://doi.org/10.1523%2FJNEUROSCI.5580-10.2011>). PMC 3505610 (<https://www.ncbi.nlm.nih.gov/pmc/article/s/PMC3505610>). PMID 21451032 (<https://pubmed.ncbi.nlm.nih.gov/21451032>).
28. Suffczynski P, Kalitzin S, Pfurtscheller G, Lopes da Silva FH (December 2001). "Computational model of thalamo-cortical networks: dynamical control of alpha rhythms in relation to focal attention". *International Journal of Psychophysiology*. **43** (1): 25–40. doi:10.1016/S0167-8760(01)00177-5 (<https://doi.org/10.1016%2FS0167-8760%2801%2900177-5>). PMID 11742683 (<https://pubmed.ncbi.nlm.nih.gov/11742683>).
29. Cabral J, Luckhoo H, Woolrich M, Joensson M, Mohseni H, Baker A, et al. (April 2014). "Exploring mechanisms of spontaneous functional connectivity in MEG: how delayed network interactions lead to structured amplitude envelopes of band-pass filtered oscillations" (<https://doi.org/10.1016%2Fj.neuroimage.2013.11.047>). *NeuroImage*. **90**: 423–35. doi:10.1016/j.neuroimage.2013.11.047 (<https://doi.org/10.1016%2Fj.neuroimage.2013.11.047>). PMID 24321555 (<https://pubmed.ncbi.nlm.nih.gov/24321555>).
30. Llinás RR (December 1988). "The intrinsic electrophysiological properties of mammalian neurons: insights into central nervous system function". *Science*. **242** (4886): 1654–64. Bibcode:1988Sci...242.1654L (<https://ui.adsabs.harvard.edu/abs/1988Sci...242.1654L>). doi:10.1126/science.3059497 (<https://doi.org/10.1126%2Fscience.3059497>). PMID 3059497 (<https://pubmed.ncbi.nlm.nih.gov/3059497>).
31. Llinás RR, Grace AA, Yarom Y (February 1991). "In vitro neurons in mammalian cortical layer 4 exhibit intrinsic oscillatory activity in the 10- to 50-Hz frequency range" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC50921>). *Proceedings of the National Academy of Sciences of the United States of America*. **88** (3): 897–901. Bibcode:1991PNAS...88..897L (<https://ui.adsabs.harvard.edu/abs/1991PNAS...88..897L>). doi:10.1073/pnas.88.3.897 (<https://doi.org/10.1073%2Fpnas.88.3.897>). PMC 50921 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC50921>). PMID 1992481 (<https://pubmed.ncbi.nlm.nih.gov/1992481>).
32. Zeitler M, Daffertshofer A, Gielen CC (June 2009). "Asymmetry in pulse-coupled oscillators with delay" (<https://research.vu.nl/ws/files/2548720/229630.pdf>) (PDF). *Physical Review E*. **79** (6 Pt 2): 065203. Bibcode:2009PhRvE..79f5203Z (<https://ui.adsabs.harvard.edu/abs/2009PhRvE..79f5203Z>). doi:10.1103/PhysRevE.79.065203 (<https://doi.org/10.1103%2FPhysRevE.79.065203>). hdl:1871/29169 (<https://hdl.handle.net/1871%2F29169>). PMID 19658549 (<https://pubmed.ncbi.nlm.nih.gov/19658549>).
33. Pikovsky A, Rosenblum M, Kurths J (2001). *Synchronization: a universal concept in nonlinear sciences*. Cambridge University Press. ISBN 978-0-521-53352-2.
34. Andrea Brovelli, Steven L. Bressler and their colleagues, 2004 (<http://www.pnas.org/cgi/reprint/101/26/9849.pdf>)
35. Muthukumaraswamy SD, Edden RA, Jones DK, Swettenham JB, Singh KD (May 2009). "Resting GABA concentration predicts peak gamma frequency and fMRI amplitude in response to visual stimulation in humans" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2688873>). *Proceedings of the National Academy of Sciences of the United States of America*. **106** (20): 8356–61. Bibcode:2009PNAS..106.8356M (<https://ui.adsabs.harvard.edu/abs/2009PNAS..106.8356M>). doi:10.1073/pnas.0900728106 (<https://doi.org/10.1073%2Fpnas.0900728106>). PMC 2688873 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2688873>). PMID 19416820 (<https://pubmed.ncbi.nlm.nih.gov/19416820>).

36. Moruzzi G, Magoun HW (November 1949). "Brain stem reticular formation and activation of the EEG". *Electroencephalography and Clinical Neurophysiology*. **1** (4): 455–73. doi:10.1016/0013-4694(49)90219-9 (<https://doi.org/10.1016%2F0013-4694%2849%2990219-9>). PMID 18421835 (<https://pubmed.ncbi.nlm.nih.gov/18421835>).
37. Buzsáki G, Draguhn A (June 2004). "Neuronal oscillations in cortical networks". *Science*. **304** (5679): 1926–9. Bibcode:2004Sci...304.1926B (<https://ui.adsabs.harvard.edu/abs/2004Sci...304.1926B>). doi:10.1126/science.1099745 (<https://doi.org/10.1126%2Fscience.1099745>). PMID 15218136 (<https://pubmed.ncbi.nlm.nih.gov/15218136>). S2CID 8002293 (<https://api.semanticscholar.org/CorpusID:8002293>).
38. Cejnar P, Vyšata O, Kukal J, Beránek M, Vališ M, Procházka A (April 2020). "Simple capacitor-switch model of excitatory and inhibitory neuron with all parts biologically explained allows input fire pattern dependent chaotic oscillations" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7192907>). *Scientific Reports*. **10** (1): 7353. Bibcode:2020NatSR..10.7353C (<https://ui.adsabs.harvard.edu/abs/2020NatSR..10.7353C>). doi:10.1038/s41598-020-63834-7 (<https://doi.org/10.1038%2Fs41598-020-63834-7>). PMC 7192907 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7192907>). PMID 32355185 (<https://pubmed.ncbi.nlm.nih.gov/32355185>).
39. Whittington MA, Traub RD, Kopell N, Ermentrout B, Buhl EH (December 2000). "Inhibition-based rhythms: experimental and mathematical observations on network dynamics". *International Journal of Psychophysiology*. **38** (3): 315–36. CiteSeerX 10.1.1.16.6410 (<https://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.16.6410>). doi:10.1016/S0167-8760(00)00173-2 (<https://doi.org/10.1016%2FS0167-8760%2800%2900173-2>). PMID 11102670 (<https://pubmed.ncbi.nlm.nih.gov/11102670>).
40. Goldental A, Vardi R, Sardi S, Sabo P, Kanter I (2015). "Broadband macroscopic cortical oscillations emerge from intrinsic neuronal response failures" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4626558>). *Frontiers in Neural Circuits*. **9**: 65. arXiv:1511.00235 (<https://arxiv.org/abs/1511.00235>). doi:10.3389/fncir.2015.00065 (<https://doi.org/10.3389%2Ffncir.2015.00065>). PMC 4626558 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4626558>). PMID 26578893 (<https://pubmed.ncbi.nlm.nih.gov/26578893>).
41. Wendling F, Bellanger JJ, Bartolomei F, Chauvel P (October 2000). "Relevance of nonlinear lumped-parameter models in the analysis of depth-EEG epileptic signals". *Biological Cybernetics*. **83** (4): 367–78. doi:10.1007/s004220000160 (<https://doi.org/10.1007%2Fs004220000160>). PMID 11039701 (<https://pubmed.ncbi.nlm.nih.gov/11039701>). S2CID 8751526 (<https://api.semanticscholar.org/CorpusID:8751526>).
42. Bressloff PC, Cowan JD (2003) Spontaneous pattern formation in primary visual cortex. In: J Hogan, AR Krauskopf, M di Bernado, RE Wilson (Eds.), *Nonlinear dynamics and chaos: where do we go from here?*
43. Kuramoto Y (1984). *Chemical Oscillations, Waves, and Turbulence*. Dover Publications.
44. Ermentrout B (1994). "An introduction to neural oscillators". In F Ventriglia (ed.). *Neural Modeling and Neural Networks*. pp. 79–110.
45. Breakspear M, Heitmann S, Daffertshofer A (2010). "Generative models of cortical oscillations: neurobiological implications of the kuramoto model" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2995481>). *Frontiers in Human Neuroscience*. **4**: 190. doi:10.3389/fnhum.2010.00190 (<https://doi.org/10.3389%2Ffnhum.2010.00190>). PMC 2995481 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2995481>). PMID 21151358 (<https://pubmed.ncbi.nlm.nih.gov/21151358>).

46. Cabral J, Hugues E, Sporns O, Deco G (July 2011). "Role of local network oscillations in resting-state functional connectivity" (<https://zenodo.org/record/3421175>). *NeuroImage*. **57** (1): 130–139. doi:10.1016/j.neuroimage.2011.04.010 (<https://doi.org/10.1016%2Fj.neuroimage.2011.04.010>). PMID 21511044 (<https://pubmed.ncbi.nlm.nih.gov/21511044>). S2CID 13959959 (<https://api.semanticscholar.org/CorpusID:13959959>).
47. Freyer F, Aquino K, Robinson PA, Ritter P, Breakspear M (July 2009). "Bistability and non-Gaussian fluctuations in spontaneous cortical activity" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6665653>). *The Journal of Neuroscience*. **29** (26): 8512–24. doi:10.1523/JNEUROSCI.0754-09.2009 (<https://doi.org/10.1523%2FJNEUROSCI.0754-09.2009>). PMC 6665653 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6665653>). PMID 19571142 (<https://pubmed.ncbi.nlm.nih.gov/19571142>).
48. Fox MD, Raichle ME (September 2007). "Spontaneous fluctuations in brain activity observed with functional magnetic resonance imaging". *Nature Reviews. Neuroscience*. **8** (9): 700–11. doi:10.1038/nrn2201 (<https://doi.org/10.1038%2Fnrn2201>). PMID 17704812 (<https://pubmed.ncbi.nlm.nih.gov/17704812>). S2CID 15979590 (<https://api.semanticscholar.org/CorpusID:15979590>).
49. Laufs H, Krakow K, Sterzer P, Eger E, Beyerle A, Salek-Haddadi A, Kleinschmidt A (September 2003). "Electroencephalographic signatures of attentional and cognitive default modes in spontaneous brain activity fluctuations at rest" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC196925>). *Proceedings of the National Academy of Sciences of the United States of America*. **100** (19): 11053–8. Bibcode:2003PNAS..10011053L (<https://ui.adsabs.harvard.edu/abs/2003PNAS..10011053L>). doi:10.1073/pnas.1831638100 (<https://doi.org/10.1073%2Fpnas.1831638100>). PMC 196925 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC196925>). PMID 12958209 (<https://pubmed.ncbi.nlm.nih.gov/12958209>).
50. Mathewson KE, Gratton G, Fabiani M, Beck DM, Ro T (March 2009). "To see or not to see: prestimulus alpha phase predicts visual awareness" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2724892>). *The Journal of Neuroscience*. **29** (9): 2725–32. doi:10.1523/JNEUROSCI.3963-08.2009 (<https://doi.org/10.1523%2FJNEUROSCI.3963-08.2009>). PMC 2724892 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2724892>). PMID 19261866 (<https://pubmed.ncbi.nlm.nih.gov/19261866>).
51. Busch NA, Dubois J, VanRullen R (June 2009). "The phase of ongoing EEG oscillations predicts visual perception" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6665641>). *The Journal of Neuroscience*. **29** (24): 7869–76. doi:10.1523/jneurosci.0113-09.2009 (<https://doi.org/10.1523%2Fjneurosci.0113-09.2009>). PMC 6665641 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6665641>). PMID 19535598 (<https://pubmed.ncbi.nlm.nih.gov/19535598>).
52. van Dijk H, Schoffelen JM, Oostenveld R, Jensen O (February 2008). "Prestimulus oscillatory activity in the alpha band predicts visual discrimination ability" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6671447>). *The Journal of Neuroscience*. **28** (8): 1816–23. doi:10.1523/jneurosci.1853-07.2008 (<https://doi.org/10.1523%2Fjneurosci.1853-07.2008>). PMC 6671447 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6671447>). PMID 18287498 (<https://pubmed.ncbi.nlm.nih.gov/18287498>).
53. Tallon-Baudry C, Bertrand O (April 1999). "Oscillatory gamma activity in humans and its role in object representation". *Trends in Cognitive Sciences*. **3** (4): 151–162. doi:10.1016/S1364-6613(99)01299-1 (<https://doi.org/10.1016%2FS1364-6613%2899%2901299-1>). PMID 10322469 (<https://pubmed.ncbi.nlm.nih.gov/10322469>). S2CID 1308261 (<https://api.semanticscholar.org/CorpusID:1308261>).

54. Pfurtscheller G, Lopes da Silva FH (November 1999). "Event-related EEG/MEG synchronization and desynchronization: basic principles". *Clinical Neurophysiology*. **110** (11): 1842–57. doi:10.1016/S1388-2457(99)00141-8 ([http://doi.org/10.1016/S1388-2457\(99\)00141-8](http://doi.org/10.1016/S1388-2457(99)00141-8)). PMID 10576479 (<http://pubmed.ncbi.nlm.nih.gov/10576479>). S2CID 24756702 (<https://api.semanticscholar.org/CorpusID:24756702>).
55. Tass PA (2007). *Phase resetting in medicine and biology: stochastic modelling and data analysis*. Berlin Heidelberg: Springer-Verlag. ISBN 978-3-540-65697-5.
56. Makeig S, Westerfield M, Jung TP, Enghoff S, Townsend J, Courchesne E, Sejnowski TJ (January 2002). "Dynamic brain sources of visual evoked responses". *Science*. **295** (5555): 690–4. Bibcode:2002Sci...295..690M (<https://ui.adsabs.harvard.edu/abs/2002Sci...295..690M>). doi:10.1126/science.1066168 (<https://doi.org/10.1126/science.1066168>). PMID 11809976 (<https://pubmed.ncbi.nlm.nih.gov/11809976>). S2CID 15200185 (<https://api.semanticscholar.org/CorpusID:15200185>).
57. Mäkinen V, Tiitinen H, May P (February 2005). "Auditory event-related responses are generated independently of ongoing brain activity". *NeuroImage*. **24** (4): 961–8. doi:10.1016/j.neuroimage.2004.10.020 (<https://doi.org/10.1016/j.neuroimage.2004.10.020>). PMID 15670673 (<https://pubmed.ncbi.nlm.nih.gov/15670673>). S2CID 16210275 (<https://api.semanticscholar.org/CorpusID:16210275>).
58. Nikulin VV, Linkenkaer-Hansen K, Nolte G, Lemm S, Müller KR, Ilmoniemi RJ, Curio G (May 2007). "A novel mechanism for evoked responses in the human brain". *The European Journal of Neuroscience*. **25** (10): 3146–54. doi:10.1111/j.1460-9568.2007.05553.x (<https://doi.org/10.1111/j.1460-9568.2007.05553.x>). PMID 17561828 (<https://pubmed.ncbi.nlm.nih.gov/17561828>). S2CID 12113334 (<https://api.semanticscholar.org/CorpusID:12113334>).
59. Mazaheri A, Jensen O (July 2008). "Asymmetric amplitude modulations of brain oscillations generate slow evoked responses" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6670375>). *The Journal of Neuroscience*. **28** (31): 7781–7. doi:10.1523/JNEUROSCI.1631-08.2008 (<https://doi.org/10.1523/JNEUROSCI.1631-08.2008>). PMC 6670375 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6670375>). PMID 18667610 (<https://pubmed.ncbi.nlm.nih.gov/18667610>).
60. Mazaheri A, Jensen O (2008). "Rhythmic pulsing: linking ongoing brain activity with evoked responses" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2972683>). *Frontiers in Human Neuroscience*. **4**: 177. doi:10.3389/fnhum.2010.00177 (<https://doi.org/10.3389/fnhum.2010.00177>). PMC 2972683 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2972683>). PMID 21060804 (<https://pubmed.ncbi.nlm.nih.gov/21060804>).
61. Hamalainen M, Hari R, Ilmoniemi RJ, Knuutila J, Lounasmaa OV (1993). "Magnetoencephalography - Theory, instrumentation, and applications to noninvasive studies of the working human brain" (<https://aaltodoc.aalto.fi/handle/123456789/18757>). *Rev Mod Phys*. **65** (2): 413–497. Bibcode:1993RvMP...65..413H (<https://ui.adsabs.harvard.edu/abs/1993RvMP...65..413H>). doi:10.1103/RevModPhys.65.413 (<https://doi.org/10.1103/RevModPhys.65.413>).
62. Singer W (1993). "Synchronization of cortical activity and its putative role in information processing and learning". *Annual Review of Physiology*. **55**: 349–74. doi:10.1146/annurev.ph.55.030193.002025 (<https://doi.org/10.1146/annurev.ph.55.030193.002025>). PMID 8466179 (<https://pubmed.ncbi.nlm.nih.gov/8466179>).

63. Singer W, Gray CM (1995). "Visual feature integration and the temporal correlation hypothesis". *Annual Review of Neuroscience*. **18**: 555–86. CiteSeerX 10.1.1.308.6735 (<https://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.308.6735>). doi:10.1146/annurev.ne.18.030195.003011 (<https://doi.org/10.1146%2Fannurev.ne.18.030195.003011>). PMID 7605074 (<https://pubmed.ncbi.nlm.nih.gov/7605074>).
64. Marder E, Bucher D (November 2001). "Central pattern generators and the control of rhythmic movements". *Current Biology*. **11** (23): R986–96. doi:10.1016/S0960-9822(01)00581-4 (<https://doi.org/10.1016%2FS0960-9822%2801%2900581-4>). PMID 11728329 (<https://pubmed.ncbi.nlm.nih.gov/11728329>). S2CID 1294374 (<https://api.semanticscholar.org/CorpusID:1294374>).
65. Dimitrijevic MR, Gerasimenko Y, Pinter MM (November 1998). "Evidence for a spinal central pattern generator in humans". *Annals of the New York Academy of Sciences*. **860** (1): 360–76. Bibcode:1998NYASA.860..360D (<https://ui.adsabs.harvard.edu/abs/1998NYASA.860..360D>). doi:10.1111/j.1749-6632.1998.tb09062.x (<https://doi.org/10.1111%2Fj.1749-6632.1998.tb09062.x>). PMID 9928325 (<https://pubmed.ncbi.nlm.nih.gov/9928325>). S2CID 102514 (<https://api.semanticscholar.org/CorpusID:102514>).
66. Danner SM, Hofstoetter US, Freundl B, Binder H, Mayr W, Rattay F, Minassian K (March 2015). "Human spinal locomotor control is based on flexibly organized burst generators" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4408427>). *Brain*. **138** (Pt 3): 577–88. doi:10.1093/brain/awu372 (<https://doi.org/10.1093%2Fbrain%2Fawu372>). PMC 4408427 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4408427>). PMID 25582580 (<https://pubmed.ncbi.nlm.nih.gov/25582580>).
67. Gupta N, Singh SS, Stopfer M (December 2016). "Oscillatory integration windows in neurons" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5171764>). *Nature Communications*. **7**: 13808. Bibcode:2016NatCo...713808G (<https://ui.adsabs.harvard.edu/abs/2016NatCo...713808G>). doi:10.1038/ncomms13808 (<https://doi.org/10.1038%2Fncomms13808>). PMC 5171764 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5171764>). PMID 27976720 (<https://pubmed.ncbi.nlm.nih.gov/27976720>).
68. Milner PM (November 1974). "A model for visual shape recognition". *Psychological Review*. **81** (6): 521–35. doi:10.1037/h0037149 (<https://doi.org/10.1037%2Fh0037149>). PMID 4445414 (<https://pubmed.ncbi.nlm.nih.gov/4445414>).
69. Gray CM, König P, Engel AK, Singer W (March 1989). "Oscillatory responses in cat visual cortex exhibit inter-columnar synchronization which reflects global stimulus properties". *Nature*. **338** (6213): 334–7. Bibcode:1989Natur.338..334G (<https://ui.adsabs.harvard.edu/abs/1989Natur.338..334G>). doi:10.1038/338334a0 (<https://doi.org/10.1038%2F338334a0>). PMID 2922061 (<https://pubmed.ncbi.nlm.nih.gov/2922061>). S2CID 4281744 (<https://api.semanticscholar.org/CorpusID:4281744>).
70. Eckhorn R, Bauer R, Jordan W, Brosch M, Kruse W, Munk M, Reitboeck HJ (1988). "Coherent oscillations: a mechanism of feature linking in the visual cortex? Multiple electrode and correlation analyses in the cat". *Biological Cybernetics*. **60** (2): 121–30. doi:10.1007/BF00202899 (<https://doi.org/10.1007%2FBF00202899>). PMID 3228555 (<https://pubmed.ncbi.nlm.nih.gov/3228555>). S2CID 206771651 (<https://api.semanticscholar.org/CorpusID:206771651>).
71. Wehr M, Laurent G (November 1996). "Odour encoding by temporal sequences of firing in oscillating neural assemblies". *Nature*. **384** (6605): 162–6. Bibcode:1996Natur.384..162W (<https://ui.adsabs.harvard.edu/abs/1996Natur.384..162W>). doi:10.1038/384162a0 (<https://doi.org/10.1038%2F384162a0>). PMID 8906790 (<https://pubmed.ncbi.nlm.nih.gov/8906790>). S2CID 4286308 (<https://api.semanticscholar.org/CorpusID:4286308>).

72. MacLeod K, Laurent G (November 1996). "Distinct mechanisms for synchronization and temporal patterning of odor-encoding neural assemblies". *Science*. **274** (5289): 976–9. Bibcode:1996Sci...274..976M (<https://ui.adsabs.harvard.edu/abs/1996Sci...274..976M>). doi:10.1126/science.274.5289.976 (<https://doi.org/10.1126/science.274.5289.976>). PMID 8875938 (<https://pubmed.ncbi.nlm.nih.gov/8875938>). S2CID 10744144 (<https://api.semanticscholar.org/CorpusID:10744144>).
73. Stopfer M, Bhagavan S, Smith BH, Laurent G (November 1997). "Impaired odour discrimination on desynchronization of odour-encoding neural assemblies". *Nature*. **390** (6655): 70–4. Bibcode:1997Natur.390...70S (<https://ui.adsabs.harvard.edu/abs/1997Natur.390...70S>). doi:10.1038/36335 (<https://doi.org/10.1038/36335>). PMID 9363891 (<https://pubmed.ncbi.nlm.nih.gov/9363891>). S2CID 205024830 (<https://api.semanticscholar.org/CorpusID:205024830>).
74. MacLeod K, Bäckér A, Laurent G (October 1998). "Who reads temporal information contained across synchronized and oscillatory spike trains?". *Nature*. **395** (6703): 693–8. Bibcode:1998Natur.395..693M (<https://ui.adsabs.harvard.edu/abs/1998Natur.395..693M>). doi:10.1038/27201 (<https://doi.org/10.1038/27201>). PMID 9790189 (<https://pubmed.ncbi.nlm.nih.gov/9790189>). S2CID 4424801 (<https://api.semanticscholar.org/CorpusID:4424801>).
75. Buhusi CV, Meck WH (October 2005). "What makes us tick? Functional and neural mechanisms of interval timing". *Nature Reviews. Neuroscience*. **6** (10): 755–65. doi:10.1038/nrn1764 (<https://doi.org/10.1038/nrn1764>). PMID 16163383 (<https://pubmed.ncbi.nlm.nih.gov/16163383>). S2CID 29616055 (<https://api.semanticscholar.org/CorpusID:29616055>).
76. Ahissar E, Zacksenhouse M (2001). *Temporal and spatial coding in the rat vibrissal system*. *Prog Brain Res*. Progress in Brain Research. Vol. 130. pp. 75–87. doi:10.1016/S0079-6123(01)30007-9 ([https://doi.org/10.1016/S0079-6123\(01\)30007-9](https://doi.org/10.1016/S0079-6123(01)30007-9)). ISBN 9780444501103. PMID 11480290 (<https://pubmed.ncbi.nlm.nih.gov/11480290>).
77. Burns SP, Xing D, Shapley RM (June 2011). "Is gamma-band activity in the local field potential of V1 cortex a "clock" or filtered noise?" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3518456>). *The Journal of Neuroscience*. **31** (26): 9658–64. doi:10.1523/jneurosci.0660-11.2011 (<https://doi.org/10.1523/jneurosci.0660-11.2011>). PMC 3518456 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3518456>). PMID 21715631 (<https://pubmed.ncbi.nlm.nih.gov/21715631>).
78. Pfurtscheller G, Aranibar A (June 1977). "Event-related cortical desynchronization detected by power measurements of scalp EEG". *Electroencephalography and Clinical Neurophysiology*. **42** (6): 817–26. doi:10.1016/0013-4694(77)90235-8 ([https://doi.org/10.1016/0013-4694\(77\)90235-8](https://doi.org/10.1016/0013-4694(77)90235-8)). PMID 67933 (<https://pubmed.ncbi.nlm.nih.gov/67933>).
79. Murthy VN, Fetz EE (December 1996). "Oscillatory activity in sensorimotor cortex of awake monkeys: synchronization of local field potentials and relation to behavior". *Journal of Neurophysiology*. **76** (6): 3949–67. doi:10.1152/jn.1996.76.6.3949 (<https://doi.org/10.1152/jn.1996.76.6.3949>). PMID 8985892 (<https://pubmed.ncbi.nlm.nih.gov/8985892>).
80. Sanes JN, Donoghue JP (May 1993). "Oscillations in local field potentials of the primate motor cortex during voluntary movement" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC46533>). *Proceedings of the National Academy of Sciences of the United States of America*. **90** (10): 4470–4. Bibcode:1993PNAS...90.4470S (<https://ui.adsabs.harvard.edu/abs/1993PNAS...90.4470S>). doi:10.1073/pnas.90.10.4470 (<https://doi.org/10.1073/pnas.90.10.4470>). PMC 46533 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC46533>). PMID 8506287 (<https://pubmed.ncbi.nlm.nih.gov/8506287>).

81. Conway BA, Halliday DM, Farmer SF, Shahani U, Maas P, Weir AI, Rosenberg JR (December 1995). "Synchronization between motor cortex and spinal motoneuronal pool during the performance of a maintained motor task in man" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1156860>). *The Journal of Physiology*. 489 (Pt 3) (3): 917–24. doi:10.1113/jphysiol.1995.sp021104 (<https://doi.org/10.1113%2Fjphysiol.1995.sp021104>). PMC 1156860 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1156860>). PMID 8788955 (<https://pubmed.ncbi.nlm.nih.gov/8788955>).
82. Salenius S, Portin K, Kajola M, Salmelin R, Hari R (June 1997). "Cortical control of human motoneuron firing during isometric contraction". *Journal of Neurophysiology*. **77** (6): 3401–5. doi:10.1152/jn.1997.77.6.3401 (<https://doi.org/10.1152%2Fjn.1997.77.6.3401>). PMID 9212286 (<https://pubmed.ncbi.nlm.nih.gov/9212286>). S2CID 2178927 (<https://api.semanticscholar.org/CorpusID:2178927>)
.
83. Baker SN, Olivier E, Lemon RN (May 1997). "Coherent oscillations in monkey motor cortex and hand muscle EMG show task-dependent modulation" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1159515>). *The Journal of Physiology*. 501 (Pt 1) (1): 225–41. doi:10.1111/j.1469-7793.1997.225bo.x (<https://doi.org/10.1111%2Fj.1469-7793.1997.225bo.x>). PMC 1159515 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1159515>). PMID 9175005 (<https://pubmed.ncbi.nlm.nih.gov/9175005>).
84. Boonstra TW, Danna-Dos-Santos A, Xie HB, Roerdink M, Stins JF, Breakspear M (December 2015). "Muscle networks: Connectivity analysis of EMG activity during postural control" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4669476>). *Scientific Reports*. **5**: 17830. Bibcode:2015NatSR...517830B (<https://ui.adsabs.harvard.edu/abs/2015NatSR...517830B>). doi:10.1038/srep17830 (<https://doi.org/10.1038%2Fsrep17830>). PMC 4669476 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4669476>). PMID 26634293 (<https://pubmed.ncbi.nlm.nih.gov/26634293>)
.
85. Kerkman JN, Daffertshofer A, Gollo LL, Breakspear M, Boonstra TW (June 2018). "Network structure of the human musculoskeletal system shapes neural interactions on multiple time scales" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6021138>). *Science Advances*. **4** (6): eaat0497. Bibcode:2018SciA....4..497K (<https://ui.adsabs.harvard.edu/abs/2018SciA....4..497K>). doi:10.1126/sciadv.aat0497 (<https://doi.org/10.1126%2Fsciadv.aat0497>). PMC 6021138 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6021138>). PMID 29963631 (<https://pubmed.ncbi.nlm.nih.gov/29963631>).
86. Rubino D, Robbins KA, Hatsopoulos NG (December 2006). "Propagating waves mediate information transfer in the motor cortex". *Nature Neuroscience*. **9** (12): 1549–57. doi:10.1038/nn1802 (<https://doi.org/10.1038%2Fnn1802>). PMID 17115042 (<https://pubmed.ncbi.nlm.nih.gov/17115042>). S2CID 16430438 (<https://api.semanticscholar.org/CorpusID:16430438>).
87. Heitmann S, Boonstra T, Gong P, Breakspear M, Ermentrout B (2015). "The rhythms of steady posture: Motor commands as spatially organized oscillation patterns". *Neurocomputing*. **170**: 3–14. doi:10.1016/j.neucom.2015.01.088 (<https://doi.org/10.1016%2Fj.neucom.2015.01.088>).

88. Heitmann S, Boonstra T, Breakspear M (October 2013). "A dendritic mechanism for decoding traveling waves: principles and applications to motor cortex" (<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3814333>). *PLOS Computational Biology*. **9** (10): e1003260. Bibcode:2013PLSCB...9E3260H (<https://ui.adsabs.harvard.edu/abs/2013PLSCB...9E3260H>). doi:10.1371/journal.pcbi.1003260 (<https://doi.org/10.1371%2Fjournal.pcbi.1003260>). PMC 3814333 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3814333>). PMID 24204220 (<https://pubmed.ncbi.nlm.nih.gov/24204220>).
89. Allum JH, Dietz V, Freund HJ (May 1978). "Neuronal mechanisms underlying physiological tremor". *Journal of Neurophysiology*. **41** (3): 557–71. doi:10.1152/jn.1978.41.3.557 (<https://doi.org/10.1152%2Fjn.1978.41.3.557>). PMID 660226 (<https://pubmed.ncbi.nlm.nih.gov/660226>).
90. Vallbo AB, Wessberg J (September 1993). "Organization of motor output in slow finger movements in man" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1143894>). *The Journal of Physiology*. **469**: 673–91. doi:10.1113/jphysiol.1993.sp019837 (<https://doi.org/10.1113%2Fjphysiol.1993.sp019837>). PMC 1143894 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1143894>). PMID 8271223 (<https://pubmed.ncbi.nlm.nih.gov/8271223>).
91. Gross J, Timmermann L, Kujala J, Dirks M, Schmitz F, Salmelin R, Schnitzler A (February 2002). "The neural basis of intermittent motor control in humans" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC122359>). *Proceedings of the National Academy of Sciences of the United States of America*. **99** (4): 2299–302. Bibcode:2002PNAS...99.2299G (<https://ui.adsabs.harvard.edu/abs/2002PNAS...99.2299G>). doi:10.1073/pnas.032682099 (<https://doi.org/10.1073%2Fpnas.032682099>). PMC 122359 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC122359>). PMID 11854526 (<https://pubmed.ncbi.nlm.nih.gov/11854526>).
92. Buszaki G (2006). *Rhythms of the brain*. Oxford University Press.
93. Nyhus E, Curran T (June 2010). "Functional role of gamma and theta oscillations in episodic memory" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2856712>). *Neuroscience and Biobehavioral Reviews*. **34** (7): 1023–35. doi:10.1016/j.neubiorev.2009.12.014 (<https://doi.org/10.1016%2Fj.neubiorev.2009.12.014>). PMC 2856712 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2856712>). PMID 20060015 (<https://pubmed.ncbi.nlm.nih.gov/20060015>).
94. Rutishauser U, Ross IB, Mamelak AN, Schuman EM (April 2010). "Human memory strength is predicted by theta-frequency phase-locking of single neurons" (<http://authors.library.caltech.edu/18277/2/nature08860-s1.pdf>) (PDF). *Nature*. **464** (7290): 903–7. Bibcode:2010Natur.464..903R (<https://ui.adsabs.harvard.edu/abs/2010Natur.464..903R>). doi:10.1038/nature08860 (<https://doi.org/10.1038%2Fnature08860>). PMID 20336071 (<https://pubmed.ncbi.nlm.nih.gov/20336071>). S2CID 4417989 (<https://api.semanticscholar.org/CorpusID:4417989>).
95. Brancaccio A, Tabarelli D, Bigica M, Baldauf D (April 2020). "Cortical source localization of sleep-stage specific oscillatory activity" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7181624>). *Scientific Reports*. **10** (1): 6976. Bibcode:2020NatSR..10.6976B (<https://ui.adsabs.harvard.edu/abs/2020NatSR..10.6976B>). doi:10.1038/s41598-020-63933-5 (<https://doi.org/10.1038%2Fs41598-020-63933-5>). PMC 7181624 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7181624>). PMID 32332806 (<https://pubmed.ncbi.nlm.nih.gov/32332806>).
96. Feller MB (July 2009). "Retinal waves are likely to instruct the formation of eye-specific retinogeniculate projections" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2706239>). *Neural Development*. **4**: 24. doi:10.1186/1749-8104-4-24 (<https://doi.org/10.1186%2F1749-8104-4-24>). PMC 2706239 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2706239>). PMID 19580682 (<https://pubmed.ncbi.nlm.nih.gov/19580682>).

97. McAuley JH, Marsden CD (August 2000). "Physiological and pathological tremors and rhythmic central motor control" (<https://doi.org/10.1093%2Fbrain%2F123.8.1545>). *Brain*. 123 (Pt 8) (8): 1545–67. doi:10.1093/brain/123.8.1545 (<https://doi.org/10.1093%2Fbrain%2F123.8.1545>). PMID 10908186 (<https://pubmed.ncbi.nlm.nih.gov/10908186>).
98. Shusterman V, Troy WC (June 2008). "From baseline to epileptiform activity: a path to synchronized rhythmicity in large-scale neural networks". *Physical Review E*. **77** (6 Pt 1): 061911. Bibcode:2008PhRvE..77f1911S (<https://ui.adsabs.harvard.edu/abs/2008PhRvE..77f1911S>). doi:10.1103/PhysRevE.77.061911 (<https://doi.org/10.1103%2FPhysRevE.77.061911>). PMID 18643304 (<https://pubmed.ncbi.nlm.nih.gov/18643304>).
99. Vanneste S, Song JJ, De Ridder D (March 2018). "Thalamocortical dysrhythmia detected by machine learning" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5856824>). *Nature Communications*. **9** (1): 1103. Bibcode:2018NatCo...9.1103V (<https://ui.adsabs.harvard.edu/abs/2018NatCo...9.1103V>). doi:10.1038/s41467-018-02820-0 (<https://doi.org/10.1038%2Fs41467-018-02820-0>). PMC 5856824 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5856824>). PMID 29549239 (<https://pubmed.ncbi.nlm.nih.gov/29549239>).
100. Birbaumer N (November 2006). "Breaking the silence: brain-computer interfaces (BCI) for communication and motor control" (<https://doi.org/10.1111%2Fj.1469-8986.2006.00456.x>). *Psychophysiology*. **43** (6): 517–32. doi:10.1111/j.1469-8986.2006.00456.x (<https://doi.org/10.1111%2Fj.1469-8986.2006.00456.x>). PMID 17076808 (<https://pubmed.ncbi.nlm.nih.gov/17076808>).
101. Vidal JJ (1973). "Toward direct brain-computer communication". *Annual Review of Biophysics and Bioengineering*. **2**: 157–80. doi:10.1146/annurev.bb.02.060173.001105 (<https://doi.org/10.1146%2Fannurev.bb.02.060173.001105>). PMID 4583653 (<https://pubmed.ncbi.nlm.nih.gov/4583653>).
102. Bozinovski S, Sestakov M, Bozinovska L (November 1988). "Using EEG alpha rhythm to control a mobile robot.". *Proceedings of the Annual International Conference of the IEEE Engineering in Medicine and Biology Society*. New Orleans: IEEE. pp. 1515–1516. doi:10.1109/IEMBS.1988.95357 (<https://doi.org/10.1109%2FIEMBS.1988.95357>). ISBN 0-7803-0785-2. S2CID 62179588 (<https://api.semanticscholar.org/CorpusID:62179588>).
103. Bozinovski S (August 1990). "Mobile robot trajectory control: From fixed rails to direct bioelectric control.". In Kaynak O (ed.). *Proceedings of the IEEE International Workshop on Intelligent Motion Control*. Vol. 2. Istanbul: IEEE. pp. 463–467. doi:10.1109/IMC.1990.687362 (<https://doi.org/10.1109%2FIMC.1990.687362>). S2CID 60642344 (<https://api.semanticscholar.org/CorpusID:60642344>).
104. Lebedev M (2016). "Augmentation of sensorimotor functions with neural prostheses". *Opera Medica et Physiologica*. **2** (3–4): 211–227. doi:10.20388/OMP.003.0035 (<https://doi.org/10.20388%2FOMP.003.0035>) (inactive 31 July 2022).
105. Lebedev MA, Nicolelis MA (April 2017). "Brain-Machine Interfaces: From Basic Science to Neuroprostheses and Neurorehabilitation". *Physiological Reviews*. **97** (2): 767–837. doi:10.1152/physrev.00027.2016 (<https://doi.org/10.1152%2Fphysrev.00027.2016>). PMID 28275048 (<https://pubmed.ncbi.nlm.nih.gov/28275048>).

Further reading

- Buzsáki G (2006). *Rhythms of the Brain*. Oxford University Press.

ISBN 978-0-19-530106-9.

- Freeman W (1975). *Mass Action in the Nervous System* (<https://web.archive.org/web/20150705004813/http://sulcus.berkeley.edu/MANSWWW/MANSWWW.html>). Academic Press. ISBN 978-0124120471. Archived from the original (<http://sulcus.berkeley.edu/MANSWWW/MANSWWW.html>) on 2015-07-05.

External links

- Binding by synchronization (http://www.scholarpedia.org/article/Binding_by_synchrony)
 - Neural Field Theory (http://www.scholarpedia.org/article/Neural_fields)
 - Spike-and-wave oscillations (http://www.scholarpedia.org/article/Spike-and-wave_oscillations)
 - Synchronization (<http://www.scholarpedia.org/article/Synchronization>)
 - Bursting (<http://www.scholarpedia.org/article/Bursting>)
-

Retrieved from "https://en.wikipedia.org/w/index.php?title=Neural_oscillation&oldid=1101829819"

This page was last edited on 2 August 2022, at 01:21 (UTC).

Text is available under the Creative Commons Attribution-ShareAlike License 3.0; additional terms may apply. By using this site, you agree to the Terms of Use and Privacy Policy. Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.