

Natural Resource Inventory for Lyme, NH
APPENDIX III
DOCUMENTATION OF BIRDS IN TWO SECTIONS OF LYME
SouthEast (SE) 1967-2006 (Holmes)
NorthWest (NW) 1981-2006 (Skelly)

Bird Species of Southeastern & Northwestern Sections of Lyme, NH 1967-2006

Common Name	Scientific Name	Comments <u>primarily from</u> <u>Holmes in SE</u>	SE	NW
Bittern, American	<i>Botaurus lentiginosus</i>	breeds in swamp below lower edge of fields	x	
Blackbird, Red-winged	<i>Agelaius phoeniceus</i>	common breeder in swamp/wet areas	x	x
Blackbird, Rusty	<i>Euphagus carolinus</i>	spring and fall migrant	x	
Bluebird, Eastern	<i>Sialia sialis</i>	sporadic breeder, occurs in fields and nest boxes	x	x
Bobolink	<i>Dolichonyx oryzivorus</i>	common breeder in hayfields	x	x
Bunting, Indigo	<i>Passerina cyanea</i>	breeder along forest edge	x	x
Cardinal, Northern	<i>Cardinalis cardinalis</i>	occasional on feeder in fall and winter	x	x
Catbird, Gray	<i>Dumetella carolinensis</i>	common breeder	x	x
Chickadee, Black-capped	<i>Poecile atricapilla</i>	common breeder in forest, at feeders	x	x
Cowbird, Brown-headed	<i>Molothrus ater</i>	common breeder	x	x
Creeper, Brown	<i>Certhia americana</i>	breeder in forest, widely scattered	x	
Crossbill, Red	<i>Loxia curvirostra</i>	occasional winter visitor	x	x
Crossbill, White-winged	<i>Loxia leucoptera</i>	occasional winter visitor	x	x
Crow, American	<i>Corvus brachyrhynchos</i>	common in valley	x	x
Cuckoo, Black-billed	<i>Coccyzus erythrophthalmus</i>	irregular breeder	x	x
Cuckoo, Yellow-billed	<i>Coccyzus americanus</i>	occasional records in mid to late summer	x	x
Dickcissel	<i>Spiza americana</i>	one record in May	x	
Dove, Mourning	<i>Zenaida macroura</i>	common year round	x	x
Dove, Rock	<i>Columba livia</i>	used to nest in our silo, now gone	x	x
Duck, Black	<i>Anas rubripes</i>	seen flying along valley, on Roby's pond in '05	x	
Duck, Wood	<i>Aix sponsa</i>	flying along valley	x	x
Finch, House	<i>Carpodacus mexicanus</i>	regularly seen in late 70s when they bred, not seen recently.	x	
Finch, Purple	<i>Carpodacus purpureus</i>	seen mostly in early spring and fall	x	x
Flicker, Common	<i>Colaptes auratus</i>	common breeder and migrant	x	x
Flycatcher, Alder	<i>Empidonax alnorum</i>	fairly common breeder and caller		x
Flycatcher, Great Crested	<i>Myiarchus crinitus</i>	occasional breeder in forest	x	x
Flycatcher, Least	<i>Empidonax minimus</i>	formerly bred along forest edges, less common recently	x	x
Flycatcher, Olive-sided	<i>Contopus cooperi</i>	heard occasionally from hillside across valley	x	x
Flycatcher, Willow	<i>Empidonax traillii</i>	common breeder in alders along swamp edge	x	

Natural Resource Inventory for Lyme, NH
Bird Species of Southeastern & Northwestern Sections of Lyme, NH 1967-2006

Common Name	Scientific Name	Comments <u>primarily from</u> <u>Holmes in SE</u>	SE	NW
Green Warbler, Cape May	<i>Dendroica tigrina</i>	forest spring migrant	x	
Warbler, Chestnut-sided	<i>Dendroica pensylvanica</i>	breeder along forest/swamp edge	x	x
Warbler, Magnolia	<i>Dendroica magnolia</i>	occasional breeder	x	x
Warbler, Nashville	<i>Vermivora ruficapilla</i>	local breeder on hillside towards Roby's	x	
Warbler, Palm	<i>Dendroica palmarum</i>	two seen next to barn on 04/08/05	x	
Warbler, Pine	<i>Dendroica pinus</i>	spring migrant	x	
Warbler, Yellow	<i>Dendroica petechia</i>	common breeder in alders below field.	X	x
Warbler, Yellow-rumped	<i>Dendroica coronata</i>	common in spring and fall, probably breeds in forest	x	x
Waxwing, Bohemian	<i>Bombycilla garrulus</i>	flock of 20 on 03/05/94, 8-10 on 02/22/96	x	x
Waxwing, Cedar	<i>Bombycilla cedrorum</i>	breeds some years, regular in summer	x	x
Whip-poor-will	<i>Caprimulgus vociferus</i>	breeder through early 1970s, then absent	x	
Woodcock, American	<i>Scolopax minor</i>	breeding in fields, edge areas	x	x
Woodpecker, Downy	<i>Picoides pubescens</i>	breeder in forest	x	x
Woodpecker, Hairy	<i>Picoides villosus</i>	breeder in forest	x	x
Woodpecker, Pileated	<i>Dryocopus pileatus</i>	breeder in forest	x	x
Wren, House	<i>Troglodytes aedon</i>	common breeder in nest boxes	x	x
Wren, Winter	<i>Troglodytes troglodytes</i>	breeder in forest	x	x
Yellowthroat, Common	<i>Geothlypis trichas</i>	common breeder in thickets	x	x

Bird Species in Southeastern (SE) Lyme : Recorded by Richard and Deborah Holmes at their home at 31 Bliss Lane in Lyme Center. Richard Holmes holds a Ph.D. in Zoology from the University of California, Berkeley, and was a Professor of Biological Sciences and Environmental Biology at Dartmouth College from 1967 to 2004. He is now a Research Professor, as well as an Emeritus Professor, at Dartmouth. Professionally, he is an ornithologist, specializing in avian ecology. His research for the last 40 years has been focused on forest birds in New Hampshire, with field studies conducted primarily at the Hubbard Brook Experimental Forest in West Thornton, NH. He is a former vice-president of the American Ornithologists' Union (AOU), and has received a lifetime achievement awards from the AOU, the Wilson Ornithological Society and the Cooper Ornithological Society.

Bird Species in Northwestern (NW) Lyme: Recorded species seen or heard by John Skelly in the area of his home at 22 Pony Hill Lane. Mr. Skelly is a forester and has lived in Lyme since 1981. He holds a bioecology degree and taught high school biology including courses on predatory birds. He did post-graduate work in forestry and wildlife at UVM. Ecological parameters - 360 degree view with best and furthest visuals east to Dimmick hill and Lambert ridge, and west to near Vermont hills on Connecticut river and "the Pinnacle" off Franklin Hill road. Habitat includes beaver meadow wetland, through which flows Trout brook, a man-made pond less than one half acre, white pine and mixed hardwood second growth forest spreading east and north into Lyme town forest (Trout pond area Q), and open farmland with mixed wood (soft and hardwood) quadrants and corridors to the south and west. Elevation is near 1100'.

Relative abundance of birds is indicated by the following: common - seen most years; uncommon - seen occasionally; rare - only one or two sightings since 1981. Breeders that are obvious, which are most summer inhabitants, are designated as such, though I may not have them nesting in my area. Birds that visit the feeder, either summer or winter, are designated as feeder birds. Birds that have killed his chickens are labeled accordingly. Birds that could be seen in this area of Lyme Center (such as Peregrine falcons!), and that are locally not uncommon, but have not been seen or heard from our property, are not included in this listing...yet. Birds that are actually more common than witnessed in our area are designated only according to his view of their relative abundance or obscurity.