


20th Biennial
conference of
the Society
for Marine
Mammalogy




© Photo credits: Micheline Jenner

DECEMBER
2013


Assessing the effects of survey planning on abundance estimates of breeding stock 'D' humpback whales (*Megaptera novaeangliae*) from North West Cape, Western Australia

Bouchet, P.J.¹, Salgado Kent, C.P.², Jenner, C.K.³ & Jenner, M.N.³




THE UNIVERSITY OF
WESTERN AUSTRALIA
Achieve International Excellence

(1)



Curtin University

(3)



Centre for
WHALE RESEARCH
WESTERN AUSTRALIA

Salgado Kent *et al.* (2012)

J. CETACEAN RES. MANAGE. 12(1): 29-38, 2012

29

Southern Hemisphere Breeding Stock D humpback whale population estimates from North West Cape, Western Australia

CHANDRA SALGADO KENT*, CURT JENNER*, MICHELINE JENNER*, PHILIPPE BOUCHET* AND ERIC REXSTAD*


Contact e-mail: c.salgado@cmst.curtin.edu.au

ABSTRACT


Estimates of the abundance of Breeding Stock D humpback whales (*Megaptera novaeangliae*) are key to the conservation and management of what is thought to be one of the largest populations of the species. Five years (2000, 2001, 2006, 2007 and 2008) of aerial surveys carried out over an eight-year period at North West Cape (Western Australia) using line transect methodology allowed trends in whale numbers to be investigated, and provided a base for comparison with estimates made approximately 400km south at Shark Bay (Western Australia). A total of 3,127 whale detections were made during 74 surveys of the 7,043km² study area west of NWC. Pod abundance for each flight was computed using a Horvitz-Thompson like estimator and converted to an absolute measure of abundance after corrections were made for estimated mean cluster size, unsurveyed time, swimming speed and animal availability. Resulting estimates from the migration model of best fit with the most credible assumptions were 7,276 (CI = 4,993–10,167) for 2000, 12,280 (CI = 6,830–49,434) for 2001, 18,692 (CI = 12,980–24,477) for 2006, 20,044 (CI = 13,815–31,646) for 2007, and 26,100 (CI = 20,152–33,272) for 2008. Based on these data, the trend model with the greatest r^2 was exponential with an annual increase rate of 13% (CI = 5.6%–18.1%). While this value is above the species' estimated maximum plausible growth rate of 11.8%, it is reasonably close to previous reports of between 10–12%. The coefficient of variation, however, was too large for a reliable trend estimate. Perception bias was also not accounted for in these calculations. Based on a crude appraisal which yielded an estimated $p(0)$ of 0.783 (from independent observer effort, CV = 0.973), the 2008 humpback population size may be as large as 33,300. In conclusion, the work here provides evidence of an increasing Breeding Stock D population, but further surveys are necessary to confirm whether the population is indeed increasing at its maximum rate.

KEYWORDS: ABUNDANCE ESTIMATE; SURVEY-AERIAL; MIGRATION; MODELLING; TRENDS


Modelling migration counts



Population growth



Empirical approach: doing worse




Growth rates (confidence interval):

n=2: 0.103 (-0.123, 0.298)

n=3: 0.133 (-0.005, 0.260)


n=4: 0.136 (0.047, 0.229)

Simulation approach: doing better (?)




Random sampling:

n=4 to 30




Simulation approach: doing better (?)



Systematic sampling:


n=4 to 30 + random start (Yes/No)

Simulation approach: doing better (?)




Focal sampling – scenarios A to F: n=6 to 15


Simulation approach: doing better (?)




Simulation approach: doing better (?)



Detecting trends



Detecting trends



Conclusions and recommendations

- (1) Minimum requirement of **5 surveys**
- (2) Choice of design function of study goals
- (3) **Trend:** the more usually the better
- (4) Based on our original data, 6 and 9 years required to detect 10 and 5% trends
- (5) **Abundance:** 10-12 surveys optimal
- (6) F and D two best designs overall
- (7) Survey planning exercises **are critical to maximising cost-effectiveness**

Acknowledgements

Co-authors

Dr Chandra Salgado Kent
Centre for Marine
Science & Technology
Curtin University of Technology



**Mr Curt and
Mrs Micheline Jenner**
Centre for Whale
Research (WA) Inc.



Scholarships

Scholarship for International Research Fees (SIRF)
UWA Postgraduate Award (UPAIS)
UWA Top-up scholarship

Sponsors



Gibbs, J. P., Ene, E. (2010). Program MONITOR: Estimating the statistical power of ecological monitoring programs. <http://www.esf.edu/efb/gibbs/monitor/>

Salgado Kent, C., Jenner, C., Jenner, M., Bouchet, P. and Rexstad, E. (2012). Southern hemisphere breeding stock 'D' humpback whale population estimates from North West cape, Western Australia. *Journal of Cetacean Research and Management*, 12: 29-38.