

Frequentist and Bayesian methods for ill-posed inverse problems

Oracle and minimax optimality



Statistical estimation, frequentist versus Bayesian approach

Ill-posed inverse problems

Quantification of the quality of a statistical method




Statistical estimation, frequentist versus Bayesian approach


Ill-posed inverse problems

Quantification of the quality of a statistical method


Frequentist estimation, mainstay




Frequentist estimation, mainstay



Frequentist estimation, mainstay



Frequentist estimation, mainstay



Frequentist estimation, data example




Figure: Estimating the drift in a Gaussian process by aggregation of projection estimators



$$dY_x = \theta_x^\circ dx + \frac{1}{\sqrt{n}} dW(x);$$


$$Y_j = \mathcal{F}(Y)(j) = \mathcal{F}(\theta^\circ)(j) + \frac{1}{\sqrt{n}} \xi_j;$$

$$\xi_j \sim_{iid} \mathcal{N}(0, 1);$$


$$\widehat{\theta}_j = \mathbb{1}_{j \leq m} Y_j;$$

$$\widetilde{\theta}_x = \mathcal{F}^{-1}(\widehat{\theta})(x).$$

Bayesian estimation, mainstay



Bayesian estimation, mainstay



Bayesian estimation, data example




Figure: Sample from the posterior distribution of a hierarchical sieve prior



$$\boldsymbol{\theta}_j \sim \mathcal{N}(0, 1) \mathbb{1}_{j \leq m} + \delta_0 \mathbb{1}_{j > m};$$

$$Y_j | \boldsymbol{\theta}_j \sim \mathcal{N}\left(\boldsymbol{\theta}_j, \frac{1}{n}\right);$$

$$\boldsymbol{\theta}_j | Y_j \sim \mathcal{N}\left(\hat{\boldsymbol{\theta}}_j, \sigma_j\right) \mathbb{1}_{j \leq m} + \delta_0 \mathbb{1}_{j > m};$$

$$\hat{\boldsymbol{\theta}}_j = \frac{n \cdot Y_j}{1+n};$$

$$\sigma_j = \frac{1}{1+n}.$$




Statistical estimation, frequentist versus Bayesian approach


III-posed inverse problems

Quantification of the quality of a statistical method


III posed-inverse problems, mainstay




III posed-inverse problems, mainstay




III posed-inverse problems, mainstay




III posed-inverse problems, mainstay




III posed-inverse problems, mainstay




III posed-inverse problems, mainstay




III posed-inverse problems, mainstay




III posed-inverse problems, mainstay




III posed-inverse problems, mainstay




III posed-inverse problems, mainstay




III posed-inverse problems, mainstay



III posed-inverse problems, mainstay



III posed-inverse problems, mainstay



III posed-inverse problems, data example





Process: $\frac{\partial Y_{x,t}}{\partial t} = \frac{\partial^2 Y_{x,t}}{\partial x^2} + \frac{1}{n} \frac{dW(x,t)}{dt};$

Solution: $Y_j(t) = \theta_j^\circ(t) + \frac{1}{n} \xi_j(t)$
 $\theta_j^\circ(t) = \mathcal{F}(\theta^\circ(\cdot, 0))(j) \cdot \exp[-j^2 t]$

Observation: $Y_j = \theta_j^\circ(T) + \frac{1}{n} \xi_j;$


Interest: $\theta_j^\circ(0) = \theta_j^\circ(T) \exp[j^2 T].$



Statistical estimation, frequentist versus Bayesian approach

Ill-posed inverse problems


Quantification of the quality of a statistical method



► Distance to the truth:

$$l(\hat{\theta}, \theta^\circ)$$

Frequentist convergence rate, mainstay




- ▶ Distance to the truth:

$$l(\hat{\theta}, \theta^\circ)$$

- ▶ Average distance to the truth:

$$\mathbb{E}_{\theta^\circ} [l(\hat{\theta}, \theta^\circ)]$$



- ▶ Distance to the truth:

$$l(\hat{\theta}, \theta^\circ)$$

- ▶ Average distance to the truth:


$$\mathbb{E}_{\theta^\circ} [l(\hat{\theta}, \theta^\circ)]$$

- ▶ Lower bound: $\Phi_n(\theta^\circ) \rightarrow 0, K > 1$

$$\inf_{\tilde{\theta} \in \mathcal{E}} \mathbb{E}_{\theta^\circ} [l(\tilde{\theta}, \theta^\circ)] \geq K^{-1} \Phi_n(\theta^\circ).$$

Upper bound:

$$\mathbb{E}_{\theta^\circ} [l(\hat{\theta}, \theta^\circ)] \leq K \Phi_n(\theta^\circ).$$



- ▶ Distance to the truth:

$$l(\hat{\theta}, \theta^\circ)$$

- ▶ Average distance to the truth:


$$\mathbb{E}_{\theta^\circ} [l(\hat{\theta}, \theta^\circ)]$$

- ▶ Lower bound: $\Phi_n(\theta^\circ) \rightarrow 0, K > 1$

$$\inf_{\tilde{\theta} \in \mathcal{E}} \mathbb{E}_{\theta^\circ} [l(\tilde{\theta}, \theta^\circ)] \geq K^{-1} \Phi_n(\theta^\circ).$$

Upper bound:

$$\mathbb{E}_{\theta^\circ} [l(\hat{\theta}, \theta^\circ)] \leq K \Phi_n(\theta^\circ).$$



- Distance to the truth:

$$l(\hat{\theta}, \theta^\circ)$$

- Average distance to the truth:

$$\mathbb{E}_{\theta^\circ} [l(\hat{\theta}, \theta^\circ)]$$

- Lower bound: $\Phi_n(\theta^\circ) \rightarrow 0, K > 1$

$$\inf_{\tilde{\theta} \in \mathcal{E}} \mathbb{E}_{\theta^\circ} [l(\tilde{\theta}, \theta^\circ)] \geq K^{-1} \Phi_n(\theta^\circ).$$


Upper bound:

$$\mathbb{E}_{\theta^\circ} [l(\hat{\theta}, \theta^\circ)] \leq K \Phi_n(\theta^\circ).$$

Frequentist convergence rate, example



► $\|\hat{\theta} - \theta^\circ\|^2 = \int_{[0,1]} (\hat{\theta}(x) - \theta^\circ(x))^2 dx;$




Frequentist convergence rate, example



12

► $\mathbb{E}_{\theta^\circ} [\|\hat{\theta} - \theta^\circ\|^2] = \int_X \|\hat{\theta}(Y^n) - \theta^\circ\|^2 d\mathbb{P}_{Y^n};$




Frequentist convergence rate, example



12


► $\mathbb{E}_{\theta^\circ} [\|\hat{\theta} - \theta^\circ\|^2] = \mathcal{O}_n(\Phi_n^\circ)$



Bayesian contraction rate, principle




13



- ▶ Probability to give a guess within a distance of the truth:


$$\mathbb{P}_{\theta|Y^n}(\theta : l(\theta, \theta^\circ) \leq \Phi_n);$$

Bayesian contraction rate, principle



- ▶ Probability to give a guess within a distance of the truth:
$$\mathbb{P}_{\theta|Y^n}(\theta : l(\theta, \theta^\circ) \leq \Phi_n);$$
- ▶ Average probability to give a guess within a distance of the truth:

$$\mathbb{E}_{\theta^\circ}[\mathbb{P}_{\theta|Y^n}(\theta : l(\theta, \theta^\circ) \leq \Phi_n)];$$



- ▶ Probability to give a guess within a distance of the truth:

$$\mathbb{P}_{\theta|Y^n}(\theta : l(\theta, \theta^\circ) \leq \Phi_n);$$

- ▶ Average probability to give a guess within a distance of the truth:


$$\mathbb{E}_{\theta^\circ}[\mathbb{P}_{\theta|Y^n}(\theta : l(\theta, \theta^\circ) \leq \Phi_n)];$$

- ▶ Lower bound:

$$\lim_{n \rightarrow \infty} \sup_{Q_\theta \in \mathcal{G}} \mathbb{E}_{\theta^\circ}[\mathbb{P}_{\theta|Y^n}(\theta : l(\theta, \theta^\circ) \leq K^{-1}\Phi_n)] < 1;$$

upper bound:

$$\lim_{n \rightarrow \infty} \mathbb{E}_{\theta^\circ}[\mathbb{P}_{\theta|Y^n}(\theta : l(\theta, \theta^\circ) \leq K\Phi_n)] = 1.$$



- ▶ Probability to give a guess within a distance of the truth:

$$\mathbb{P}_{\theta|Y^n}(\theta : l(\theta, \theta^\circ) \leq \Phi_n);$$

- ▶ Average probability to give a guess within a distance of the truth:


$$\mathbb{E}_{\theta^\circ}[\mathbb{P}_{\theta|Y^n}(\theta : l(\theta, \theta^\circ) \leq \Phi_n)];$$

- ▶ Lower bound:

$$\lim_{n \rightarrow \infty} \sup_{Q_\theta \in \mathcal{G}} \mathbb{E}_{\theta^\circ}[\mathbb{P}_{\theta|Y^n}(\theta : l(\theta, \theta^\circ) \leq K^{-1}\Phi_n)] < 1;$$

upper bound:

$$\lim_{n \rightarrow \infty} \mathbb{E}_{\theta^\circ}[\mathbb{P}_{\theta|Y^n}(\theta : l(\theta, \theta^\circ) \leq K\Phi_n)] = 1.$$



- ▶ Probability to give a guess within a distance of the truth:

$$\mathbb{P}_{\theta|Y^n}(\theta : l(\theta, \theta^\circ) \leq \Phi_n);$$

- ▶ Average probability to give a guess within a distance of the truth:


$$\mathbb{E}_{\theta^\circ}[\mathbb{P}_{\theta|Y^n}(\theta : l(\theta, \theta^\circ) \leq \Phi_n)];$$

- ▶ Lower bound:

$$\lim_{n \rightarrow \infty} \sup_{Q_\theta \in \mathcal{G}} \mathbb{E}_{\theta^\circ}[\mathbb{P}_{\theta|Y^n}(\theta : l(\theta, \theta^\circ) \leq K^{-1}\Phi_n)] < 1;$$

upper bound:

$$\lim_{n \rightarrow \infty} \mathbb{E}_{\theta^\circ}[\mathbb{P}_{\theta|Y^n}(\theta : l(\theta, \theta^\circ) \leq K\Phi_n)] = 1.$$



- ▶ Probability to give a guess within a distance of the truth:

$$\mathbb{P}_{\theta|Y^n}(\theta : l(\theta, \theta^\circ) \leq \Phi_n);$$

- ▶ Average probability to give a guess within a distance of the truth:

$$\mathbb{E}_{\theta^\circ}[\mathbb{P}_{\theta|Y^n}(\theta : l(\theta, \theta^\circ) \leq \Phi_n)];$$

- ▶ Lower bound:

$$\lim_{n \rightarrow \infty} \sup_{Q_\theta \in \mathcal{G}} \mathbb{E}_{\theta^\circ}[\mathbb{P}_{\theta|Y^n}(\theta : l(\theta, \theta^\circ) \leq K^{-1}\Phi_n)] < 1;$$

upper bound:

$$\lim_{n \rightarrow \infty} \mathbb{E}_{\theta^\circ}[\mathbb{P}_{\theta|Y^n}(\theta : l(\theta, \theta^\circ) \leq K\Phi_n)] = 1.$$

Bibliography



- Comte, F., Cuenod, C.-A., Pensky, M., and Rozenholc, Y. (2017). Laplace deconvolution on the basis of time domain data and its application to dynamic contrast-enhanced imaging. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 79(1):69–94.
- Engl, H. W. and Kügler, P. (2005). Nonlinear inverse problems: theoretical aspects and some industrial applications. In *Multidisciplinary methods for analysis optimization and control of complex systems*, pages 3–47. Springer.
- Mohammad-Djafari, A. (2002). Bayesian inference for inverse problems. In *AIP Conference Proceedings*, volume 617, pages 477–496. AIP.
- Mohammad-Djafari, A., Giovannelli, J.-F., Demoment, G., and Idier, J. (2002). Regularization, maximum entropy and probabilistic methods in mass spectrometry data processing problems. *International Journal of Mass Spectrometry*, 215(1):175 – 193. Detectors and the Measurement of Mass Spectra.
- Naidu, P. S. and Westphal, K. O. (1965). The inverse problem of particle motion and its application. *Applied Scientific Research, Section B*, 12(5):435–450.