

Optimal aggregation in circular deconvolution by a frequentist Bayesian approach

StatMathAppli

Fréjus

Xavier Loizeau
4th of September 2017

In collaboration with
Fabienne Comte and Jan Johannes

Contents



Introduction

Suggested method

Optimality results


Simulations

Considered model

Circular deconvolution model




1



Considered model

Circular deconvolution model




Considered model

Circular deconvolution model




1



- ▶ Object of interest : density of a circular r.v. $X \sim f^X$;

Considered model


Circular deconvolution model



- ▶ Object of interest : density of a circular r.v. $X \sim f^X$;
- ▶ Noise : circular r.v. $\epsilon \sim f^\epsilon$;

Considered model

Circular deconvolution model




- ▶ Object of interest : density of a circular r.v. $X \sim f^X$;
- ▶ Noise : circular r.v. $\epsilon \sim f^\epsilon$;
- ▶ Observation : noisy circular r.v. $Y = X + \epsilon \sim f^Y = f^X \star f^\epsilon$

Considered model

Circular deconvolution model



1




- ▶ Object of interest : density of a circular r.v. $X \sim f^X$;
- ▶ Noise : circular r.v. $\epsilon \sim f^\epsilon$;
- ▶ Observation : noisy circular r.v. $Y = X + \epsilon \sim f^Y = f^X \star f^\epsilon$

Considered model

Circular deconvolution model



1



- ▶ Object of interest : density of a circular r.v. $X \sim f^X$;
- ▶ Noise : circular r.v. $\epsilon \sim f^\epsilon$;
- ▶ Observation : noisy circular r.v. $Y = X + \epsilon \sim f^Y = f^X \star f^\epsilon$

Considered model

Frequency domain



As all our variables have values in $[0,1]$, their densities can be extended in Fourier series. Hence we give the following definitions :

- ▶ $\forall j \in \mathbb{Z}, x \in [0,1], \quad e_j(x) = \exp[-2\pi \cdot i \cdot j \cdot x]$, the Fourier basis ;
- ▶ $[f^X]_j = \mathbb{E}_{f^X} [e_j(-X_1)]$, Fourier coefficients of f^X ;
- ▶ $[f^\epsilon]_j = \mathbb{E}_{f^\epsilon} [e_j(-\epsilon_1)]$, the Fourier coefficients of f^ϵ ;
- ▶ $[f^Y]_j = \mathbb{E}_{f^Y} [e_j(-Y_1)] = [f^X]_j \cdot [f^\epsilon]_j$, the Fourier coefficients of f^Y ;

Considered model

Illustration



Bayesian point of view

Bayesian fundamental paradigm



4

The problem is here treated from a Bayesian point of view :

- ▶ the parameter $[f]$ is a random variable with prior $\mathbb{P}_{[f]}$ with density $\pi_{[f]}$ with respect to some measure ;

Bayesian point of view

Bayesian fundamental paradigm



The problem is here treated from a Bayesian point of view :

- ▶ the parameter $[f]$ is a random variable with **prior** $\mathbb{P}_{[f]}$ with density $\pi_{[f]}$ with respect to some measure ;
- ▶ given $[f]$, the **likelihood** of Y is $\mathbb{P}_{Y|[f]}^n$ with density
$$\pi_{Y|[f]}^n(y_1, \dots, y_n, [f]) = \prod_{k \in [1, n]} \sum_{j \in \mathbb{Z}} [f]_j [f^c]_j e_j(y_k);$$

Bayesian point of view

Bayesian fundamental paradigm



The problem is here treated from a Bayesian point of view :

- ▶ the parameter $[f]$ is a random variable with prior $\mathbb{P}_{[f]}$ with density $\pi_{[f]}$ with respect to some measure ;
- ▶ given $[f]$, the likelihood of Y is $\mathbb{P}_{Y|[f]}^n$ with density
$$\pi_{Y|[f]}^n(y_1, \dots, y_n, [f]) = \prod_{k \in [1, n]} \sum_{j \in \mathbb{Z}} [f]_j [f^c]_j e_j(y_k);$$
- ▶ the marginal distribution of Y is \mathbb{P}_Y^n ;

Bayesian point of view

Bayesian fundamental paradigm



The problem is here treated from a Bayesian point of view :


- ▶ the parameter $[f]$ is a random variable with **prior** $\mathbb{P}_{[f]}$ with density $\pi_{[f]}$ with respect to some measure ;
- ▶ given $[f]$, the **likelihood** of Y is $\mathbb{P}_{Y|[f]}^n$ with density
$$\pi_{Y|[f]}^n(y_1, \dots, y_n, [f]) = \prod_{k \in [1, n]} \sum_{j \in \mathbb{Z}} [f]_j [f^c]_j e_j(y_k);$$
- ▶ the marginal distribution of Y is \mathbb{P}_Y^n ;
- ▶ we are interested in the **posterior distribution** $\mathbb{P}_{[f]|Y}^n$ with density
$$\pi_{[f]|Y}^n([f], y) \propto \pi_{Y|[f]}^n(y, [f]) \cdot \pi_{[f]}([f]).$$

Formulation of optimality

Frequentist case



5



Formulation of optimality


Frequentist case



5

- ▶ For each frequentist estimator, measure the performance by its risk

$$\mathbb{E}_{f^X}^n \left[d\left(\tilde{f}, f^X\right)^2 \right].$$




Formulation of optimality

Frequentist case



5

- For each frequentist estimator, measure the performance by its risk



- Goal : derive a lower bound for this risk

$$\inf_{\tilde{f}} \mathbb{E}_{f^X}^n \left[d(\tilde{f}, f^X)^2 \right] \geq \mathcal{R}_n^\circ(f^X)$$

over a family of estimators.

Formulation of optimality

Frequentist case



5


- ▶ For each frequentist estimator, measure the performance by its risk

$$\mathbb{E}_{f^X}^n \left[d(\tilde{f}, f^X)^2 \right].$$

- ▶ Goal : derive a lower bound for this risk

$$\inf_{\tilde{f}} \mathbb{E}_{f^X}^n \left[d(\tilde{f}, f^X)^2 \right] \geq \mathcal{R}_n^\circ(f^X)$$

over a family of estimators.



Formulation of optimality

Frequentist case



5


- For each frequentist estimator, measure the performance by its risk

$$\mathbb{E}_{f^X}^n \left[d(\tilde{f}, f^X)^2 \right].$$

- Goal : derive a lower bound for this risk

$$\inf_{\tilde{f}} \mathbb{E}_{f^X}^n \left[d(\tilde{f}, f^X)^2 \right] \geq \mathcal{R}_n^\circ(f^X)$$

over a family of estimators.




Formulation of optimality

Frequentist case



5



- For each frequentist estimator, measure the performance by its risk

$$\mathbb{E}_{f^X}^n \left[d(\tilde{f}, f^X)^2 \right].$$

- Goal : derive a lower bound for this risk

$$\inf_{\tilde{f}} \mathbb{E}_{f^X}^n \left[d(\tilde{f}, f^X)^2 \right] \geq \mathcal{R}_n^\circ(f^X)$$

over a family of estimators.

- Finding \hat{f} in this family such that

$$\mathbb{E}_{f^X}^n \left[d(\hat{f}, f^X)^2 \right] \leq K^\circ \cdot \mathcal{R}_n^\circ(f^X),$$


it is then **oracle rate optimal** and **adaptive** if \hat{f} does not depend on f^X .

Formulation of optimality

Frequentist case



5



- For each frequentist estimator, measure the performance by its risk

$$\mathbb{E}_{f^X}^n \left[d(\tilde{f}, f^X)^2 \right].$$

- Goal : derive a lower bound for this risk

$$\inf_{\tilde{f}} \mathbb{E}_{f^X}^n \left[d(\tilde{f}, f^X)^2 \right] \geq \mathcal{R}_n^\circ(f^X)$$

over a family of estimators.

- Finding \hat{f} in this family such that

$$\mathbb{E}_{f^X}^n \left[d(\hat{f}, f^X)^2 \right] \leq K^\circ \cdot \mathcal{R}_n^\circ(f^X),$$


it is then **oracle rate optimal** and **adaptive** if \hat{f} does not depend on f^X .

Formulation of optimality

Frequentist case



5



- For each frequentist estimator, measure the performance by its risk

$$\mathbb{E}_{f^X}^n \left[d(\tilde{f}, f^X)^2 \right].$$

- Goal : derive a lower bound for this risk

$$\inf_{\tilde{f}} \mathbb{E}_{f^X}^n \left[d(\tilde{f}, f^X)^2 \right] \geq \mathcal{R}_n^\circ(f^X)$$

over a family of estimators.

- Finding \hat{f} in this family such that

$$\mathbb{E}_{f^X}^n \left[d(\hat{f}, f^X)^2 \right] \leq K^\circ \cdot \mathcal{R}_n^\circ(f^X),$$

it is then **oracle rate optimal** and **adaptive** if \hat{f} does not depend on f^X .

Formulation of optimality

Pragmatic Bayesian paradigm



6

How to transfer this in a Bayesian point of view ?

Taking a "pragmatic Bayesian" point of view :

- ▶ f^X the **true parameter**.

- ▶ Is $\mathbb{P}_{f|Y}^n$ **shrinking** around f^X as n tends to ∞ ?

- ▶ How fast ?


Formulation of optimality

Pragmatic Bayesian formulation of optimality



7

► Concentration rate $(\phi_n)_{n \in \mathbb{N}}$




$$\exists c \in \mathbb{R}_+, \quad \lim_{n \rightarrow \infty} \mathbb{E}_{f^X}^n \left[\mathbb{P}_{f|Y}^n \left(d(f, f^X)^2 \geq c \phi_n \right) \right] = 0.$$

Formulation of optimality

Pragmatic Bayesian formulation of optimality



► Concentration rate $(\phi_n)_{n \in \mathbb{N}}$




$$\exists c \in \mathbb{R}_+, \quad \lim_{n \rightarrow \infty} \mathbb{E}_{f|Y}^n \left[\mathbb{P}_{f|Y}^n \left(d(f, f^X) \geq c \phi_n \right) \right] = 0.$$

Formulation of optimality

Pragmatic Bayesian formulation of optimality



► Concentration rate $(\phi_n)_{n \in \mathbb{N}}$



$$\exists c \in \mathbb{R}_+, \quad \lim_{n \rightarrow \infty} \mathbb{E}_{f^X}^n \left[\mathbb{P}_{f|Y}^n \left(d(f, f^X)^2 \geq c \phi_n \right) \right] = 0.$$


Formulation of optimality

Pragmatic Bayesian formulation of optimality



7

► Concentration rate $(\phi_n)_{n \in \mathbb{N}}$




$$\exists c \in \mathbb{R}_+, \quad \lim_{n \rightarrow \infty} \mathbb{E}_{f^X}^n \left[\mathbb{P}_{f|Y}^n \left(d(f, f^X)^2 \geq c \phi_n \right) \right] = 0.$$

Formulation of optimality

Pragmatic Bayesian formulation of optimality



- ▶ Concentration rate $(\phi_n)_{n \in \mathbb{N}}$



$$\exists c \in \mathbb{R}_+, \quad \lim_{n \rightarrow \infty} \mathbb{E}_{f^X}^n \left[\mathbb{P}_{f|Y}^n \left(d(f, f^X)^2 \geq c \phi_n \right) \right] = 0.$$

- ▶ Exact concentration rate $(\phi_n)_{n \in \mathbb{N}}$ if in addition


$$\lim_{n \rightarrow \infty} \mathbb{E}_{f^X}^n \left[\mathbb{P}_{f|Y}^n \left(d(f, f^X)^2 \leq c^{-1} \phi_n \right) \right] = 0.$$

Formulation of optimality

Pragmatic Bayesian formulation of optimality



- ▶ Concentration rate $(\phi_n)_{n \in \mathbb{N}}$



$$\exists c \in \mathbb{R}_+, \quad \lim_{n \rightarrow \infty} \mathbb{E}_{f^X}^n \left[\mathbb{P}_{f|Y}^n \left(d(f, f^X)^2 \geq c \phi_n \right) \right] = 0.$$

- ▶ Exact concentration rate $(\phi_n)_{n \in \mathbb{N}}$ if in addition

$$\lim_{n \rightarrow \infty} \mathbb{E}_{f^X}^n \left[\mathbb{P}_{f|Y}^n \left(d(f, f^X)^2 \leq c^{-1} \phi_n \right) \right] = 0.$$


Formulation of optimality

Pragmatic Bayesian formulation of optimality



7

- ▶ Concentration rate $(\phi_n)_{n \in \mathbb{N}}$



$$\exists c \in \mathbb{R}_+, \quad \lim_{n \rightarrow \infty} \mathbb{E}_{f^X}^n \left[\mathbb{P}_{f|Y}^n \left(d(f, f^X)^2 \geq c \phi_n \right) \right] = 0.$$

- ▶ Exact concentration rate $(\phi_n)_{n \in \mathbb{N}}$ if in addition

$$\lim_{n \rightarrow \infty} \mathbb{E}_{f^X}^n \left[\mathbb{P}_{f|Y}^n \left(d(f, f^X)^2 \leq c^{-1} \phi_n \right) \right] = 0.$$


Formulation of optimality

Pragmatic Bayesian formulation of optimality



7

- ▶ Concentration rate $(\phi_n)_{n \in \mathbb{N}}$



$$\exists c \in \mathbb{R}_+, \quad \lim_{n \rightarrow \infty} \mathbb{E}_{f^X}^n \left[\mathbb{P}_{f|Y}^n \left(d(f, f^X)^2 \geq c \phi_n \right) \right] = 0.$$

- ▶ Exact concentration rate $(\phi_n)_{n \in \mathbb{N}}$ if in addition

$$\lim_{n \rightarrow \infty} \mathbb{E}_{f^X}^n \left[\mathbb{P}_{f|Y}^n \left(d(f, f^X)^2 \leq c^{-1} \phi_n \right) \right] = 0.$$

A popular frequentist method

Projection estimators



From a frequentist point of view, a natural method to answer this problem is :

- ▶ define $\widehat{[f^Y]}_j = \frac{1}{n} \sum_{k=1}^n e_j(-Y_k)$, an unbiased estimator for $[f^Y]_j = \mathbb{E}_{f^Y} [e_j(-Y_1)]$;

A popular frequentist method

Projection estimators



From a frequentist point of view, a natural method to answer this problem is :

- ▶ define $\widehat{[f^Y]}_j = \frac{1}{n} \sum_{k=1}^n e_j(-Y_k)$, an unbiased estimator for $[f^Y]_j = \mathbb{E}_{f^Y} [e_j(-Y_1)]$;
- ▶ consider $\widehat{[f]}_j = \frac{\widehat{[f^Y]}_j}{\widehat{[f^e]}_j}$, an unbiased estimator for $[f^X]_j$;

A popular frequentist method

Projection estimators



From a frequentist point of view, a natural method to answer this problem is :

- ▶ define $\widehat{[f^Y]}_j = \frac{1}{n} \sum_{k=1}^n e_j(-Y_k)$, an unbiased estimator for $[f^Y]_j = \mathbb{E}_{f^Y} [e_j(-Y_1)]$;
- ▶ consider $\widehat{[f]}_j = \frac{\widehat{[f^Y]}_j}{\widehat{[f^e]}_j}$, an unbiased estimator for $[f^X]_j$;
- ▶ use an element from the family of projection estimators :
$$\left\{ \left(\widehat{[f]}_j^m \right)_{j \in \mathbb{Z}} : \forall j \in \mathbb{Z}, \quad \widehat{[f]}_j^m = \widehat{[f]}_j \mathbb{1}_{\{0 < |j| \leq m\}} + \mathbb{1}_{\{j=0\}} \right\};$$

A popular frequentist method

Projection estimators



From a frequentist point of view, a natural method to answer this problem is :

- ▶ define $\widehat{[f^Y]}_j = \frac{1}{n} \sum_{k=1}^n e_j(-Y_k)$, an unbiased estimator for $[f^Y]_j = \mathbb{E}_{f^Y} [e_j(-Y_1)]$;
- ▶ consider $\widehat{[f]}_j = \frac{\widehat{[f^Y]}_j}{\widehat{[f^e]}_j}$, an unbiased estimator for $[f^X]_j$;
- ▶ use an element from the family of projection estimators :
$$\left\{ \left(\widehat{[f]}_j^m \right)_{j \in \mathbb{Z}} : \forall j \in \mathbb{Z}, \quad \widehat{[f]}_j^m = \widehat{[f]}_j \mathbb{1}_{\{0 < |j| \leq m\}} + \mathbb{1}_{\{j=0\}} \right\};$$
- ▶ define a method to select m in a satisfactory way (e.g. Massart [2003]).

Illustration

Direct case



squared bias, variance and MSE




Figure: MSE of projection estimators in the direct case

Illustration

Direct case



10

Illustration

Inverse case



11

squared bias, variance and MSE




Figure: MSE of projection estimators in the mildly ill-posed case

Considered model

Illustration



12

Contents



12

Introduction

Suggested method

Optimality results

Simulations



We first introduce the so called sieve priors with deterministic threshold parameter m :

$$\pi_{[f]}([f]) = \prod_{|j|>m} \delta_0([f]_j) \cdot \exp[\Delta_1([f])].$$



The posterior is then of the form

$$\pi_{[f]|Y^n}([f], y^n) = \prod_{|j|>m} \delta_0([f]_j) \cdot \\ \exp \left[\underbrace{\Delta_1([f]) + \sum_{k=1}^n \log \left(1 + \sum_{0<|j|\leq m} [f]_j \cdot [f^\epsilon]_j \cdot e_j(y_k) \right) + \Delta_2(y^n)}_{R([f], y^n)} \right].$$



Open question : By using a Taylor expansion of the log around 1, it seems like this distribution could be approximated by a Gaussian, indeed :

$$\begin{aligned}
 R([f], y^n) = & - \sum_{0 < |j| \leq m} \frac{|[f]_j - [\widehat{f}]_j|^2}{2 \cdot \sigma_j} \\
 & + \sum_{l=2}^{\infty} \sum_{k=1}^n \frac{(-1)^{l+1}}{l} \left(\sum_{0 < |j| \leq m} [f]_j \cdot [f^\epsilon]_j \cdot e_j(y_k) \right)^l + \delta_1([f]) + \delta_2(y^n);
 \end{aligned}$$

$$\text{where } \sigma_j = \frac{1}{1+n|[f^\epsilon]_j|^2} \text{ and } [\widehat{f}]_j = \frac{[f^\epsilon]_j \cdot \sum_{k=1}^n e_j(y_k)}{1+n|[f^\epsilon]_j|^2}$$

Looking for a prior

Hierarchical prior



- ▶ Consider a **random hyper-parameter M** , with values in $\llbracket 1, n \rrbracket$, acting like a threshold :

$$\forall j \in \mathbb{Z} : |j| > m, \quad \mathbb{P}_{[\mathbf{f}]_j | M=m} = \delta_0$$

- ▶ Write the density of M $\pi_M(m) \propto \exp[\text{pen}(m)]$ with respect to the counting measure, then, the posterior can be written

$$\pi_{M|Y^n}(m, y^n) = \exp[\text{pen}(m) - 2\Delta_2(y^n)]$$

$$\pi_{[\mathbf{f}]|Y^n} = \sum_{m \in \mathbb{N}} \pi_{[\mathbf{f}]|M=m, Y^n} \cdot \pi_{M=m|Y^n};$$

$$\widetilde{[\mathbf{f}]} = \left(\mathbb{E}_{[\mathbf{f}]|M \geq j, Y^n} \left[[\mathbf{f}]_j \right] \cdot \mathbb{P}_{M|Y^n}(M \geq j) \right)_{j \in \mathbb{N}}.$$

Hierarchical prior

Graphical representation



Contents



17

Introduction

Suggested method

Optimality results

Simulations

Optimality results

Definitions



18

We here note : $\Phi_n^\circ = \Phi_n^\circ([f^X])$ the optimal convergence rate for the family of projection estimators for each $[f^X]$; this objects has analytical formulation (see for example Comte and Taupin [2003]).



Theorem : oracle optimality J & C & L [2017]

For each $[f^X]$, there exist C° such that

$$\forall n \in \mathbb{N}^*, \quad \mathbb{E}_{[f^X]}^n \left[\left\| [\widetilde{f}] - [f^X] \right\|^2 \right] \leq C^\circ \Phi_n^\circ.$$



Theorem J & C & L [2017]

Φ_n° can be considered as upper bound for the family of sieve priors :

$$\lim_{n \rightarrow \infty} \inf_{\mathbb{Q}_{[f]}} \mathbb{E}_{[f^X]}^n \left[\mathbb{Q}_{[f] \mid Y^n} \left(\left\| [f] - [f^X] \right\|^2 \geq \Phi_n^\circ \right) \right] = 1,$$

where $\inf_{\mathbb{Q}_{[f]}}$ is taken over the family of sieve priors with Gaussian margins.

This gives a purely Bayesian formulation of oracle optimality.



Theorem : oracle concentration J & C & L [2017]

For all $[f^X]$ in Θ° , there exist K° such that

$$\lim_{n \rightarrow \infty} \mathbb{E}_{[f^X]}^n \left[\mathbb{P}_{[\mathbf{f}], M | Y^n} \left((K^\circ)^{-1} \Phi_n^\circ \leq \| [\mathbf{f}] - [f^X] \|^2 \leq K^\circ \Phi_n^\circ \right) \right] = 1.$$

Contents



Introduction

Suggested method

Optimality results

Simulations

Direct case : visualisation of the estimates



Direct case : sampling from the approximated posterior



23




Figure: Evolution of the empirical mean squared error of the two estimates with respect to the number of observations in the direct case

Direct case : empirical mean square errors of the estimates



24




Figure: Sample from the posterior distribution and the posterior mean in the direct case



- ▶ Bayesian method for circular deconvolution with interpretable prior ;
- ▶ providing a fully data driven estimator, fulfilling oracle optimality ;
- ▶ giving a Bayesian formulation of Oracle optimality...

And much more !

- ▶ Also fulfills minimax optimality (no log term !)
- ▶ actually provides a complete family of optimal estimators using iterated posterior method
- ▶ works with β -mixing dependance
- ▶ suitable for severely ill-posed problems
- ▶ can be adapted to real line deconvolution