



A

PrEP Transcriptomic Cohort




B

IBDMDB



C

PrEP Proteomic Cohort



Microbes

Akkermansia muciniphila	Bacteroides stercoris	Dorea formicigenerans	Lactobacillus hamsteri	Parabacteroides distasonis
Atopobium vaginae	Bacteroides thetaiotaomicron	Escherichia coli	Lactobacillus helveticus	Parabacteroides merdae
Bacteroides caccae	Bacteroides uniformis	Fusobacterium nucleatum	Lactobacillus hominis	Prevotella
Bacteroides cellulosilyticus	Bacteroides vulgatus	Gardnerella vaginalis	Lactobacillus iners	Pseudomonas
Bacteroides dorei	Bifidobacterium	Lactobacillus	Lactobacillus jensenii	Pseudomonas fulva
Bacteroides eggerthii	Bifidobacterium breve	Lactobacillus crispatus	Lactobacillus ultunensis	Pseudomonas syringae
Bacteroides fragilis	Collinsella aerofaciens	Lactobacillus gallinarum	Lactobacillus vaginalis	Undistinguishable
Bacteroides ovatus	Coprococcus comes	Lactobacillus gasseri	Mobiluncus mulieris	Other Microbes

Supplementary Figure 1: The microbiome composition of the cohorts for the vaginal models (A and C) and gut models (B). The vaginal cohorts share 24 microbes which are colored, microbes which are not shared are designated as "Other Microbes". The gut microbes represented are the 27 that are shared between the IBDMDB gut cohort and the Zimmerman et al 2019 data used to assign metabolic relationships to the microbe-drug pairs. Microbes that are not shared with the Zimmerman et al 2019 data are denoted as "Other Microbes".