



## Review

## Role of Estrogen in Type 1 and Type 2 Diabetes Mellitus: A Review of Clinical and Preclinical Data

Monica De Paoli MD<sup>a,b</sup>; Geoff H. Werstuck PhD<sup>a,b,\*</sup><sup>a</sup>Thrombosis and Atherosclerosis Research Institute, McMaster University, Hamilton, Ontario, Canada<sup>b</sup>Department of Medicine, McMaster University, Hamilton, Ontario, Canada

## Key Messages

- Premenopausal women are protected from developing diabetes and its complications.
- Preclinical studies suggest that estrogen directly affects beta-cell health and function.

## ARTICLE INFO

## Article history:

Received 1 August 2019

Received in revised form

17 September 2019

Accepted 6 January 2020

## Keywords:

beta cell  
cardiovascular disease  
diabetes  
estrogen

## Mots clés :

cellule bêta  
maladie cardiovasculaire  
diabète  
œstrogène

## ABSTRACT

The incidence and prevalence of diabetes mellitus, and the cardiovascular complications associated with this disease, are rapidly increasing worldwide. Individuals with diabetes have a higher mortality rate due to cardiovascular diseases and a reduced life expectancy compared to those without diabetes. This poses a significant economic burden on health-care systems worldwide, making the diabetes epidemic a global health crisis. Sex differences in the presentation and outcome of diabetes do exist. Premenopausal women are protected from developing diabetes and its cardiovascular complications relative to males and postmenopausal women. However, women with diabetes tend to have a higher mortality as a result of cardiovascular complications than age-matched men. Despite this evidence, preclinical and clinical research looking at sex as a biologic variable in metabolic disorders and their cardiovascular complications is very limited. The aim of this review is to highlight the current knowledge of the potential protective role of estrogens in humans as well as rodent models of diabetes mellitus, and the possible pathways by which this protection is conferred. We stress the importance of increasing knowledge of sex-specific differences to facilitate the development of more targeted prevention strategies.

© 2020 Canadian Diabetes Association.

## R É S U M É

L'incidence et la prévalence du diabète sucré et les complications cardiovasculaires associées à cette maladie ne cessent d'augmenter dans le monde. Les personnes diabétiques ont un taux de mortalité plus élevé en raison des maladies cardiovasculaires et une espérance de vie plus courte que les personnes non diabétiques. Pour les systèmes de soins de santé du monde entier, il s'agit d'un fardeau économique important, qui fait de l'épidémie du diabète une crise mondiale de la santé. Le tableau clinique et l'évolution du diabète montrent des différences entre les sexes. Par rapport aux hommes et aux femmes postménopausées, les femmes préménopausées sont à l'abri du diabète et de ses complications cardiovasculaires. Toutefois, du fait des complications cardiovasculaires, les femmes diabétiques présentent une mortalité plus élevée que les hommes appariés selon l'âge. En dépit de ces données scientifiques, la recherche préclinique et clinique, qui montre que le sexe est une variable biologique des perturbations métaboliques et de leurs complications cardiovasculaires, est très limitée. L'objectif de la présente étude est de présenter les connaissances actuelles sur le rôle protecteur potentiel des œstrogènes dans des modèles de diabète sucré chez les humains et chez les rongeurs, et les voies par lesquelles cette protection peut être obtenue. Nous insistons sur l'importance d'accroître les connaissances sur les différences liées au sexe pour faciliter l'élaboration de stratégies de prévention plus ciblées.

© 2020 Canadian Diabetes Association.

\* Address for correspondence: Geoff H. Werstuck PhD, Thrombosis and Atherosclerosis Research Institute, McMaster University, 237 Barton Street East, Hamilton, Ontario L8L 2X2, Canada.

E-mail address: [geoff.werstuck@taari.ca](mailto:geoff.werstuck@taari.ca)

1499-2671/© 2020 Canadian Diabetes Association.

The Canadian Diabetes Association is the registered owner of the name Diabetes Canada.

<https://doi.org/10.1016/j.cjcd.2020.01.003>

## Introduction

Diabetes mellitus is a group of metabolic disorders characterized by the presence of elevated blood glucose levels (hyperglycemia). The underlying characteristic of all forms of diabetes mellitus is the inability of pancreatic beta cells to produce/secrete sufficient quantities of insulin to regulate glucose levels. In type 1 diabetes mellitus (T1DM), the inability of regulating blood glucose levels results from the rapid destruction of pancreatic beta cells due to autoimmune processes, which leads to insulin deficiency. In type 2 diabetes mellitus (T2DM) hyperglycemia is the result of an insufficient insulin secretion combined with an impaired response to insulin (insulin resistance) in the hepatic, skeletal and/or adipose tissues. The early stages of T2DM are characterized by pancreatic beta-cell hyperfunction, which leads to dysfunction and, ultimately, apoptosis of the beta cells (1).

Diabetes mellitus is associated with the development of microvascular and macrovascular complications. Microvascular complications include diabetic nephropathy, neuropathy and retinopathy. Macrovascular complications include coronary artery disease, peripheral arterial disease and stroke (2). Atherosclerosis is the main underlying cause of macrovascular complications (2), and people with diabetes have a 2-fold increased risk of death (3) as well as a reduced life expectancy by up to 15 years (4,5) when compared to those without diabetes. Individuals with diabetes are also at a higher risk of developing blindness, renal failure, chronic neuropathic pain, below-knee amputations, cancer, depression, cognitive decline, cirrhosis, frailty and a significant increase in the risk of cardiovascular disease (CVD), even when controlling for other CV risk factors (6,7).

In the last few decades, as a result of changes in lifestyle and increased rates of obesity, there has been a dramatic increase in the incidence and prevalence of diabetes mellitus worldwide (8–10). The costs associated with the treatment of diabetes and its complications present a major burden on health-care systems around the world (11,12). Herein, we review the clinical data pertaining to sex differences in the prevalence of diabetes and its CV complications. In addition, we examine rodent models of diabetes and CV complications in which similar sex-specific differences have been observed. We discuss how these models have been used to highlight the protective role of estrogen and to suggest potential mechanism(s) by which this protection is conferred.

## Sex Differences in Diabetes and Cardiovascular Complications

Sex differences in the pathophysiology of diabetes mellitus do exist and females have a protective advantage compared with males. Many studies have shown that there is a higher prevalence of T1DM in males than in females (13,14). Moreover, sexual dimorphism can also be observed in T2DM. Specifically, premenopausal women have a lower prevalence of T2DM when compared with men or postmenopausal women (15). This trend has been observed in many studies worldwide that show a persistent male predominance in the incidence of T2DM in the long-term follow up of patients (16–20). This evidence suggests a potential protective role of estrogens in the development of diabetes. This hypothesis is supported by 2 large, randomized, double-blind, placebo-controlled clinical trials, the Women's Health Initiative study and the Heart and Estrogen/Progestin Replacement Study, which showed that women taking hormone replacement therapy, were less prone to develop diabetes than those of a placebo group (21,22).

Sex differences can also be observed in diabetes-associated CVDs, which represent the major cause of mortality and morbidity among patients with diabetes (23). Evidence shows that

there is a 3- to 5-fold increased risk in coronary heart disease (CHD) in men without diabetes vs premenopausal females (24). Furthermore, women tend to develop CHD later than men, supporting a potential protective role of female sex hormones (25). However, when women are diagnosed with diabetes, this protective factor is erased, as shown by a recent meta-analysis concluding that women with diabetes have a 40% increased risk of developing CHD compared to men with diabetes (26). Consistent with these findings, a meta-analysis of prospective cohort studies showed that women with diabetes have about 50% greater risk of CHD than men with diabetes, even after consideration of other major cardiovascular risk factors (27). Another systematic review showed that the relative risk of stroke in women with diabetes was 27% higher than males with diabetes, after taking into account baseline and major cardiovascular risk factors (28). Similarly, a Japanese study showed that women with diabetes have a higher risk of a stroke event compared with males (29). A collaborative meta-analysis of 102 prospective studies worldwide showed similar outcomes in terms of the presence of diabetes in women and the risk of CVD (20).

The higher rate of CVD in women with diabetes results in a higher CV mortality compared with women or men without diabetes (30). A Canadian study showed that hospitalization and mortality risk due to CVD was significantly increased in women with diabetes vs those without diabetes. Strikingly, the mortality rate was 2-fold higher in women than men (31). Consistent with these data, 2 population-based studies in Sweden and Italy, respectively, highlighted that the survival rate after stroke or other CVDs was significantly lower in women with diabetes compared with men (32,33). It was previously thought that the sex-specific severity of diabetes-associated CV complications was a result of disparities in the management of diabetes between male and female patients; however, as treatment has become more equal between sexes, the greater risk of severe CVD in women has persisted (26). This finding supports a potential protective role of estrogens, because women tend to develop diabetes after menopause and have a higher risk of severe diabetes-associated CVD complications and consequent higher mortality rate compared to men with diabetes.

The current trends for diabetes and its cardiovascular complications predict a progressive increase in their prevalence, which will pose a significant economic and managerial burden in health-care systems worldwide (34). Further research needs to be carried out to increase our understanding of the role of sex in the development and progression of diabetes mellitus and its cardiovascular complications. This will facilitate the establishment of targeted strategies to more effectively prevent and manage these diseases.

## Potential Role of Estrogen in Diabetes and Its Cardiovascular Complications

The increased CV risk in postmenopausal women may result from the reduction in circulating estrogen and the associated risk of increased body weight, which can result in the onset of insulin resistance (35). Insulin resistance causes beta cells to significantly hypersecrete insulin. Elevated circulating insulin concentrations can have deleterious effects on the vasculature, such as sodium and liquid retention, vasoconstriction, proliferation of smooth muscle cells and proinflammatory activity, which can induce high blood pressure as well as development of atherosclerosis (36,37). Hyperglycemia, high blood pressure and atherosclerosis represent predominant risk factors for CVDs, hence they pose a significant risk in postmenopausal women (36). The underlying mechanisms by which estrogens may confer protection against diabetes and its associated vascular complications have begun to be investigated in new rodent models of cardiometabolic disease (38).

## Rodent Models of Sexual Dimorphism in Diabetes Mellitus

Rodent models of diabetes mellitus are very useful when it comes to delineating the pathogenesis of metabolic disorders. Many rodent models of diabetes and/or hyperglycemia present sexual dimorphism in terms of glucose regulation and associated complications that are similar to those observed in human patients (Table 1). To mimic menopause and to more clearly investigate the roles of ovarian hormones, ovariectomized rodents are commonly used as an experimental model (39).

To our knowledge, the only study to address sex-specific differences in the development of hyperglycemia-induced atherosclerosis involved the ApoE<sup>-/-</sup>:Ins2<sup>+Akita</sup> mouse model (38). This mouse strain shows sex dimorphism in glucose regulation with male mice developing chronic hyperglycemia and female mice being transiently hyperglycemic (38). Furthermore, atherosclerotic analysis showed that both male and female ApoE<sup>-/-</sup>:Ins2<sup>+Akita</sup> mice developed atherosclerosis; however, ovariectomized ApoE<sup>-/-</sup>:Ins2<sup>+Akita</sup> mice developed chronic hyperglycemia and showed signs of advanced atherosclerosis, similar to what was observed in male hyperglycemic ApoE<sup>-/-</sup>:Ins2<sup>+Akita</sup> mice (38). This suggests that the loss of estrogens contributes to the development of hyperglycemia and atherosclerosis, both significant risk factors for CVD.

## Mechanisms of Estrogen Protection

The sex differences in glucose homeostasis observed in both humans and rodent models suggest a potential protective role of female sex hormones in the prevention of diabetes. The mechanisms by which this protection is conferred are still not clear.

Estradiol has been shown to increase insulin content and glucose-stimulated insulin secretion in isolated mouse pancreatic islets, suggesting an active role for this hormone in glucose homeostasis (40). It was shown that ovariectomized C57BL/6J mice have an altered glucose homeostasis characterized by impaired glycemia, reduced glucose tolerance, insulinemia and impaired insulin secretion when compared to females supplemented with estradiol (41). Another study in leptin-resistant female Zucker Diabetic Fatty Rats showed that fasting blood glucose levels were higher and glucose tolerance was impaired in an ovariectomized group as opposed to sham controls or ovariectomized females

given estradiol treatment (42). Consistent with these observations, many other studies have highlighted the beneficial role of estradiol in glucose homeostasis (43–45).

Pancreatic beta cells express all 3 major estrogen receptors (ERs) ER-alpha, ER-beta and the G protein-coupled estrogen receptor, and there is evidence suggesting a role for each of these receptors in the maintenance and enhancement of beta-cell function (46–48). A study conducted in mice lacking either ER-alpha or ER-beta showed that beta-cell survival was significantly impaired in both models when compared with controls (49). The same study also showed that female mice lacking G protein-coupled estrogen receptor lost their protection against streptozotocin-induced beta-cell apoptosis (49).

Oxidative stress has been implicated as a mechanism by which hyperglycemia may promote beta-cell dysfunction. However, antioxidant therapies have failed to significantly protect beta cells from apoptosis (50,51). The role of endoplasmic reticulum (ER) stress in beta-cell dysfunction is currently being investigated as an alternative pathway by which elevated concentrations of glucose may damage beta cells. The ER is an organelle in the cytoplasm of the cell whose function is to synthesize, properly fold and deliver proteins to the location of interest. Pancreatic beta cells have a complex, well-developed ER because they need to synthesize and secrete large quantities of insulin in response to the physiologic needs of glucose regulation (52). However, in situations of chronic hyperglycemia and/or insulin resistance, the demand of insulin is significantly increased and, as a result, the folding capacity of the ER is also increased. When the functional folding capacity of the ER is exceeded, unfolded proteins accumulate and induce a condition known as ER stress (53,54). The accumulated misfolded proteins trigger the unfolded protein response, which is an adaptive mechanism aimed at restoring homeostasis within the ER (55). The unfolded protein response activates the expression of ER-resident protein chaperones that function to increase the protein folding capacity of the cell. If the ER homeostasis cannot be restored, apoptotic mechanisms are triggered (56,57).

An in vitro study conducted using the INS-1 cell line showed that elevated concentrations of glucose can induce the expression of ER stress markers, but treatment of these cells with estradiol can significantly reduce stress marker expression (58). Furthermore, estradiol protected cells from apoptosis when challenged with an ER stress inducer, thapsigargin, in the presence of high glucose

**Table 1**  
Rodent models of sexual dimorphism with diabetes and/or hyperglycemia

Rodent model	Type of diabetes	Main features
ZDF rats, TALLYHO/JngJ mice	Polygenic	<ul style="list-style-type: none"> <li>• Males develop chronic hyperglycemia, females remain normoglycemic (63,64)</li> </ul>
OLETF rats	Polygenic	<ul style="list-style-type: none"> <li>• Males develop chronic hyperglycemia, females remain normoglycemic</li> <li>• Females increase beta-cell mass after partial pancreatectomy</li> <li>• This compensatory effect is lost after ovariectomy and restored by estradiol replacement (65,66)</li> </ul>
NZO mice	Polygenic	<ul style="list-style-type: none"> <li>• Males and females develop obesity when given a high-fat diet</li> <li>• Males develop chronic hyperglycemia, females remain normoglycemic unless ovariectomized</li> <li>• Male diabetic mice have an impaired cardiac function compared with normoglycemic females (67,68)</li> </ul>
Wfs1 knockout mouse (Wfs1 <sup>-/-</sup> )	Monogenic	<ul style="list-style-type: none"> <li>• Hyperglycemia is less severe in female Wfs1<sup>-/-</sup> compared with age-matched males</li> </ul>
Ins2 <sup>+Akita</sup> mouse, ApoE <sup>-/-</sup> :Ins2 <sup>+Akita</sup> mouse	Monogenic	<ul style="list-style-type: none"> <li>• This phenotype is similar to that in humans with mutations in the Wfs1 gene (69,70)</li> <li>• Point mutation in one allele of the Ins2 gene (C96Y), which impairs formation of functional insulin</li> <li>• Female Ins2<sup>+Akita</sup> mice develop less severe hyperglycemia compared with males</li> <li>• Female ApoE<sup>-/-</sup>:Ins2<sup>+Akita</sup> mice are only transiently hyperglycemic (by 10 weeks they become normoglycemic); ovariectomy eliminates normalization of blood glucose levels (38,71)</li> </ul>
STZ-injected rodents	Chemically induced hyperglycemia	<ul style="list-style-type: none"> <li>• STZ is a naturally occurring alkylating antineoplastic agent that is selectively toxic to pancreatic beta cells</li> <li>• Female rodents are more resistant to the toxic effects of STZ (higher concentrations and/or more frequent injections are required to induce diabetes in females) (43,72,73)</li> </ul>
Alloxan-injected rodents	Chemically induced hyperglycemia	<ul style="list-style-type: none"> <li>• Alloxan is a toxic analogue of glucose that has been used to selectively eliminate pancreatic beta cells</li> <li>• Administration of estrogen in male rats protects against alloxan-induced hyperglycemia (74)</li> </ul>

Apo, apolipoprotein; Ins, insulin; NZO, New Zealand Obese; OLETF, Otsuka Long-Evans Tokushima Fatty (rats); STZ, streptozotocin; Wfs, Wolfram syndrome; ZDF, Zucker Diabetic Fatty (rats).

concentration (58). The results of the study suggested that all 3 estrogen receptors play a role in signalling (58). Another study using INS-1 cells and isolated murine pancreatic islets demonstrated that estradiol protects beta cells from apoptosis when exposed to high levels of glucose (59). Ovariectomized rats treated with glucosamine, a supplement that in high doses can cause ER stress, had significantly reduced levels of insulin in their pancreatic islets as well as increased glucosamine-induced beta-cell apoptosis when compared with nonovariectomized controls. This supports a protective role for estrogens against beta-cell ER stress (60). Similarly, improved beta-cell function and health was observed in ovariectomized females and male hyperglycemic *Ins2<sup>+/-</sup>/Akita* mice receiving estrogen treatment (61).

Accumulating evidence is highlighting the importance of ER homeostasis in the maintenance and function of healthy beta cells. Given the systemic effects of estrogens and the multifaceted nature of intracellular hormone signalling, it is of high importance to conduct further research that can identify how sex hormones can affect relevant pathways and mechanisms that regulate beta-cell health.

## Conclusions

Despite the strong evidence for sexual dimorphism in diseases, such as diabetes, and its cardiovascular complications in humans as well as animal models, these differences remain poorly studied and represent a critical research gap. Moreover, preclinical studies are still predominantly conducted using male animal models and male-derived cell lines, resulting in an accumulation of incomplete, sex-biased data (62). It is, therefore, critical to address this gap in both clinical and preclinical research conducted in these fields so we may gain better understanding of sex differences in diabetes-associated cardiometabolic diseases and the impact of menopause, aging and hormone replacement therapies on health outcomes. This may lead to more targeted, effective and sex-specific management and prevention of cardiometabolic diseases.

## Acknowledgments

This work was supported by the Canadian Institutes of Health Research (PJT-166092) and the Heart and Stroke Foundation of Canada (G-17-0017029). G.H.W. is supported by an Ontario Mid-Career Investigator Award from the Heart and Stroke Foundation of Canada.

## Author Disclosures

Conflicts of interest: None.

## Author Contributions

MDP and GHW wrote and edited the manuscript.

## References

- Cnop M, Welsh N, Jonas J-C, Jörns A, Lenzen S, Eizirik DL. Mechanisms of pancreatic beta-cell death in type 1 and type 2 diabetes: Many differences, few similarities. *Diabetes* 2005;54(Suppl. 2):S97–107.
- Fowler MJ. Microvascular and macrovascular complications of diabetes. *Clin Diabetes* 2008;26:77–82.
- Emerging Risk Factors Collaboration, Di Angelantonio E, Kaptoge S, Wormser D, et al. Association of cardiometabolic multimorbidity with mortality. *JAMA* 2015;314:52–60.
- Booth GL, Kapral MK, Fung K, Tu JV. Relation between age and cardiovascular disease in men and women with diabetes compared with non-diabetic people: A population-based retrospective cohort study. *Lancet* 2006;368:29–36.
- Narayan KMV, Boyle JP, Thompson TJ, Sorensen SW, Williamson DF. Lifetime risk for diabetes mellitus in the United States. *JAMA* 2003;290:1884–90.
- Lehto S, Rönkämaa T, Pyörälä K, Laakso M. Cardiovascular risk factors clustering with endogenous hyperinsulinaemia predict death from coronary heart disease in patients with Type II diabetes. *Diabetologia* 2000;43:148–55.
- Soedamah-Muthu SS, Fuller JH, Mulnier HE, Raleigh VS, Lawrenson RA, Colhoun HM. High risk of cardiovascular disease in patients with type 1 diabetes in the U.K.: A cohort study using the general practice research database. *Diabetes Care* 2006;29:798–804.
- Patterson CC, Dahlquist GG, Gyürüs E, Green A, Soltész G. Incidence trends for childhood type 1 diabetes in Europe during 1989–2003 and predicted new cases 2005–20: A multicentre prospective registration study. *Lancet* 2009;373:2027–33.
- Kaiser AB, Zhang N, Pluijm WVD. Global prevalence of type 2 diabetes over the next ten years (2018–2028). *Diabetes* 2018;67:202–LB.
- Anja B, Laura R. The cost of diabetes in Canada over 10 years: Applying attributable health care costs to a diabetes incidence prediction model. *Health Promot Chronic Dis Prev Can Res Policy Pract* 2017;37:49–53.
- Rowley WR, Bezold C, Arikian Y, Byrne E, Krohe S. Diabetes 2030: Insights from yesterday, today, and future trends. *Popul Health Manag* 2017;20:6–12.
- Authors/Task Force Members, Rydén L, Grant PJ, Anker SD, et al. ESC guidelines on diabetes, pre-diabetes, and cardiovascular diseases developed in collaboration with the EASD: The Task Force on Diabetes, Pre-diabetes, and Cardiovascular Diseases of the European Society of Cardiology (ESC) and developed in collaboration with the European Association for the Study of Diabetes (EASD). *Eur Heart J* 2013;34:3035–87.
- Gale EA, Gillespie KM. Diabetes and gender. *Diabetologia* 2001;44:3–15.
- Beeson PB. Age and sex associations of 40 autoimmune diseases. *Am J Med* 1994;96:457–62.
- Wild S, Roglic G, Green A, Sicree R, King H. Global prevalence of diabetes: Estimates for the year 2000 and projections for 2030. *Diabetes Care* 2004;27:1047–53.
- Wändell PE, Carlsson AC. Gender differences and time trends in incidence and prevalence of type 2 diabetes in Sweden—a model explaining the diabetes epidemic worldwide today? *Diabetes Res Clin Pract* 2014;106:e90–2.
- Choi YJ, Kim HC, Kim HM, Park SW, Kim J, Kim DJ. Prevalence and management of diabetes in Korean adults: Korea National Health and Nutrition Examination Surveys 1998–2005. *Diabetes Care* 2009;32:2016–20.
- Ha KH, Kim DJ. Trends in the diabetes epidemic in Korea. *Endocrinol Metab* 2015;30:142–6.
- Lebovitz HE, Banerji MA. Ketosis-prone diabetes (Flatbush diabetes): An emerging worldwide clinically important entity. *Curr Diab Rep* 2018;18:120.
- The Emerging Risk Factors Collaboration. Diabetes mellitus, fasting blood glucose concentration, and risk of vascular disease: A collaborative meta-analysis of 102 prospective studies. *Lancet* 2010;375:2215–22.
- Margolis KL, Bonds DE, Rodabough RJ, et al. Effect of oestrogen plus progestin on the incidence of diabetes in postmenopausal women: Results from the Women's Health Initiative Hormone Trial. *Diabetologia* 2004;47:1175–87.
- Kanaya AM, Herrington D, Vittinghoff E, et al. Glycemic effects of postmenopausal hormone therapy: The Heart and Estrogen/Progestin Replacement Study: A randomized, double-blind, placebo-controlled trial. *Ann Intern Med* 2003;138:1–9.
- Gregg EW, Gu Q, Cheng YJ, Narayan KMV, Cowie CC. Mortality trends in men and women with diabetes, 1971 to 2000. *Ann Intern Med* 2007;147:149–55.
- Colhoun H. Coronary heart disease in women: Why the disproportionate risk? *Curr Diab Rep* 2006;6:22–8.
- Anand SS, Islam S, Rosengren A, et al. Risk factors for myocardial infarction in women and men: Insights from the INTERHEART study. *Eur Heart J* 2008;29:932–40.
- Peters SAE, Huxley RR, Woodward M. Diabetes as risk factor for incident coronary heart disease in women compared with men: A systematic review and meta-analysis of 64 cohorts including 858,507 individuals and 28,203 coronary events. *Diabetologia* 2014;57:1542–51.
- Huxley R, Barzi F, Woodward M. Excess risk of fatal coronary heart disease associated with diabetes in men and women: Meta-analysis of 37 prospective cohort studies. *BMJ* 2006;332:73–8.
- Peters SAE, Huxley RR, Woodward M. Diabetes as a risk factor for stroke in women compared with men: A systematic review and meta-analysis of 64 cohorts, including 775 385 individuals and 12 539 strokes. *Lancet* 2014;383:1793–80.
- Renzhe C, Hiroyasu I, Kazumasa Y, et al. Diabetes mellitus and risk of stroke and its subtypes among Japanese. *Stroke* 2011;42:2611–4.
- Abbott RD, Donahue RP, Kannel WB, Wilson PWF. The impact of diabetes on survival following myocardial infarction in men vs women: The Framingham Study. *JAMA* 1988;260:3456–60.
- Roche MM, Wang PP. Sex differences in all-cause and cardiovascular mortality, hospitalization for individuals with and without diabetes, and patients with diabetes diagnosed early and late. *Diabetes Care* 2013;36:2582–90.
- Eriksson M, Carlberg B, Eliasson M. The disparity in long-term survival after a first stroke in patients with and without diabetes persists: The Northern Sweden MONICA study. *Cerebrovasc Dis* 2012;34:153–60.
- Ballotari P, Ranieri SC, Luberto F, et al. Sex differences in cardiovascular mortality in diabetics and nondiabetic subjects: A population-based study (Italy). *Int J Endocrinol* 2015;2015:914057.
- NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in diabetes since 1980: A pooled analysis of 751 population-based studies with 4.4 million participants. *Lancet* 2016;387:1513–30.



35. Ferrannini E. Physiological and metabolic consequences of obesity. *Metabolism* 1995;44:15–7.
36. Rosano DGM, Vitale C, Marazzi G, Volterrani M. Menopause and cardiovascular disease: The evidence. *Climacteric* 2007;10:19–24.
37. Ormazabal V, Nair S, Elfeky O, Aguayo C, Salomon C, Zúñiga FA. Association between insulin resistance and the development of cardiovascular disease. *Cardiovasc Diabetol* 2018;17.
38. Venegas-Pino DE, Wang P-W, Stoute HK, et al. Sex-specific differences in an ApoE(-/-):Ins2(+)/Akita mouse model of accelerated atherosclerosis. *Am J Pathol* 2016;186:67–77.
39. Rettberg JR, Yao J, Brinton RD. Estrogen: A master regulator of bioenergetic systems in the brain and body. *Front Neuroendocrinol* 2014;35:8–30.
40. Alonso-Magdalena P, Ropero AB, Carrera MP, et al. Pancreatic insulin content regulation by the estrogen receptor ER $\alpha$ . *PLoS One* 2008;3:e2069.
41. Santos RS, Batista TM, Camargo RL, et al. Lack of estradiol reduces insulin exocytosis from pancreatic  $\beta$ -cells and increases hepatic insulin degradation. *Steroids* 2016;114:16–24.
42. Weigt C, Hertrampf T, Flenker U, et al. Effects of estradiol, estrogen receptor subtype-selective agonists and genistein on glucose metabolism in leptin resistant female Zucker Diabetic Fatty (ZDF) rats. *J Steroid Biochem Mol Biol* 2015;154:12–22.
43. Le May C, Chu K, Hu M, et al. Estrogens protect pancreatic beta-cells from apoptosis and prevent insulin-deficient diabetes mellitus in mice. *Proc Natl Acad Sci USA* 2006;103:9232–7.
44. Yamabe N, Kang KS, Zhu BT. Beneficial effect of 17 $\beta$ -estradiol on hyperglycemia and islet  $\beta$ -cell functions in a streptozotocin-induced diabetic rat model. *Toxicol Appl Pharmacol* 2010;249:76–85.
45. de Souza CF, Stopa LRS, Santos GF, et al. Estradiol protects against ovariectomy-induced susceptibility to the anabolic effects of glucocorticoids in rats. *Life Sci* 2019;218:185–96.
46. Tian JP, Delghingaro-Augusto V, Le May C, et al. Estrogen receptor activation reduces lipid synthesis in pancreatic islets and prevents  $\beta$  cell failure in rodent models of type 2 diabetes. *J Clin Invest* 2011;121:3331–42.
47. Yuchi Y, Cai Y, Legein B, et al. Estrogen receptor  $\alpha$  regulates  $\beta$ -cell formation during pancreas development and following injury. *Diabetes* 2015;64:3218–28.
48. Alonso-Magdalena P, Ropero AB, García-Arévalo M, et al. Antidiabetic actions of an estrogen receptor  $\beta$  selective agonist. *Diabetes* 2013;62:2015–25.
49. Liu S, Le May C, Wong WPS, et al. Importance of extranuclear estrogen receptor- $\alpha$  and membrane G protein-coupled estrogen receptor in pancreatic islet survival. *Diabetes* 2009;58:2292–302.
50. Belch J, MacCuish A, Campbell I, et al. The prevention of progression of arterial disease and diabetes (POPADAD) trial: Factorial randomised placebo controlled trial of aspirin and antioxidants in patients with diabetes and asymptomatic peripheral arterial disease. *BMJ* 2008;337:a1840.
51. Lonn E, Bosch J, Yusuf S, et al. Effects of long-term vitamin E supplementation on cardiovascular events and cancer: A randomized controlled trial. *JAMA* 2005;293:1338–47.
52. Lipson KL, Fonseca SG, Ishigaki S, et al. Regulation of insulin biosynthesis in pancreatic beta cells by an endoplasmic reticulum-resident protein kinase IRE1. *Cell Metab* 2006;4:245–54.
53. Scheuner D, Mierde DV, Song B, et al. Control of mRNA translation preserves endoplasmic reticulum function in beta cells and maintains glucose homeostasis. *Nat Med* 2005;11:757.
54. Back SH, Kaufman RJ. Endoplasmic reticulum stress and type 2 diabetes. *Annu Rev Biochem* 2012;81:767–93.
55. Wu J, Kaufman RJ. From acute ER stress to physiological roles of the unfolded protein response. *Cell Death Differ* 2006;13:374–84.
56. Lindholm D, Korhonen L, Eriksson O, Köks S. Recent insights into the role of unfolded protein response in ER stress in health and disease. *Front Cell Dev Biol* 2017;5:48.
57. Ozcan L, Tabas L. Role of endoplasmic reticulum stress in metabolic disease and other disorders. *Annu Rev Med* 2012;63:317–28.
58. Kooptiwut S, Mahawong P, Hanchang W, et al. Estrogen reduces endoplasmic reticulum stress to protect against glucotoxicity induced-pancreatic  $\beta$ -cell death. *J Steroid Biochem Mol Biol* 2014;139:25–32.
59. Kooptiwut S, Kaewin S, Semprasert N, et al. Estradiol prevents high glucose-induced  $\beta$ -cell apoptosis by decreased BTG2 expression. *Sci Rep* 2018;8:12256.
60. Kang L, Chen C-H, Wu M-H, Chang J-K, Chang F-M, Cheng J-T. 17 $\beta$ -Estradiol protects against glucosamine-induced pancreatic  $\beta$ -cell dysfunction. *Meno-pause* 2014;21:1239.
61. Xu B, Allard C, Alvarez-Mercado AI, et al. Estrogens promote misfolded pro-insulin degradation to protect insulin production and delay diabetes. *Cell Rep* 2018;24:181–96.
62. Reusch JEB, Kumar TR, Regensteiner JG, et al. Identifying the critical gaps in research on sex differences in metabolism across the life span. *Endocrinology* 2018;159:9–19.
63. Yokoi N, Hoshino M, Hidaka S, et al. A novel rat model of type 2 diabetes: The Zucker fatty diabetes mellitus ZFDM rat. *J Diabetes Res* 2013;2013:103731.
64. Kim JH, Stewart TP, Soltani-Bejnood M, et al. Phenotypic characterization of polygenic type 2 diabetes in TALLYHO/Jngj mice. *J Endocrinol* 2006;191:437–46.
65. Kawano K, Hirashima T, Mori S, Natori T. OLETF (Otsuka Long-Evans Tokushima Fatty) rat: A new NIDDM rat strain. *Diabetes Res Clin Pract* 1994;24(Suppl.): S317–20.
66. Zhu M, Mizuno A, Kuwajima M, et al. Ovarian hormone-induced beta-cell hypertrophy contributes to the homeostatic control of beta-cell mass in OLETF female rat, a model of type II diabetes. *Diabetologia* 1998;41:799–805.
67. John C, Grune J, Ott C, et al. Sex differences in cardiac mitochondria in the New Zealand obese mouse. *Front Endocrinol* 2018;9.
68. Vogel H, Mirhashemi F, Liehl B, et al. Estrogen deficiency aggravates insulin resistance and induces  $\beta$ -cell loss and diabetes in female New Zealand obese mice. *Horm Metab Res* 2013;45:430–5.
69. Noormets K, Köks S, Muldmaa M, Mauring L, Vasar E, Tillmann V. Sex differences in the development of diabetes in mice with deleted wolframin (Wfs1) gene. *Exp Clin Endocrinol* 2011;119:271–5.
70. Smith CJA, Crock PA, King BR, Meldrum CJ, Scott RJ. Phenotype-genotype correlations in a series of Wolfram syndrome families. *Diabetes Care* 2004;27:2003–9.
71. Yoshioka M, Kayo T, Ikeda T, Koizumi A. A novel locus, Mody4, distal to D7Mit189 on chromosome 7 determines early-onset NIDDM in nonobese C57BL/6 (Akita) mutant mice. *Diabetes* 1997;46:887–94.
72. Paik SG, Michelis MA, Kim YT, Shin S. Induction of insulin-dependent diabetes by streptozotocin. Inhibition by estrogens and potentiation by androgens. *Diabetes* 1982;31:724–9.
73. Li Y, Huang J, Yan Y, et al. Preventative effects of resveratrol and estradiol on streptozotocin-induced diabetes in ovariectomized mice and the related mechanisms. *PLoS One* 2018;13:e0204499.
74. Hamden K, Carreau S, Boujbiha MA, et al. Hyperglycaemia, stress oxidant, liver dysfunction and histological changes in diabetic male rat pancreas and liver: Protective effect of 17 $\beta$ -estradiol. *Steroids* 2008;73:495–501.