

Machine Learning Homework Sheet 11

Clustering

1 Gaussian Mixture Model

Problem 1: Consider a mixture of K Gaussians

$$p(\mathbf{x}) = \sum_k \pi_k \mathcal{N}(\mathbf{x} \mid \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k).$$

Derive the expected value $\mathbb{E}[\mathbf{x}]$ and the covariance $\text{Cov}[\mathbf{x}]$.

Hint: it is helpful to remember the identity $\text{Cov}[\mathbf{x}] = \mathbb{E}[\mathbf{x}\mathbf{x}^T] - \mathbb{E}[\mathbf{x}]\mathbb{E}[\mathbf{x}]^T$.

For $\mathbb{E}[\mathbf{x}]$ we use the law of iterated expectations

$$\begin{aligned} \mathbb{E}[\mathbf{x}] &= \mathbb{E}_{p(\mathbf{z})}[\mathbb{E}_{p(\mathbf{x}|\mathbf{z})}[\mathbf{x}|\mathbf{z}]] \\ &= \sum_{k=1}^K \pi_k \mathbb{E}_{p(\mathbf{x}|\mathbf{z})}[\mathbf{x}|\mathbf{z}] \end{aligned}$$

plugging in the mean of multivariate Gaussian, we get

$$= \sum_{k=1}^K \pi_k \boldsymbol{\mu}_k$$

For covariance, we first compute $\mathbb{E}[\mathbf{x}\mathbf{x}^T]$ again using the law of iterated expectations

$$\begin{aligned} \mathbb{E}[\mathbf{x}\mathbf{x}^T] &= \mathbb{E}_{p(\mathbf{z})}[\mathbb{E}_{p(\mathbf{x}|\mathbf{z})}[\mathbf{x}\mathbf{x}^T \mid \mathbf{z}]] \\ &= \sum_k \pi_k \mathbb{E}_{p(\mathbf{x}|\mathbf{z})}[\mathbf{x}\mathbf{x}^T \mid \mathbf{z}] \end{aligned}$$

plugging in the covariance of multivariate Gaussian, we get

$$= \sum_{k=1}^K \pi_k (\boldsymbol{\Sigma}_k + \boldsymbol{\mu}_k \boldsymbol{\mu}_k^T)$$

and thus

$$\begin{aligned} \text{Cov}[\mathbf{x}] &= \mathbb{E}[\mathbf{x}\mathbf{x}^T] - \mathbb{E}[\mathbf{x}]\mathbb{E}[\mathbf{x}]^T \\ &= \sum_{k=1}^K \pi_k (\boldsymbol{\Sigma}_k + \boldsymbol{\mu}_k \boldsymbol{\mu}_k^T) - \sum_{k=1}^K \sum_{j=1}^K \pi_k \pi_j \boldsymbol{\mu}_k \boldsymbol{\mu}_j^T \end{aligned}$$

Upload a single PDF file with your solution to Moodle by 27.01.2019, 11:59pm CET. We recommend to typeset your solution (using L^AT_EX or Word), but handwritten solutions are also accepted.

If your handwritten solution is illegible, it won't be graded and you waive your right to dispute that.

Problem 2: Consider a mixture of K isotropic Gaussians, all with the same *known* covariances $\Sigma_k = \sigma^2 \mathbf{I}$. Derive the EM algorithm for the case when $\sigma^2 \rightarrow 0$, and show that it's equivalent to Lloyd's algorithm for K -means.

We consider a GMM with identical, isotropic covariances. In that case, the responsibilities take the following form:

$$p(z_{ik} = 1 \mid \mathbf{x}, \theta) = \frac{p(\mathbf{x} \mid z_{ik} = 1, \boldsymbol{\mu}_k, \Sigma_k) p(z_{ik} = 1 \mid \pi_k)}{\int p(\mathbf{x} \mid \mathbf{z}) p(\mathbf{z}) d\mathbf{z}} \quad (1)$$

$$= \frac{\pi_k \mathcal{N}(\mathbf{x} \mid \boldsymbol{\mu}_k, \Sigma_k)}{\sum_{l=1}^K \pi_l \mathcal{N}(\mathbf{x} \mid \boldsymbol{\mu}_l, \Sigma_l)} \quad (2)$$

$$= \frac{\pi_k \exp\left(\frac{-\|\mathbf{x} - \boldsymbol{\mu}_k\|^2}{2\sigma^2}\right)}{\sum_l \pi_l \exp\left(\frac{-\|\mathbf{x} - \boldsymbol{\mu}_l\|^2}{2\sigma^2}\right)} \quad (3)$$

$$= \frac{1}{\sum_l \frac{\pi_l}{\pi_k} \exp\left(\frac{-\|\mathbf{x} - \boldsymbol{\mu}_l\|^2 + \|\mathbf{x} - \boldsymbol{\mu}_k\|^2}{2\sigma^2}\right)} \quad (4)$$

If $\boldsymbol{\mu}_k$ denotes the center that is closest to \mathbf{x} , then

$$\frac{-\|\mathbf{x} - \boldsymbol{\mu}_l\|^2 + \|\mathbf{x} - \boldsymbol{\mu}_k\|^2}{2\sigma^2} \leq 0$$

for all l , with equality if and only if $k = l$. For $\sigma \rightarrow 0$, the denominator of Equation 4 converges to 1: If $k = l$, the argument of $\exp(\cdot)$ is exactly zero, while for $k \neq l$ we are exponentiating increasingly negative arguments.

If $\boldsymbol{\mu}_k$ denotes a center that is *not* closest to \mathbf{x} , there is at least one $l \neq k$ for which

$$0 < \frac{-\|\mathbf{x} - \boldsymbol{\mu}_l\|^2 + \|\mathbf{x} - \boldsymbol{\mu}_k\|^2}{2\sigma^2} \rightarrow \infty, \quad \text{as } \sigma \rightarrow 0.$$

Consequently, the denominator of Equation 4 converges to ∞ , too.

This means that the responsibilities degenerate to a hard one-hot assignment of the data point \mathbf{x} to the component closest to \mathbf{x} . This coincides with step 1 of Lloyd's algorithm.

Inserting one-hot responsibilities into the general GMM M-step immediately yields step 2 in Lloyd's algorithm. Notice that we do not learn covariances, they are assumed fixed. Moreover, we don't have to worry about π_k 's, because they are irrelevant as the term π_l/π_k always gets overshadowed by the $\exp(\cdot)$ next to it.

We can conclude that Lloyd's algorithm for K-Means is a special case of the more general EM algorithm for GMM.

Problem 3: The dataset displayed on the right has been generated using a Gaussian mixture model with $K = 4$ components, each with its own mean $\boldsymbol{\mu}_k$ and covariance matrix Σ_k .


Match the covariance matrices in the table on the left with their corresponding Gaussian components in

Upload a single PDF file with your solution to Moodle by 27.01.2019, 11:59pm CET. We recommend to typeset your solution (using L^AT_EX or Word), but handwritten solutions are also accepted.

If your handwritten solution is illegible, it won't be graded and you waive your right to dispute that.

the plot on the right. Explain each of the answers with 1 sentence.

Σ_k	Cluster
$\begin{bmatrix} 2 & -1.7 \\ -1.7 & 2 \end{bmatrix}$	
$\begin{bmatrix} 0.9 & -0.8 \\ -0.8 & 1.2 \end{bmatrix}$	
$\begin{bmatrix} 3 & 0 \\ 0 & 0.5 \end{bmatrix}$	
$\begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix}$	



1. plot (a) - Has same shape as 2 (plot (c)), but has higher variance in all directions.
2. plot (c) - Has same shape as 1 (plot (a)), but has lower variance in all directions.
3. plot (b) - Aligned with the axes (offdiagonal entries are 0); more variance along the x_1 dimension.
4. plot (d) - Isotropic Gaussian - same variance in all directions.

Problem 4:

- a) Given is the dataset displayed in the figure below. Apply the K-means algorithm to this data using $K = 2$ and using the circled points as initial centroids.




Figure 1: K-Means Dataset

What are the clusters after K-Means converges? Draw your solution in the figure above, i.e. mark the location of the centroids with \times 's and show the clusters by drawing two bounding boxes around the points assigned to each cluster.

How many iterations did it take for K-Means to converge in the above problem?

One.

- b) Provide a different initialization, for which the algorithm will take **more** iterations to converge to the **same** solution. Make sure that your initialization does not lead to ties. Draw your initialization in the figure below.




Figure 2: Provide your initialization