

# Exercise

07

TUM Department of Informatics

<b>Supervised by</b>	Prof. Dr. Stephan Günnemann Informatics 3 - Professorship of Data Mining and Analytics
<b>Submitted by</b>	Marcel Bruckner (03674122) Julian Hohenadel (03673879) Kevin Bein (03707775)
<b>Submission date</b>	Munich, November 29, 2019

## Constrained Optimization

### Problem 1:

Constraints:

$$\theta_1 + \theta_2 \leq 12 \implies \theta_2 \leq 12 - \theta_1 \quad (1)$$


$$-\theta_1 + 2\theta_2 \geq -3 \implies \theta_2 \geq \frac{\theta_1 - 3}{2} \quad (2)$$

$$-5\theta_1 + 3\theta_2 \leq -4 \implies \theta_2 \leq \frac{1}{3}(5\theta_1 - 4) \quad (3)$$

$$\theta_2 \geq 2 \quad (4)$$

$$\theta_1 \geq 2 \quad (5)$$

Plot: Axis along  $x$ -dimension:  $\theta_1$ , Axis along  $y$ -dimension:  $\theta_2$ , simply plot the functions (1) up to (5):



$f(\theta) = 2\theta_1 - 3\theta_2$  Minimizer and maximizer both need to be a corner vertex of the domain.

Simple testing against (2, 2), (7, 2), (9, 3) and (5, 7) shows the solution:

Minimizer  $\theta_{min} = (5, 7) f(\theta_{min}) = -11$

Maximizer  $\theta_{max} = (9, 3) f(\theta_{max}) = 9$

### Problem 2:

### Problem 3:

### Problem 4:

# Appendix

We confirm that the submitted solution is original work and was written by us without further assistance.  
Appropriate credit has been given where reference has been made to the work of others.

---

Munich, November 29, 2019, Signature Marcel Bruckner (03674122)

---

Munich, November 29, 2019, Signature Julian Hohenadel (03673879)

---

Munich, November 29, 2019, Signature Kevin Bein (03707775)