

¹ *METADATA FOR:*² **Demography of the understory herb *Heliconia acuminata* (Heliconiaceae) in an**
³ **experimentally fragmented tropical landscape**⁴ Emilio M. Bruna^{1,2,3}, María Uriarte⁴, Maria Rosa Darrigo³, Paulo Rubim³, Cristiane F.
⁵ Jurinitz³, Eric R. Scott¹, Osmaildo Ferreira da Silva³, & W. John Kress⁵⁶ ¹ Department of Wildlife Ecology and Conservation, University of Florida, PO Box 110430,
⁷ Gainesville, FL 32611-0430, USA⁸ ² Center for Latin American Studies, University of Florida, PO Box 115530, Gainesville, FL
⁹ 32611-5530, USA¹⁰ ³ Biological Dynamics of Forest Fragments Project, INPA-PDBFF, CP 478, Manaus, AM
¹¹ 69011-970, Brazil¹² ⁴ Department of Ecology, Evolution and Environmental Biology, Columbia University, 1200
¹³ Amsterdam Ave., New York, New York 10027, USA¹⁴ ⁵ Department of Botany, National Museum of Natural History, PO Box 37012, Smithsonian
¹⁵ Institution, PO Box 37012, Washington DC, USA¹⁶ *Corresponding author:* Emilio M. Bruna (embruna@ufl.edu)¹⁷ *Open Research Statement:* The data described here are available as Supporting Information
¹⁸ at [url to be added] and have been archived at the Dryad Digital Repository [url to be added].

19 The version of the code used to review, correct, and prepare this archive is at Zenodo [*url to*
20 *be added*]. The code used to prepare this manuscript – including all statistical summaries,
21 tables, and figures – has also been archived at Zenodo [*url to be added*]. Post-publication
22 updates to code and the data sets, along with other project-related information, can be
23 found in the HDP Github Repository (<https://github.com/BrunaLab/HeliconiaSurveys>).

24 ## NULL

25 **METADATA**

26 **I. CLASS I. Data Set Descriptors**

27 **A. Data set identity:** Demographic data from populations of the understory herb
28 *Heliconia acuminata* (Heliconiaceae) in an experimentally fragmented tropical landscape
29 (1998-2009).

30 **B. Data set identification code:**

- 31 1. Data set File 1: HDP_plots.csv
32 2. Data set File 2: HDP_survey.csv

33 **C. Data set description:**

34 **1. Originators:** Emilio M. Bruna, Department of Wildlife Ecology and Conservation,
35 University of Florida, PO Box 110430, Gainesville, FL 32611-0430, USA and Center for
36 Latin American Studies, University of Florida, PO Box 115530, Gainesville, FL 32611,
37 USA.

38 **2. Abstract:** Habitat fragmentation remains a major focus of research by ecologists
39 decades after being put forward as a threat to the integrity of ecosystems. While
40 studies have documented myriad biotic changes in fragmented landscapes, including
41 the local extinction of species from fragments, the demographic mechanisms underlying
42 these extinctions are rarely known. However, many of them – especially in lowland
43 tropical forests – are thought to be driven by one of two mechanisms: (1) reduced
44 recruitment in fragments resulting from changes in the diversity or abundance of
45 pollinators and seed dispersers or (2) increased rates of individual mortality in
46 fragments due to dramatically altered abiotic conditions, especially near fragment

edges. Unfortunately, there have been few tests of these potential mechanisms due to the paucity of long-term and comprehensive demographic data collected in both forest fragments and continuous forest sites. Here we report 11 years (1998-2009) of demographic data from populations of the Amazonian understory herb *Heliconia acuminata* (LC Rich.) found at Brazil's Biological Dynamics of Forest Fragments Project (BDFFP). The resulting data set comprises >66000 plant×year records of 8586 plants, including 3464 seedlings that became established after the initial census. Seven populations were in experimentally isolated fragments (one in each of four 1-ha fragments and one in each of three 10-ha fragments), with the remaining six populations in continuous forest. Each population was in a 50 × 100m permanent plot, with the distance between plots ranging from 500 m-60 km. The plants in each plot were censused annually, at which time we recorded, identified, marked, and measured new seedlings, identified any previously marked plants that died, and recorded the size of surviving individuals. Each plot was also surveyed 4-5 times during the flowering season to identify reproductive plants and record the number of inflorescences each produced. These data have been used to investigate topics ranging from the way fragmentation-related reductions in germination influence population dynamics to statistical methods for analyzing reproductive rates. This breadth of prior use reflects the value of these data to future researchers. In addition to analyses of plant responses to habitat fragmentation, these data can be used to address fundamental questions in plant demography, the evolutionary ecology of tropical plants, and for developing and testing demographic models and tools. Though we welcome opportunities to collaborate with interested users, there are no restrictions on the use this data set. However, we do request that those using the data for teaching or research inform us of how they are doing so and cite this paper and the data archive when appropriate. Any publication using the data must also include a BDFFP Technical Series Number in the Acknowledgments. Authors can request this series number upon the acceptance of their

74 article by contacting the BDFFP's Scientific Coordinator or E. M. Bruna.

75 **D. Key words:** Amazon, Brazil, deforestation, demography, edge effects, flowering, forest
76 fragments, habitat fragmentation, integral projection models, matrix models, population
77 dynamics, vital rates.

78 **CLASS II. RESEARCH ORIGIN DESCRIPTORS**

79 **A. Overall project description:**

80 **1. Identity:** The *Heliconia* Demography Project

81 **2. Originators:** Emilio M. Bruna, W. John Kress, and María Uriarte

82 **3. Period of study:** 1998-2009

83 **4. Objectives:** Habitat fragmentation remains a major focus of research by ecologists
84 (Didham et al. 2012, Haddad et al. 2015, Brudvig et al. 2017, Resasco et al. 2017,
85 Fletcher et al. 2018) decades after it was first put forward as a threat to the integrity
86 of ecosystems (Harris 1984, Wilcove et al. 1986). Studies have documented myriad
87 biotic changes in fragmented landscapes, including the local extinction of species from
88 fragments (Harrison and Bruna 1999, Laurance et al. 2011). The demographic
89 mechanisms underlying these extinctions are rarely known (Bruna et al. 2009).

90 However, many of them – especially in lowland tropical forests – are thought to be
91 driven by one of two mechanisms: (1) reduced recruitment in fragments resulting from
92 changes in the diversity or abundance of specialized pollinators and seed dispersers
93 (Murcia 1996, Silva and Tabarelli 2000), or (2) increased rates of individual mortality
94 in fragments (Laurance et al. 1998, Zartman et al. 2015) due to dramatically altered
95 abiotic conditions, especially near fragment edges (reviewed in Broadbent et al. 2008).
96 Unfortunately, tests of these potential mechanisms remain limited due to the paucity of
97 long-term demographic data collected in both forest fragments and continuous forest

98 sites (Bruna et al. 2009).

99 Most studies investigating the effects of forest fragmentation on tropical plants
100 focus on trees (Cordeiro et al. 2009, Jurinitz et al. 2013, Zambrano and
101 Salguero-Gómez 2014), in part because they are major reservoirs of carbon (Slik
102 et al. 2010, Lasky et al. 2014). However, herbaceous species can comprise up to
103 30% of the plant species in lowland tropical forests (Gentry and Emmons 1987,
104 Ribeiro et al. 2010, Iannone and Vargas 2022, Spicer et al. 2022), where they are
105 habitat and food for myriad animal taxa as well as economically and culturally
106 important non-timber forest products (Nakazono et al. 2004, Athayde et al.
107 2006). Despite their biocultural importance, however, the way in which habitat
108 fragmentation and other global change phenomena influences the population
109 dynamics of tropical understory plants remains conspicuously understudied
110 (Bruna et al. 2009).

111 The *Heliconia* Demography Project (HDP) was established to address the lack of
112 data on the demography of understory plants in fragmented tropical landscapes.
113 The core of the HDP is annual censuses of thirteen populations of *Heliconia*
114 *acuminata* located in either continuous forest or experimentally isolated forest
115 fragments at Brazil's Biological Dynamiocs of Forest Fragments Project
116 (Laurance et al. 2011) The primary purpose behind their initial collection was to
117 parameterize size-structured demographic models (Caswell 2000, Ellner and Rees
118 2006) with which to (1) compare the demography and population dynamics of *H.*
119 *acuminata* populations in fragments and continuous forest, and (2) test the
120 hypothesis that reductions in seedling establishment in forest fragments would
121 lead to population declines. Simulations suggested extreme reductions in
122 recruitment were necessary for population declines (Bruna 2003), and empirical
123 estimates of seedling establishment in fragments were frequently below these

thresholds (Bruna 2002). Other studies demonstrated that the growth rates of plants in fragments, where abiotic conditions are often(Broadbent et al. 2008) severely altered, are much lower than in continuous forest (Bruna et al. 2002). Chronically reduced growth (Gagnon et al. 2011), especially of large plants, is a primary contributor to lower population growth rates in forest fragments (Bruna and Oli 2005). More recent projects using the data set described have has, for example, assessed the effects of local environmental conditions, disperser diversity, and disperser behavior on safe-site vs. seed-limitation (Uriarte et al. 2010, 2011), quantified population genetic structure (Côrtes et al. 2013), and compared statistical methods for modeling reproductive rates (Brooks et al. 2019).

5. Abstract: Habitat fragmentation remains a major focus of research by ecologists decades after being put forward as a threat to the integrity of ecosystems. While studies have documented myriad biotic changes in fragmented landscapes, including the local extinction of species from fragments, the demographic mechanisms underlying these extinctions are rarely known. However, many of them – especially in lowland tropical forests – are thought to be driven by one of two mechanisms: (1) reduced recruitment in fragments resulting from changes in the diversity or abundance of pollinators and seed dispersers or (2) increased rates of individual mortality in fragments due to dramatically altered abiotic conditions, especially near fragment edges. Unfortunately, there have been few tests of these potential mechanisms due to the paucity of long-term and comprehensive demographic data collected in both forest fragments and continuous forest sites. Here we report 11 years (1998-2009) of demographic data from populations of the Amazonian understory herb *Heliconia acuminata* (LC Rich.) found at Brazil's Biological Dynamics of Forest Fragments Project (BDFFP). The resulting data set comprises >66000 plant×year records of 8586 plants, including 3464 seedlings that became established after the initial census. Seven populations were in experimentally isolated fragments (one in each of four 1-ha

fragments and one in each of three 10-ha fragments), with the remaining six populations in continuous forest. Each population was in a 50 × 100m permanent plot, with the distance between plots ranging from 500 m-60 km. The plants in each plot were censused annually, at which time we recorded, identified, marked, and measured new seedlings, identified any previously marked plants that died, and recorded the size of surviving individuals. Each plot was also surveyed 4-5 times during the flowering season to identify reproductive plants and record the number of inflorescences each produced. These data have been used to investigate topics ranging from the way fragmentation-related reductions in germination influence population dynamics to statistical methods for analyzing reproductive rates. This breadth of prior use reflects the value of these data to future researchers. In addition to analyses of plant responses to habitat fragmentation, these data can be used to address fundamental questions in plant demography, the evolutionary ecology of tropical plants, and for developing and testing demographic models and tools. Though we welcome opportunities to collaborate with interested users, there are no restrictions on the use this data set. However, we do request that those using the data for teaching or research inform us of how they are doing so and cite this paper and the data archive when appropriate. Any publication using the data must also include a BDFFP Technical Series Number in the Acknowledgments. Authors can request this series number upon the acceptance of their article by contacting the BDFFP's Scientific Coordinator or E. M. Bruna.

6. Sources of funding: The initial establishment of plots and the 1998-2002 surveys were supported by grants to E. M. Bruna from the Smithsonian Institution (Graduate Student Research Award), the University of California, Davis (Center for Population Biology Graduate Research Grant, M. E. Mathias Graduate Research Grant), the Biological Dynamics of Forest Fragments Project (Graduate Student Logistics Grant), the National Science Foundation (Dissertation Improvement Grant INT 98-06351), and the Ford Foundation (Dissertation Year Fellowship). The 2001-2005 surveys were

178 supported a grant from the National Science Foundation to E. M. Bruna (Research
179 Starter Grant DEB-0309819). The 2006-2009 surveys were supported by grants from
180 the National Science Foundation to E. M. Bruna (DEB-0614149) and María Uriarte
181 (DEB-0614339). Subsequent analyses and the preparation of these data for archiving
182 were supported by the National Science Foundation (DEB-1948607).

183 **B. Subproject description**

184 **1. Site description**

185 a. **Site type:** Lowland tropical forest

186 b. **Geography:** The data were collected at the Biological Dynamics of Forest
187 Fragments Project (BDFFP, 2°30'S, 60°W), a 1000-km mosaic of lowland
188 forest, forest fragments, secondary forests, and pastures located
189 approximately 70 km north of Manaus, Amazonas, Brazil (Fig. 1).

190 c. **Habitat:** The BDFFP is dominated by tropical evergreen lowland forest
191 (i.e., ‘tropical moist forest’, *sensu* Holdridge (1967). The forest canopy at
192 the sites is ~35–40 m tall, with emergent trees of up to ~45 m
193 (Rankin-de-Mérona et al. 1992). The tree community at the BDFFP is
194 highly diverse: ~1300 species total (Laurance 2001), with as many as 280
195 tree species ha⁻¹ (Oliveira and Mori 1999). The understory is dominated by
196 stemless palms (Scariot 1999). All HDP plots are located in *terra-firme* (i.e.,
197 non-flooded) forest and none are bisected by streams.

198 d. **Geology:** Soils in the sites are nutrient-poor xanthic ferralsols, known as
199 yellow latosols in the Brazilian soil classification system. Despite their high
200 clay content they have poor water-retention capacity (Fearnside and
201 Leal-Filho 2001). The often rugged topography at the BDFFP ranges in

202 elevation from 50-150 m elevation (Gascon and Bierregaard 2001).

203 e. **Watersheds:** The BDFFP landscape includes catchments of the Urubu,
204 Cuieiras, and Preto da Eva rivers (Nessimian et al. 2008).

205 f. **Site history:** A complete history of the BDFFP can be found in Gascon
206 and Bierregaard (2001) and Bierregaard et al. (2002). Briefly, the BDFFP
207 reserves were established on three cattle ranches. Fragments were isolated
208 between 1980-1984 by felling the trees surrounding the patch of forest to be
209 isolated (Lovejoy et al. 1986). Fragment reserves were fenced to prevent the
210 incursion of cattle from the surrounding pastures, and the vegetation in a
211 100 m strip around each fragment is mechanically cleared every 5-6 years to
212 ensure fragments remain isolated (Gascon and Bierregaard 2001). The
213 structure and species composition of the secondary growth that surrounds a
214 fragment, which is strongly dependent on whether fire was used to clear
215 land prior to planting pasture grasses (Mesquita et al. 2001), can have large
216 effects on the species composition, ecological processes, and abiotic
217 conditions in fragments (reviewed in Laurance et al. 2002, 2011). The
218 BDFFP is currently administered collaboratively by the Smithsonian
219 Tropical Research Institute and Brazil's Instituto Nacional de Pesquisas da
220 Amazônia (INPA).

221 g. **Climate:** Mean annual temperature at the site is 26°C (range 19-39°C).
222 Annual rainfall ranges from 1900-2300 mm (Scott et al. 2022), with a
223 pronounced dry season from June-December in which there is <100 mm
224 rain per month.

225 2. Sampling Design

226 a. **Design characteristics:** Annual demographic surveys of *Heliconia*

227 *acuminata* populations were carried out in 13 permanent plots distributed
228 across the BDFFP landscape (Bruna and Kress 2002). Six plots are located
229 in continuous forest, four in 1-ha fragments, and three in 10-ha fragments
230 (one plot per fragment; Fig. 1).

231 *Heliconia acuminata* (Heliconiaceae) is a perennial, self-incompatible
232 monocot native to Amazonia (Kress 1990) and widely distributed
233 throughout the Amazon basin (Kress 1990). Although many species of
234 *Heliconia* grow in large aggregations on roadsides, gaps, and in other
235 disturbed habitats, others, including *H. acuminata*, grow primarily in
236 the shaded forest understory (Kress 1983). *Heliconia acuminata* is the
237 most abundant understory herb throughout most of the BDFFP
238 (Ribeiro et al. 2010); the other two *Heliconia* species found in the
239 BDFFP reserves are either very rare (*H. latispatha*) or restricted to
240 saturated soils adjacent to streams (*H. tarumaensis*).

241 Each *Heliconia acuminata* has a basal rhizome from which erect
242 vegetative shoots with broad leaves emerge. Plants grow very slowly –
243 an analysis based on the 1998–2006 survey data found that the average
244 change in shoot number was < 10% per year (averaged across all
245 plants), with negative growth rates (i.e., plants lost shoots) in three of
246 the years (Gagnon et al. 2011). The slow growth rate of *H. acuminata*
247 observed in demographic plots is similar to that observed in
248 experimental transplants (Bruna and Ribeiro 2005a). Stem loss is not
249 the result of herbivory. The primary foliar herbivores of *Heliconia*
250 species are Hispine beetles, whose larvae and adults perforate or scrape
251 the surface of unrolled immature leaves or feed on inflorescences
252 (Strong 1977). The beetle species associated with *H. acuminata* is

253 *Cephaloleia nigriceps* Baly (Staines and Garcia-Robledo 2014); it does
254 little damage to most leaves but can cause extensive damage to bracts,
255 flowers, and developing ovaries.

256 *Heliconia acuminata* flowers during the rainy season, and the likelihood
257 an individual will flower is strongly size size-dependent (Bruna 2002,
258 Bruna and Kress 2002 fig. 2). Reproductive plants produce one or
259 more flowering shoots, each of which terminates in single inflorescence
260 comprising red bracts (i.e., modified leaves) subtending white flowers.
261 Reproductive plants can produce multiple flowering shoots, but 75% of
262 the reproductive plants recorded in our demographic suveys produced
263 a single inflorescence (range = 1-7). Flowers remain open for a single
264 day, after which the style and perianth abscise and fall from the plant.
265 In 1998 we documented the phenology of N = 112 flowering plants
266 found along the trails of BDFFP Reserve #1501 (Bruna 2021) and
267 found that they produced an average of 20.62 ± 8.61 SD flowers each
268 (range: 0-48). Flowers remain open for a single day, after which the
269 style and perianth abscise and fall from the plant. Pollen transfer
270 experiments indicate self-compatibility is extremely low (E. M. Bruna
271 and M. R. Darrigo, *unpubl. data*).

272 *Heliconia* in the Americas are hummingbird-pollinated. In our field
273 sites *H. acuminata* is pollinated by the ‘traplining’ hummingbirds
274 *Phaeothornis superciliosus* and *P. bourcieri* (Stouffer and Bierregaard
275 1995, 1996), whose visitation rates to flowers are extremely low (<1
276 visit hour⁻¹, Bruna et al. 2004). This, coupled with damage by *C.*
277 *nigriceps* to the developing ovaries, results in low rates of fruit
278 production (Bruna and Kress 2002). A successfully pollinated flower

279 produces a fleshy blue fruit with up to 3 seeds (average = 1.90 ± 0.81
280 SD, N = 873 fruits, Bruna 2014).

281 *Heliconia* fruits are consumed by frugivorous birds, which then disperse
282 the seeds. In our study sites the primary dispersers of *Heliconia*
283 *acuminata* seeds are several species of manakin (*Pipra erythrocephala*,
284 *P. pipra*, *Lepidothrix serena*, *Schiffornis turdinus*, *Corapipo gutturalis*)
285 and the White-necked Thrush (*Turdus albicollis*) (Uriarte et al. 2011).
286 Seeds germinate 6-7 months after they are dispersed, which coincides
287 with the onset of the rainy season (Bruna 1999, 2002). Rates of seed
288 germination and seedling establishment in field experiments were
289 generally low, but they were significantly higher in continuous forest
290 than forest fragments (Bruna 1999, 2002). Experiments also indicate
291 that post-dispersal seed predation is negligible. While some seeds did
292 germinate >1 year after experimental dispersal, this was generally rare
293 – especially in fragments. These results are consistent with others
294 suggesting long-lived seed banks are rare in lowland tropical forests
295 (Vázquez-Yanes and Orozco-Segovia 1993), as well as observations that
296 understory *Heliconia* species recruit primarily via seeds and not clonal
297 spread (Stiles 1975).

298 *Heliconia acuminata* individuals are easily collected in the field and can
299 be grown in pots or transplanted directly into the ground (Bruna et al.
300 2002, Bruna and Ribeiro 2005a). Plants can be readily propagated for
301 use in transplant experiments (e.g., Bruna and Andrade 2011) by
302 segmenting the rhizome (Berry and Kress 1991). Seeds collected from
303 ripe fruits and treated using protocols described in Bruna (2002) can
304 be stored for weeks for use in experiments (e.g., Bruna 2002, Bruna

305 and Ribeiro 2005b) and they have high germination rates in field shade
306 houses. Finally, plants from the demographic surveys were used to
307 develop a library of microsatellite markers (Côrtes et al. 2009).

308 b. **Permanent Plots:** Each demographic plot is 50 × 100m and is subdivided
309 into 50 contiguous subplots of 10 × 10m to facilitate the surveys. Plots in
310 1-ha fragments were established in a randomly selected half of the fragment
311 (Fig. 3), plots in 10-ha fragments are located in the center of the fragment
312 (Fig. 4), and plots in continuous forest are located 500-4000 m from any
313 borders with cattle pastures or secondary forest (Fig. 5). The plots furthest
314 apart are from each other are separated by ~70 km.

315 In January 1997 we demarcated four demographic plots in fragments
316 (FF-3, FF-4, FF-6, FF-7) and one in a continuous forest (CF 1) to test
317 survey protocols and sample plants for genetic analyses. The remaining
318 plots were established in 1998, which was also when the first complete
319 census was conducted in all plots. To find and mark the plants, a team
320 of 2-3 people walked slowly through each subplot to locate any
321 *Heliconia acuminata*, which they then marked with a wooden stake to
322 which was attached an individually numbered aluminum tag. They
323 also recorded (1) how many vegetative shoots the plant had, and (2)
324 its height, measured as the distance from the ground to the top of the
325 tallest leaf (rounded to the nearest cm). Both of these size metrics are
326 highly correlated with the total leaf-area of a plant.

327 c. **Frequency of Data Collection:** Plots were censused annually at the
328 onset of the rainy season to coincide with seedling establishment (generally
329 late January to February). The exception to this was the three continuous
330 forest plots established in August 1998, which were censused in August 1999.

331 Regular visits were made to all 13 plots throughout the rainy season to
332 identify reproductive individuals and record the number of flowering shoots
333 and inflorescences that they had produced.

334 **3. Research Methods**

335 a. **Demographic Surveys:** During each census team members recorded
336 which plants died, the size (i.e., height and number of shoots) of all
337 surviving plants, and the size of all new seedlings, which were also marked
338 with a numbered tag. Survey team members also noted any new canopy
339 gaps created by fallen trees or limbs, estimated the proportion of any subplot
340 that was affected by a treefall, and recorded if plants were under treefalls or
341 damaged by fallen branches or palm fronds. These treefall records
342 (`treefall_impacts.csv` and `subplot_treefalls.csv`) are in the
343 `data/survey_clean` subfolder of the HDP Repository
344 (<https://github.com/BrunaLab/HeliconiaSurveys>).

345 b. **Taxonomy and systematics:** *Heliconia* is the only genus in the family
346 Heliconiaceae. This family is distinguished from the others in the order
347 Zingiberales by having inverted flowers, a single staminode, and drupaceous
348 fruits (Kress 1990). It is estimated that there are 200-250 species of
349 *Heliconia*, almost all of which are native to the Neotropics. *Heliconia*
350 *acuminata* L. C. (Rich.) (Richard 1831) is one of the approximately 20
351 *Heliconia* species found in the Brazilian Amazon (Kress 1990). We
352 deposited voucher specimens of *H. acuminata* collected in areas adjacent to
353 demographic plots at the herbaria of the Instituto Nacional de Pesquisas da
354 Amazônia (Accession Numbers INPA 189569-189573) and the University of
355 California, Davis (Accession Numbers DAV 69391-69396).

356 **4. Project personnel:** In addition to the Project Originators, other key personnel
357 include the Project Managers that were responsible for coordinating the annual
358 censuses and other field activities, BDFFP Technicians (“*Mateiros*”) that assisted with
359 data collection and provided logistical support in the field, and undergraduate and
360 postgraduate field assistants hired to assist with the surveys.

361 a. **Project Managers:** Paulo Rubim (2007-2012), Maria Beatriz Nogueira
362 (2002), Maria Rosa Darrigo (2002-2003), Cris Follman Jurinitz (2003),
363 Simone Benedet (2004).

364 b. **BDFFP Technicians:** Osmaildo Ferreira da Silva, Francisco Marques,
365 Alaercio Marajó dos Reis, João de Deus Fragata, Romeu Cardoso.

366 c. **Undergraduate & Postgraduate Field Assistants:** Olavo Nardy
367 (2000), Obed Garcia (2001), Sylvia Heredia (2001-2002), Maria Beatriz
368 Nogueira (2002), Cris Follman Jurinitz (2003), David M. Lapola (2003),
369 Denise Cruz (2003), Cristina Escate (2004), Bruno Turbiani (2005),
370 Elisabete Marques da Costa (2006), Wesley Dátilo da Cruz (2007),
371 Jefferson José Valsko da Silva (2007).

372 **CLASS III. DATA SET STATUS AND ACCESSIBILITY**

373 **A. Status**

374 **1. Latest updates:**

- 375 a. File 1 (`HDP_plots.csv`): 2023-06-08.
376 b. File 2 (`HDP_survey.csv`): 2023-06-08

377 **2. Latest archive date:** [date of archiving at Dryad to be added upon acceptance]

378 **3. Metadata status:** Complete (last update: NA)

379 **4. Data verification:** An extensive review of the data was conducted in preparation for
380 archiving. We began by generating a list of potential anomalies that could indicate
381 errors (e.g. extremely large changes in size from one year to the next, plants marked as
382 dead that had subsequent measurements), and then wrote code to search for these
383 anomalies using the R statistical programming language (R Core Development Team
384 2014). We also used the `pointblank` library (Iannone and Vargas 2022) to identify
385 cases in the data set for review and validation. All records flagged were evaluated by E.
386 M. Bruna by checking the values in the electronic records against the original data
387 sheets. Corrections to the data set were made using R scripts; the code documenting
388 and implementing these changes is archived at Zenodo [*url to be added upon
389 acceptance*]. Questions regarding the data set or code should be posted as `Issues` on
390 the HDP Repository (<https://github.com/BrunaLab/HeliconiaSurveys/issues>) or
391 referred to E. M. Bruna, who will investigate and update the database or code as
392 needed. Summaries of any post-publication updates will be posted to the `NEWS.md` file
393 of the HDP Github Repository (<https://github.com/BrunaLab/HeliconiaSurveys>),
394 with revised data sets assigned new version numbers based on the ‘Frictionless Data’
395 guidelines (<https://frictionlessdata.io/specs/patterns/>). This will allow users to
396 reference the version of the data used in their analyses to ensure their reproducibility
397 (reviewed in Yenni et al. 2019).

398 **B. Accessibility**

399 **1. Storage location and medium:** The data described here are available as Supporting
400 Information at [*url for ESA Archives to be added*] and have been archived at the Dryad
401 Digital Repository [*url to be added*]. The version of the code used to review, correct,
402 and prepare this archive is at Zenodo [*url to be added*]. The code used to prepare this
403 manuscript – including all statistical summaries, tables, and figures – has also been
404 archived at Zenodo [*url to be added*]. Post-publication updates to code and the data

405 sets, along with other project-related information, can be found in the HDP Github
406 Repository (<https://github.com/BrunaLab/HeliconiaSurveys>).

407 **2. Contact person:** Emilio M. Bruna, Department of Wildlife Ecology and
408 Conservation, Box 110430, Gainesville, FL 32611 USA. Phone: (352) 846-0634. Email:
409 embruna@ufl.edu

410 **3. Copyright restrictions:** None

411 **4. Proprietary restrictions:** None.

412 a. **Conditions of Reuse:** Though we welcome opportunities to collaborate
413 with interested users, there are no restrictions on the use this data set.
414 However, any publication using data collected at the BDFFP – including
415 this data set – must include a BDFFP Technical Series Number in the
416 Acknowledgments. Authors can request this series number upon the
417 acceptance of their article by contacting the BDFFP's Scientific
418 Coordinator (pdbff@inpa.gov.br) or E. M. Bruna (embruna@ufl.edu).

419 b. **Citation:** Authors of any publications or products using these data should
420 cite both this data paper and the Dryad data archive [*citation of Dryad*
421 *archive to be added upon acceptance*]. We also request that they provide E.
422 M. Bruna a copy of their article upon acceptance, which allows us to track
423 the data set's usage, inform users of any corrections or updates, report
424 articles using the data to the funding agencies that provided support, and
425 document the different ways in which the scientific community uses the data.

426 c. **Disclaimers:** While the data are provided in good faith and are accurate
427 to the best of our knowledge, they are provided “as is”. We do not assume
428 any legal liability or responsibility for their accuracy, completeness, or

429 utility. The responsibility for use and analysis of these data lies completely
430 with the user.

431 **5. Costs of acquiring data:** None.

432 **CLASS IV. DATA STRUCTURAL DESCRIPTORS**

433 **A. Data set File 1:** Descriptors of demographic plots

434 **1. Identity:** HDP_plots.csv

435 **2. Size:** 14 rows (including header), 407 Bytes

436 **3. Format and storage mode:** ASCII text, comma delimited. No compression scheme
437 used.

438 **4. Header information:** The first row of the file contains the variable names.

439 **5. Alphanumeric attributes:** Mixed

440 **6. Special Characters:** Missing values are represented with NA.

441 **7. Authentication Procedures:** checksum (MD5 of the file downloaded to computer
442 from the online repository: 7c39b8905ecef0a99c18ceb615ba6ca8.

443 **8. Start & End Columns:** Start: plot_id, End: yr_isolated

444 **9. Variable Information:** Each row is one plot, with the columns providing
445 plot-specific values for each variable (Table 1).

446 **B. Data set File 2:** *Heliconia* Demographic Data

447 **1. Identity:** HDP_survey.csv

448 **2. Size:** 66397 rows (including header), 3.61 MB

449 **3. Format and storage mode:** ASCII text, comma delimited. No compression scheme
450 used.

451 **4. Header information:** The first row of the file contains the variable names.

452 **5. Alphanumeric attributes:** Mixed.

453 **6. Special Characters:** Missing values are represented with NA.

454 **7. Authentication Procedures:** checksum (MD5 of the file downloaded to computer
455 from the online repository: 464706803cae7ea78aa6da8cc84124e0).

456 **8. Start & End Columns:** Start: `plot_id`, End: `tag_number`

457 **9. Data anomalies:** Plants that could not be found during a survey were recorded as
458 ‘missing’ but maintained on the survey list to be searched for in subsequent years. The
459 same is true of plants under branches or the crowns of fallen trees, which might not be
460 found for several years when the crown’s leaves dried and fell or the area under the
461 crown could be safely searched. The codes used to denote such cases are defined in
462 Table 2.

463 The stakes and numbered tags used to mark plants were sometimes displaced,
464 broken, or buried under leaf litter as a result of tree falls or other disturbances. If
465 a plant’s tag couldn’t be found after an extensive search, it would be marked
466 with a new tag. In some cases, it was straightforward to determine such a plant’s
467 original number when entering the survey data (e.g., when all plants in a
468 low-density subplot were found except one, which in the prior year was similar in
469 size as the plant found without a tag). In those cases, the plant’s prior
470 measurements were transferred to the new number and we logged the details of
471 the change in tag number; the `tag_changes.csv` file is available in the
472 `data/survey_clean` subfolder of the HDP Repository

(<https://github.com/BrunaLab/HeliconiaSurveys>). In other cases, it was impossible to definitively determine a plant's original number (e.g., when two similarly sized plants in a subplots were both missing their tags). In these cases, the original number was maintained in the database with the plant's status noted as 'missing' in subsequent surveys. The record for the new number indicates the plant with which it is associated is an established plant that was found without a tag (see Section IV, Table 2) and not a new seedling.

There were also cases in which established plants were found without tags in subplots where all previously tagged plants had already been located and measured, indicating previous survey teams had failed to find and mark them. These plants were marked, measured, and added to the database with a code indicating they were a established (i.e., post-seedling) but previously unmarked plant (See Table 2). Of the $N = 1562$ plants in the data set, 18.2% were found without tags after the plot had been established. Almost half of these (45%) were in the three plots where *H. acuminata* density was highest (CF-1, FF-7, CF-3).

Due to logistical or financial constraints, no surveys were conducted in plot CF-6 in 2003; in plots CF-4, CF-5, and CF-6 in 2000; or plots FF-5, FF-6, and FF-7 in 2008-2009.

10. **Variable information:** Each row in the data set is a demographic plot, with columns of data describing that plot (Table 2). Blanks do not denote missing information, but rather nothing relevant to report.

494 CLASS V. SUPPLEMENTAL DESCRIPTORS

495 A. Data acquisition:

1. **Data forms:** Examples of the forms used to collect survey data can be found at

497 `survey_records.md` in the `docs/survey_records/survey_sheets` subfolder of the
498 HDP Github Repository (<https://github.com/BrunaLab/HeliconiaSurveys>).

499 **2. Location of original data forms, electronic files, and archived copies:** Original
500 data sheets are stored at the University of Florida. Scanned copies of the data sheets
501 (in `.pdf` format) and the electronic copies of the data (in `.csv` format) are stored on a
502 desktop computer at the University of Florida that is backed up daily to two portable
503 hard drives and two cloud storage services. The integrity of digital files is verified
504 semi-annually.

505 **3. Data entry verification procedures:** Surveys were typically conducted in field trips
506 of 7-14 days with a 2-4 day interval in Manaus. During each interval we made backup
507 paper copies of the data sheets which we stored in the BDFFP office, and the PI or
508 Project Manager entered the data into plot-specific spreadsheets. These spreadsheets
509 had been prepared in advance; they were based on the previous census and the printed
510 versions were used to record data in the field. This helped reduce data entry errors by
511 allowing the person entering the data to verify a plant's tag number on both the paper
512 data sheet and the electronic spreadsheet prior. Questions regarding the interpretation
513 of field observations or values recorded were clarified immediately with the person
514 recording the data. We identified potential errors or outliers with histograms of plant
515 height and shoot number as well plots of individual plant height vs. shoot number.

516 **B. Quality assurance/quality control procedures:** Once the data for a plot had been
517 entered and verified, they were added to previous years' surveys by using tag ID and subplot
518 as the join keys. The measurements of plant height and stem number were then compared
519 with those from the previous year to identify potential errors in either plant measurement or
520 entry (e.g., a plant with 1 shoot in year t and 11 shoots in year t+1 is likely an error in data
521 entry). Discrepancies were investigated by referring to the original data sheets and, on
522 occasion, returning to the field to remeasure plants.

523 **C. Related materials:** Each demographic plots' location, orientation, and subdivision into
524 subplots can be seen in Fig. 1. Links to photographs, the output of data validation
525 algorithms, summaries of other data sets collected by HDP researchers, and other related
526 materials can be found on the `README.md` file of the HDP Github Repository.

527 **D. Computer programs and data-processing algorithms:** The version of the R code
528 used to prepare this data archive can be found at Zenodo [*url to be added*]. Post-publication
529 updates to the code and data will be available on the `NEWS.md` file of the HDP Repository
530 (<https://github.com/BrunaLab/HeliconiaSurveys>) until updated version of the archives are
531 uploaded to Zenodo and Dryad.

532 **F. Publications using the demographic data set:** The following publications include
533 analyses of the demographic data set. Links to an updated publication list and downloadable
534 *BibTeX* file can be found on the `README.md` file of the HDP Github repository
535 (<https://github.com/BrunaLab/HeliconiaSurveys>)

- 536 1. Bruna, E. M. and W. J. Kress. 2002. Habitat fragmentation and the
537 demographic structure of an Amazonian understory herb (*Heliconia*
538 *acuminata*). *Conservation Biology* 16(5): 1256-1266.
- 539 2. Bruna, E. M., O. Nardy, S. Y. Strauss, and S. P. Harrison. 2002.
540 Experimental assessment of *Heliconia acuminata* growth in a fragmented
541 Amazonian landscape. *Journal of Ecology* 90(4): 639-649.
- 542 3. Bruna, E. M. 2002. Effects of forest fragmentation on *Heliconia acuminata*
543 seedling recruitment in the central Amazon. *Oecologia* 132:235-243.
- 544 4. Bruna, E. M. 2003. Are plant populations in fragmented habitats
545 recruitment limited? Tests with an Amazonian herb. *Ecology* 84(4):
546 932-947.

- 547 5. Bruna, E. M. 2004. Biological impacts of deforestation and fragmentation.
548 Pages 85-90 in *The Encyclopaedia of Forest Sciences*. J. Burley, J Evans,
549 and J Youngquist, (eds.). Elsevier Press, London.
- 550 6. Bruna, E. M., and M. K. Oli. 2005. Demographic effects of habitat
551 fragmentation on a tropical herb: Life-table response experiments. *Ecology*
552 86: 1816-1824.
- 553 7. Bruna E. M. & W. J. Kress. 2005. Forest fragments and plant reproduction
554 in Amazonian Brazil. pp. 141-146 in G. A. Krupnick & W. J. Kress (eds).
555 *Plant conservation: a natural history approach*. University of Chicago Press,
556 Chicago.
- 557 8. Morris, W. F., C. A. Pfister, S. Tuljapurkar, C. V. Haridas, C. Boggs, M. S.
558 Boyce, E. M. Bruna, D. R. Church, T. Coulson, D. F. Doak,, S. Forsyth,
559 J-M. Gaillard, C. C. Horvitz, S. Kalisz, B. E. Kendall, T. M. Knight, C. T.
560 Lee, and E. S. Menges. 2008. Longevity can buffer plant and animal
561 populations against changing climatic variability. *Ecology* 89(1): 19-25.
- 562 9. Fiske, I., E. M. Bruna, and B. M. Bolker. 2008. Effect of sample size on
563 estimates of population growth rates calculated with matrix models. *PLoS
564 ONE* 3(8): e3080.
- 565 10. Fiske, I. and E. M. Bruna. 2010. Alternative spatial sampling in studies of
566 plant demography: consequences for estimates of population growth rate.
567 *Plant Ecology* 207(2): 213-225.
- 568 11. Uriarte, M., E. M. Bruna, P. Rubim, M. Anciaes, and I. Jonckeeere. 2010.
569 Effects of forest fragmentation on seedling recruitment of an understory
570 herb: assessing seed vs. safe-site limitation. *Ecology* 91(5): 1317-1328.

- 571 12. Gagnon, P. R., E. M. Bruna, P. Rubim, M. R. Darrigo, R. C. Littlel, M.
572 Uriarte, and W. J. Kress. 2011. The growth of an understory herb is
573 chronically reduced in Amazonian forest fragments. *Biological Conservation*
574 144: 830-835.
- 575 13. Uriarte, M. Anciães, M. T.B. da Silva, P. Rubim, E. Johnson, and E. M.
576 Bruna. 2011. Disentangling the drivers of reduced long-distance seed
577 dispersal by birds in an experimentally fragmented landscape. *Ecology*
578 92(4): 924-93.
- 579 14. Côrtes, M., M. Uriarte, M. Lemes, R. Gribel, W. J. Kress, P. Smouse, E. M.
580 Bruna. 2013. Low plant density enhances gene flow in the Amazonian
581 understory herb *Heliconia acuminata*. *Molecular Ecology* 22: 5716-5729.
- 582 15. Brooks, M. E., K. Kristensen, M. R. Darrigo, P. Rubim, M. Uriarte, E. M.
583 Bruna, and B. M Bolker. 2019. Statistical modeling of patterns in annual
584 reproductive rates. *Ecology* 100(7): e02706.
- 585 16. Leite, MCA, R. Sauchuk, F.B. Agusto, O. G. Gaoue, B. Chen-Charpentier.
586 2021. Modeling the persistence of plant populations in fragmented
587 ecosystems. *Ecological Modelling* 457:109681.
- 588 17. Scott, E. R., M. Uriarte, and E. M. Bruna. 2022. Delayed effects of climate
589 on vital rates lead to demographic divergence in Amazonian forest
590 fragments. *Global Change Biology* 28(2): 463-479.
- 591 18. Leite, M. C. A., R. Sauchuk, F. B. Agusto, O. G. Gaoue, and B.
592 Chen-Charpentier. 2021. Modeling the persistence of plant populations in
593 fragmented ecosystems. *Ecological Modelling* 457: 109681.

594 **G. Related publications and data sets:** The information in the following data archives

595 and publications can be used in concert with the census data to conduct demographic
596 modeling or carry out other analyses. The HDP Github Repository's `README.md` file
597 provides links to an updated list and downloadable BibTeX file of these publications and
598 data sets (<https://github.com/BrunaLab/HeliconiaSurveys>).

- 599 1. Bruna, E. M. 1999. Seed germination in rainforest fragments. *Nature*
600 402(6758):139–139.
- 601 2. Bruna, E. M., W. John Kress, Francisco Marques, and Osmaildo Ferreira da
602 Silva. 2004. *Heliconia acuminata* reproductive success is independent of
603 local floral density. *Acta Amazonica* 34(3):467–471.
- 604 3. E. M. Bruna and M. B. N. Ribeiro. 2005. The compensatory responses of
605 an understory herb to experimental damage are habitat-dependent.
606 *American Journal of Botany* 92(12):2101–2106.
- 607 4. E. M. Bruna and M. B. N. Ribeiro. 2005. Regeneration and population
608 structure of *Heliconia acuminata* in Amazonian secondary forests with
609 contrasting land-use histories. *Journal of Tropical Ecology* 21(1):127–131.
- 610 5. Côrtes, M. C., V. Gowda, W. J. Kress, E. M. Bruna, and M. Uriarte. 2009.
611 Characterization of 10 microsatellite markers for the understorey Amazonian
612 herb *Heliconia acuminata*. *Molecular Ecology Resources* 9(4):1261–1264.
- 613 6. Ribeiro, M. B. N., E. M. Bruna, and Waldir Mantovani. 2010. Influence of
614 post-clearing treatment on the recovery of herbaceous plant communities in
615 Amazonian secondary forests. *Restoration Ecology* 18: 50–58.
- 616 7. Bruna, E. M. and A. S. Andrade. 2011. Edge effects on growth and biomass
617 partitioning of an Amazonian understory herb (*Heliconia acuminata*;
618 Heliconiaceae). *American Journal of Botany*. 98(10):1727–1734.

- 619 8. Bruna, E. M. 2014. *Heliconia acuminata* seed set (seeds per fruit). 2008.
620 https://figshare.com/articles/dataset/Heliconia_acuminata_seedset_2008/1273926
- 622 9. Bruna, E. M. 2021. Leaf number, leaf area, shoot number, and height of
623 reproductive *Heliconia acuminata*. <https://doi.org/10.5281/zenodo.5041931>

624 **ACKNOWLEDGMENTS**

625 This work would not have been possible without the technicians and field assistants
626 participating in the annual surveys, the logistical support provided by the BDFFP staff, and
627 the organizations providing financial support. Sharon Y. Strauss, Susan P. Harrison, and
628 Patricia Delamônica Sampaio provided invaluable feedback during the project's
629 establishment. Charles L. Staines identified the Hispine beetle. Financial support for the
630 preparation of this manuscript was provided by the U.S. National Science Foundation (award
631 DEB-1948607). This paper is publication number -- in the BDFFP Technical Series.

References

- 632 Athayde, S. F. de, G. M. Da Silva, J. Kaiabi, M. Kaiabi, H. R. De Souza, K. Ono, and E. M.
633 Bruna. 2006. Participatory research and management of arumã (*Ischnosiphon gracilis*
634 [Rudge] Köern., Marantaceae) by the Kaiabi people in the Brazilian Amazon. *Journal of*
635 *Ethnobiology* 26: 36–59.
- 636 Berry, F., and W. J. Kress. 1991. *Heliconia*: An identification guide. Smithsonian
637 Institution Press, Washington D.C., USA.
- 638 Bierregaard, R. O., C. Gascon, T. E. Lovejoy, and R. Mesquita, editors. 2002. Lessons from
639 Amazonia: The ecology and conservation of a fragmented forest. Yale University Press,
640 New Haven, Connecticut, USA.
- 641 Broadbent, E. N., G. P. Asner, M. Keller, D. E. Knapp, P. J. C. Oliveira, and J. N. Silva.
642 2008. Forest fragmentation and edge effects from deforestation and selective logging in
643 the Brazilian Amazon. *Biological Conservation* 141: 1745–1757.
- 644 Brooks, M. E., K. Kristensen, M. R. Darrigo, P. Rubim, M. Uriarte, E. Bruna, and B. M.
645 Bolker. 2019. Statistical modeling of patterns in annual reproductive rates. *Ecology* 100:
646 e02706.
- 647 Brudvig, L. A., S. J. Leroux, C. H. Albert, E. M. Bruna, K. F. Davies, R. M. Ewers, D. J.
648 Levey, R. Pardini, and J. Resasco. 2017. Evaluating conceptual models of landscape
649 change. *Ecography* 40: 74–84.
- 650 Bruna, E. M. 1999. Seed germination in rainforest fragments. *Nature* 402: 139–139.
- 651 Bruna, E. M. 2002. Effects of forest fragmentation on *Heliconia acuminata* seedling
652 recruitment in central Amazonia. *Oecologia* 132: 235–243.
- 653 Bruna, E. M. 2003. Are plant populations in fragmented habitats recruitment limited? Tests
654 with an Amazonian herb. *Ecology* 84: 932–947.
- 655 Bruna, E. M. 2014. *Heliconia acuminata* seed set (seeds per fruit) 2008.
656 https://figshare.com/articles/dataset/Heliconia_acuminata_seedset_2008/1273926.
- 657 Bruna, E. M. 2021. Leaf number, leaf area, shoot number, and height of reproductive

- 659 *Heliconia acuminata*. <https://doi.org/10.5281/zenodo.5041931>.
- 660 Bruna, E. M., and A. S. de Andrade. 2011. Edge effects on growth and biomass partitioning
661 of an Amazonian understory herb (*Heliconia acuminata*; Heliconiaceae). *American
662 Journal of Botany* 98: 1727–1734.
- 663 Bruna, E. M., I. J. Fiske, and M. D. Trager. 2009. Habitat fragmentation and plant
664 populations: Is what we know demographically irrelevant? *Journal of Vegetation Science*
665 20: 569–576.
- 666 Bruna, E. M., and W. J. Kress. 2002. Habitat fragmentation and the demographic structure
667 of an Amazonian understory herb (*Heliconia acuminata*). *Conservation Biology* 16:
668 1256–1266.
- 669 Bruna, E. M., W. J. Kress, O. F. da Silva, and F. Marques. 2004. *Heliconia acuminata*
670 reproductive success is independent of local flowering plant density. *Acta Amazonica* 34:
671 467–471.
- 672 Bruna, E. M., O. Nardy, S. Y. Strauss, and S. P. Harrison. 2002. Experimental assessment
673 of *Heliconia acuminata* growth in a fragmented Amazonian landscape. *Journal of
674 Ecology* 90: 639–649.
- 675 Bruna, E. M., and M. B. N. Ribeiro. 2005a. The compensatory responses of an understory
676 herb to experimental damage are habitat-dependent. *American Journal of Botany* 92:
677 2101–210.
- 678 Bruna, E. M., and M. B. N. Ribeiro. 2005b. Regeneration and population structure of
679 *Heliconia acuminata* in Amazonian secondary forests with contrasting land-use histories.
680 *Journal of Tropical Ecology* 21: 127–131.
- 681 Bruna, E., and M. Oli. 2005. Demographic effects of habitat fragmentation on a tropical
682 herb: Life-table response experiments. *Ecology* 86: 1816–1824.
- 683 Caswell, H. 2000. Matrix population models. Sinauer Associates, Sunderland, Massachusetts,
684 USA.
- 685 Cordeiro, N. J., H. J. Ndangalasi, J. P. McEntee, and H. F. Howe. 2009. Disperser limitation

- 686 and recruitment of an endemic African tree in a fragmented landscape. *Ecology* 90:
687 1030–1041.
- 688 Côrtes, M. C., V. Gowda, W. J. Kress, E. M. Bruna, and M. Uriarte. 2009. Characterization
689 of 10 microsatellite markers for the understorey Amazonian herb *Heliconia acuminata*.
690 *Molecular Ecology Resources* 9: 1261–1264.
- 691 Côrtes, M. C., M. Uriarte, M. R. Lemes, R. Gribel, W. John Kress, P. E. Smouse, and E. M.
692 Bruna. 2013. Low plant density enhances gene dispersal in the Amazonian understory
693 herb *Heliconia acuminata*. *Molecular Ecology* 22: 5716–5729.
- 694 Didham, R. K., V. Kapos, and R. M. Ewers. 2012. Rethinking the conceptual foundations of
695 habitat fragmentation research. *Oikos* 121: 161–170.
- 696 Ellner, S. P., and M. Rees. 2006. Integral projection models for species with complex
697 demography. *The American Naturalist* 167: 410–428.
- 698 Fearnside, P. M., and N. Leal-Filho. 2001. Soil and development in Amazonia: Lessons from
699 the Biological Dynamics of Forest Fragments Project. Yale University Press, New Haven,
700 Connecticut, USA.
- 701 Fletcher, R. J., R. K. Didham, C. Banks-Leite, J. Barlow, R. M. Ewers, J. Rosindell, R. D.
702 Holt, A. Gonzalez, R. Pardini, E. I. Damschen, F. P. L. Melo, L. Ries, J. A. Prevedello,
703 T. Tscharntke, W. F. Laurance, T. Lovejoy, and N. M. Haddad. 2018. Is habitat
704 fragmentation good for biodiversity? *Biological Conservation* 226: 9–15.
- 705 Gagnon, P. R., E. M. Bruna, P. Rubim, M. R. Darrigo, R. C. Littell, M. Uriarte, and W. J.
706 Kress. 2011. Growth of an understory herb is chronically reduced in Amazonian forest
707 fragments. *Biological Conservation* 144: 830–835.
- 708 Gascon, C., and Jr. Bierregaard Richard O. 2001. The Biological Dynamics of Forest
709 Fragments Project: The study site, experimental design, and research activity. Pages
710 31–42 *Lessons from Amazonia: The ecology and conservation of a fragmented forest*.
711 Yale University Press, New Haven, Connecticut, USA.
- 712 Gentry, A. H., and L. H. Emmons. 1987. Geographical variation in fertility, phenology and

- 713 composition of the understory of neotropical forests. *Biotropica* 19: 216–217.
- 714 Haddad, N. M., L. A. Brudvig, J. Clobert, K. F. Davies, A. Gonzalez, R. D. Holt, T. E.
715 Lovejoy, J. O. Sexton, M. P. Austin, C. D. Collins, W. M. Cook, E. I. Damschen, R. M.
716 Ewers, B. L. Foster, C. N. Jenkins, A. J. King, W. F. Laurance, D. J. Levey, C. R.
717 Margules, B. A. Melbourne, A. O. Nicholls, J. L. Orrock, D. X. Song, and J. R.
718 Townshend. 2015. Habitat fragmentation and its lasting impact on Earth's ecosystems.
719 *Science Advances* 1: e1500052.
- 720 Harris, L. D. 1984. The fragmented forest: Island Biogeography Theory and the preservation
721 of biotic diversity. University of Chicago Press, Chicago, Illinois, USA.
- 722 Harrison, S., and E. Bruna. 1999. Habitat fragmentation and large-scale conservation: What
723 do we know for sure? *Ecography* 22: 225–232.
- 724 Holdridge, L. R. 1967. Life zone ecology. Tropical Science Center, San Jose, Costa Rica.
- 725 Iannone, R., and M. Vargas. 2022. Pointblank: Data validation and organization of
726 metadata for local and remote tables. R package version 0.10.0.
727 [Https://CRAN.R-project.org/package=pointblank](https://CRAN.R-project.org/package=pointblank).
- 728 Jurinitz, C. F., A. A. de Oliveira, and E. M. Bruna. 2013. Abiotic and biotic influences on
729 early-stage survival in two shade-tolerant tree species in Brazil's Atlantic Forest.
730 *Biotropica* 45: 728–736.
- 731 Kress, J. 1990. The diversity and distribution of *Heliconia* (Heliconiaceae) in Brazil. *Acta
732 Botanica Brasileira* 4: 159–167.
- 733 Kress, W. J. 1983. Self-incompatibility in central American *Heliconia*. *Evolution* 37:
734 735–744.
- 735 Lasky, J. R., M. Uriarte, V. K. Boukili, D. L. Erickson, W. John Kress, and R. L. Chazdon.
736 2014. The relationship between tree biodiversity and biomass dynamics changes with
737 tropical forest succession. *Ecology Letters* 17: 1158–1167.
- 738 Laurance, W. F. 2001. The hyper-diverse flora of the central Amazon. Pages 47–53 in R. O.
739 Bierregaard Jr., C. Gascon, T. E. Lovejoy, and R. Mesquita, editors. Lessons from

- 740 Amazonia: The ecology conservation of a fragmented forest. Yale University Press, New
741 Haven, Connecticut, USA.
- 742 Laurance, W. F., J. L. C. Camargo, R. C. C. Luizão, S. G. Laurance, S. L. Pimm, E. M.
743 Bruna, P. C. Stouffer, G. Bruce Williamson, J. Benítez-Malvido, H. L. Vasconcelos, K. S.
744 Van Houtan, C. E. Zartman, S. A. Boyle, R. K. Didham, A. Andrade, and T. E. Lovejoy.
745 2011. The fate of Amazonian forest fragments: A 32-year investigation. *Biological
746 Conservation* 144: 56–67.
- 747 Laurance, W. F., L. V. Ferreira, J. M. Rankin de Merona, and S. G. Laurance. 1998. Rain
748 forest fragmentation and the dynamics of Amazonian tree communities. *Ecology* 79:
749 2032–2040.
- 750 Laurance, W. F., T. E. Lovejoy, H. L. Vasconcelos, E. M. Bruna, R. K. Didham, P. C.
751 Stouffer, C. Gascon, R. O. Bierregaard, S. G. Laurance, and E. Sampaio. 2002.
752 Ecosystem Decay of Amazonian Forest Fragments: A 22-Year Investigation.
753 *Conservation Biology* 16: 605–618.
- 754 Lovejoy, T. E., R. O. Bierregaard, A. B. Rylands, J. R. Malcolm, C. E. Quintela, L. H.
755 Harper, K. S. Brown, A. H. Powell, C. V. N. Powell, H. O. R. Schubart, and M. B. Hays.
756 1986. Edge and other effects of isolation on Amazon forest fragments. Pages 257–285 *in*
757 M. Soulé, editor. *Conservation Biology: The science of scarcity and diversity*. Sinauer
758 Associates, Sunderland, Massachusetts, USA.
- 759 Mesquita, R. C. G., K. Ickes, G. Ganade, and G. B. Williamson. 2001. Alternative
760 successional pathways in the Amazon Basin. *Journal of Ecology* 89: 528–537.
- 761 Murcia, C. 1996. Forest fragmentation and the pollination of neotropical plants. Pages 19–36
762 *in* Schelhas, J. and Greenberg, R., editors. *Forest patches in tropical landscapes*. Island
763 Press, Washington D.C., USA.
- 764 Nakazono, E. M., E. M. Bruna, and R. C. G. Mesquita. 2004. Experimental harvesting of
765 the non-timber forest product *Ischnosiphon polypyllus* in central Amazonia. *Forest
766 Ecology and Management* 190: 219–225.

- 767 Nessimian, J. L., E. M. Venticinque, J. Zuanon, P. De Marco, M. Gordo, L. Fidelis, J. D.
768 Batista, and L. Juen. 2008. Land use, habitat integrity, and aquatic insect assemblages
769 in Central Amazonian streams. *Hydrobiologia* 614: 117–131.
- 770 Oliveira, A. A. de, and S. A. Mori. 1999. A central Amazonian terra firme forest. I. High
771 tree species richness on poor soils. *Biodiversity and Conservation* 8: 1219–1244.
- 772 R Core Development Team. 2014. R: A language and environment for statistical computing.
773 R Foundation for Statistical Computing, Vienna, Austria.
- 774 Rankin-de-Mérona, J. M., G. T. Prance, R. W. Hutchings, M. F. da Silva, W. A. Rodrigues,
775 and M. E. Uehling. 1992. Preliminary results of a large-scale tree inventory of upland
776 rain forest in the central Amazon. *Acta Amazonica* 22: 493–534.
- 777 Resasco, J., E. M. Bruna, N. M. Haddad, C. Banks-Leite, and C. R. Margules. 2017. The
778 contribution of theory and experiments to conservation in fragmented landscapes.
779 *Ecography* 40: 109–118.
- 780 Ribeiro, M. B. N., E. M. Bruna, and W. Mantovani. 2010. Influence of post-clearing
781 treatment on the recovery of herbaceous plant communities in Amazonian secondary
782 forests. *Restoration Ecology* 18: 50–58.
- 783 Scariot, A. 1999. Forest fragmentation effects on palm diversity in central Amazonia.
784 *Journal of Ecology* 87: 66–76.
- 785 Scott, E. R., M. Uriarte, and E. M. Bruna. 2022. Delayed effects of climate on vital rates
786 lead to demographic divergence in Amazonian forest fragments. *Global Change Biology*
787 28: 463–479.
- 788 Silva, J. M. C. da, and M. Tabarelli. 2000. Tree species impoverishment and the future flora
789 of the Atlantic forest of northeast Brazil. *Nature* 404: 72–74.
- 790 Slik, J. W. F., S.-I. Aiba, F. Q. Brarley, C. H. Cannon, O. Forshed, K. Kitayama, H.
791 Nagamasu, R. Nilus, J. Payne, G. Paoli, A. D. Poulsen, N. Raes, D. Sheil, K. Sidiyasa, E.
792 Suzuki, and J. L. C. H. van Valkenburg. 2010. Environmental correlates of tree biomass,
793 basal area, wood specific gravity and stem density gradients in Borneo's tropical forests.

- 794 *Global Ecology and Biogeography* 19: 50–60.
- 795 Spicer, M. E., H. V. N. Radhamoni, M. C. Duguid, S. A. Queenborough, and L. S. Comita.
796 2022. Herbaceous plant diversity in forest ecosystems: Patterns, mechanisms, and threats.
- 797 *Plant Ecology* 223: 117–129.
- 798 Staines, C. L., and C. Garcia-Robledo. 2014. The genus *Cephaloleia* Chevrolat, 1836
799 (Coleoptera, Chrysomelidae, Cassidinae). *ZooKeys* 436: 1–355.
- 800 Stiles, F. G. 1975. Ecology, flowering phenology, and hummingbird pollination of some Costa
801 Rican *heliconia* species. *Ecology* 56: 285–301.
- 802 Stouffer, P. C., and R. O. Bierregaard. 1995. Effects of forest fragmentation on understory
803 hummingbirds in Amazonian Brazil. *Conservation Biology* 9: 1085–1094.
- 804 Stouffer, P. C., and R. O. Bierregaard. 1996. Forest fragmentation and seasonal patterns of
805 hummingbird abundance in Amazonian Brazil. *Ararajuba* 4: 9–14.
- 806 Strong, D. R. 1977. Rolled-leaf Hispine beetles (Chrysomelidae) and their Zingiberales host
807 plants in Middle America. *Biotropica* 9: 156–169.
- 808 Uriarte, M., M. Anciães, M. T. B. da Silva, P. Rubim, E. Johnson, and E. M. Bruna. 2011.
809 Disentangling the drivers of reduced long-distance seed dispersal by birds in an
810 experimentally fragmented landscape. *Ecology* 92: 924–937.
- 811 Uriarte, M., E. M. Bruna, P. Rubim, M. Anciães, and I. Jonckheere. 2010. Effects of forest
812 fragmentation on the seedling recruitment of a tropical herb: Assessing seed vs. Safe-site
813 limitation. *Ecology* 91: 1317–1328.
- 814 Vázquez-Yanes, C., and A. Orozco-Segovia. 1993. Patterns of seed longevity and germination
815 in the tropical rainforest. *Annual Review of Ecology and Systematics* 24: 69–87.
- 816 Wilcove, D. S., C. H. McLellan, and A. P. Dobson. 1986. Habitat fragmentation in the
817 temperate zone. Pages 237–256. *in* M. E. Soulé, editor. *Conservation biology: The*
818 *science of scarcity and diversity*. Sinauer Associates, Sunderland, Massachusetts, USA.
- 819 Yenni, G. M., E. M. Christensen, E. K. Bledsoe, S. R. Supp, R. M. Diaz, E. P. White, and S.
820 K. M. Ernest. 2019. Developing a modern data workflow for regularly updated data.

821 *PLOS Biology* 17: 1–12.

822 Zambrano, J., and R. Salguero-Gómez. 2014. Forest fragmentation alters the population
823 dynamics of a late-successional tropical tree. *Biotropica* 46: 556–564.

824 Zartman, C. E., J. A. Amaral, J. N. Figueiredo, and C. D. Dambros. 2015. Drought impacts
825 survivorship and reproductive strategies of an epiphyllous leafy liverwort in central
826 Amazonia. *Biotropica* 47: 172–178.




Figure 1. Satellite image of the Biological Dynamics of Forest Fragments Project (ca. 1995) showing the location of the *Heliconia* Demographic Plots. Plots are located in Continuous Forest (CF1-CF6) or Forest Fragments (FF1-FF7), both of which are dark green. Light green areas are regenerating forest, while red indicates pasture. The BDFFP is located 70 km north of Manaus, Brazil (inset map). For additional details on each plot see Table 1.




Figure 2. The abundance and reproduction of *Heliconia acuminata* in Continuous Forest.

In each of survey year (gray lines) the data from the six demographic plots in Continuous Forest were pooled to give (A) the total number of post-seedling *H. acuminata* in size classes based on shoot number, and (B) the percentage of plants in each size class that produced at least 1 inflorescence. The green line is the average of the annual values for each size category.




Figure 3. Schematic of the *Heliconia* Demographic Plots in the BDFFP 1-hectare forest fragment reserves (note: not to scale). The rectangles filled with a letter-number combination (i.e., F-3, A-2, E-2) are coordinate stakes marking the permanent plots of the BDFFP Phytodemographic Project (Rankin-de-Mérona et al. 1992).




Figure 4. Schematic of the *Heliconia* Demographic Plots in the BDFFP 10-ha forest fragment reserves (note: not to scale). The rectangles filled with a letter-number combination (i.e., I-8, J-8) are coordinate stakes marking the permanent plots of the BDFFP Phytodemographic Project (Rankin-de-Mérona et al. 1992).




Figure 5. Heliconia Demographic Plots in Continuous Forest (note: not to scale). Rectangles with a letter-number combination (i.e., L-40, J-40, C-1, C-2) are coordinate stakes marking permanent plots of the BDFFP Phytodemographic Project (Rankin-de-Mérona et al. 1992).

Table 1. Variable Information for “Data set File 1: *Descriptors of demographic plots*”.

Variable	Definition	Codes	Storage
plot_id	Code used to identify a plot	FF1-FF7: plots in fragments CF1-CF6: plots in continuous forest	string
habitat	Habitat in which a plot is located	one: 1-ha fragment ten: 10-ha fragment forest: continuous forest	string
ranch	Ranch in which a plot is located	porto alegre, esteio, dimona	string
bdffp_no	BDFFPs Reserve ID Number ¹	1104, 1202, 1301, 1501, 2107, 2108, 2206, 3209, 3402, NA	string
yr_isolated	For fragments, the year initially isolated	1980, 1983, 1984	integer

¹ See Gascon and Bierregaard (2001) for details of the reserve numbering scheme. ‘NA’ indicates the plot is not inside a formally demarcated BDFFP reserve.

Table 2. Variable Information for “Data set File 2: *Heliconia Demographic Data*”.

Variable	Definition	Codes or Range of Values	Storage
plot_id	Plot in which plant is located	FF1-FF7, CF1-CF6	string
subplot	Subplot in which plant is located	A1-E10 except in CF3, where F6-J10 ¹	string
plant_id	Unique ID no. assigned to plant	range = 1-8660 (units: number, precision: 1)	integer
tag_number	Number on tag attached to plant	range = 1-3751 (units: number, precision: 1)	integer
year	Calendar year of survey	range = 1998-2009 (units: year, precision: 1)	integer
shts	No. of shoots when surveyed	range = 0-24 (units: shoots, precision: 1) NA: data missing	integer
ht	Plant height when surveyed	range = 0-226 (units: cm, precision: 1) NA: data missing	integer
infl	No. of inflorescences (if flowering)	range = 1-7 (units: shoots, precision: 1) NA: data missing	integer
recorded_sdlg	New seedling	TRUE, FALSE	logical
adult_no_tag	Established (i.e., post-seedling) individual without tag	TRUE, FALSE	logical
treefall_status	Plant found under fallen tree crown, branches, or leaf litter at time of survey	branch: under fallen tree limbs tree: under tree crown or fallen trees litter: under accumulated leaf-litter NA: not relevant or no observation recorded	string

census_status	Plant status in a census	measured: alive, measured dead: died prior to census missing: not found during census	string
---------------	--------------------------	---	--------

¹ For the arrangement of the subplots see Figures 3-5