

The effects of climate change and land-use change on demographic rates and population viability

Katherine E. Selwood^{1,*}, Melodie A. McGeoch¹ and Ralph Mac Nally²

¹*School of Biological Sciences, Monash University, Clayton, Victoria 3800, Australia*

²*Institute for Applied Ecology, The University of Canberra, Bruce, Australian Capital Territory 2617, Australia*

ABSTRACT

Understanding the processes that lead to species extinctions is vital for lessening pressures on biodiversity. While species diversity, presence and abundance are most commonly used to measure the effects of human pressures, demographic responses give a more proximal indication of how pressures affect population viability and contribute to extinction risk. We reviewed how demographic rates are affected by the major anthropogenic pressures, changed landscape condition caused by human land use, and climate change. We synthesized the results of 147 empirical studies to compare the relative effect size of climate and landscape condition on birth, death, immigration and emigration rates in plant and animal populations. While changed landscape condition is recognized as the major driver of species declines and losses worldwide, we found that, on average, climate variables had equally strong effects on demographic rates in plant and animal populations. This is significant given that the pressures of climate change will continue to intensify in coming decades. The effects of climate change on some populations may be underestimated because changes in climate conditions during critical windows of species life cycles may have disproportionate effects on demographic rates. The combined pressures of land-use change and climate change may result in species declines and extinctions occurring faster than otherwise predicted, particularly if their effects are multiplicative.

Key words: climate variation, extinction risk, extirpation, emigration, immigration, land-use intensification, landscape condition, mortality, natality.

CONTENTS

I. Introduction	838
(1) Factors affecting population viability	839
II. Conceptual model	839
(1) Overview of land-use change and climate change	839
(2) Model description	840
III. Quantifying the effects of human pressures on demographic rates	840
(1) Literature search	841
(2) Quantification of effect sizes	841
(3) Results	842
IV. Mechanisms affecting demographic rates	842
(1) Birth rates	843
(2) Death rates	844
(3) Emigration and immigration	845
V. Synthesis and future work	846
VI. Conclusions	847
VII. Acknowledgements	847
VIII. References	847
IX. Supporting Information	853

* Address for correspondence (Tel: +61 399020860; E-mail: katherine.selwood@monash.edu).

I. INTRODUCTION

Biodiversity continues to decline; of species that have been assessed for extinction risk around the world, 38% are considered to be under threat (Vié, Hilton-Taylor & Stuart, 2009). The abundances of vertebrate populations fell by one-third between 1970 and 2006, and continue to decline; 70% of assessed plant species have been classified as threatened by the IUCN (Vié *et al.*, 2009). The principal pressures causing biodiversity loss are unabated, and, in most cases, are increasing (Butchart *et al.*, 2010). Human land-use change, leading to the loss, fragmentation and degradation of native vegetation, is the predominant driver of terrestrial species decline (Sala *et al.*, 2000). Climate change has been recognized comparatively recently as a major driver, and its effect on plant and animal populations is increasing (Bellard *et al.*, 2012; Foden *et al.*, 2013).

The most widely used measure of biodiversity is species richness, although subspecies, races and genotypes are important components. However, it is the extinction of individual species, especially iconic ones, that causes most consternation among practitioners and the public, so that it is important to understand the processes leading to species extinction. While there is a relatively good understanding of the identity of the pressures acting on species, the mechanisms by which these pressures operate and interact to affect the viability of species and populations is poorly understood (Akçakaya *et al.*, 2006). Understanding the processes that ultimately cause species extinctions is critical for deciding on the most appropriate actions for conservation management (Cushman, 2006).

The effects of land-use change have been a focus for conservation biology for several decades, particularly the effects of habitat fragmentation (Fischer & Lindenmayer, 2007). The most common measures for quantifying the effects on biota are species richness, species occurrence and the abundance patterns of individual species (Debinski & Holt, 2000). Few studies on fragmentation measure demographic responses, with most studies measuring presence/absence, diversity, or abundance (McGarigal & Cushman, 2002); these are 'static' rather than dynamic measures, and so generally do not provide much information on the trajectories of change. There has been much less focus on the demographic effects of land-use change on populations, which provide indications of trajectories of change (Lampila, Mönkkönen & Desrochers, 2005).

Climate change is expected to become an equally, or more important, driver of global biodiversity loss over the next century (Heller & Zavaleta, 2009). Climate change and climatic events (e.g. drought) have already caused range shifts (Chen *et al.*, 2011), severe and long-term population declines (Sanderson *et al.*, 2006; Newton, 2008b) and extinctions (Thomas, Franco & Hill, 2006). While interest in the effects of climate on biodiversity has escalated in recent decades, studies on the effects of climate have predominantly focused on observed and potential shifts in species ranges (Dawson *et al.*, 2011) and changes in species phenology (Parmesan,

2006; Chambers & Keatley, 2010) and physiology (Buckley, Nufio & Kingsolver, 2013). These factors may indicate or lead to a change in the likelihood of a species' persistence, but they do not directly reveal the changes in demographic rates that determine the chances that a population will persist. Changes in the phenology, such as timing of breeding, do not in themselves indicate a deleterious effect on population viability. The population is affected when these changes alter demographic rates.

Geographic distribution is the spatial expression of demographic rates, but change in distribution is one of the last signals to be detected as a species declines (Martinez-Meyer, 2012). Focusing on shifts in species ranges misses the population-level processes leading to these shifts, including local extinctions and recolonizations, and the changes in demographic rates that lead to these. While species-distribution models may predict range expansions with climate change, demographic studies may indicate the opposite effect (Campbell *et al.*, 2012). Organisms may colonize or remain in poor-quality habitat if there is asynchrony between the cues used for habitat selection and declines in habitat suitability caused by climate change (van de Pol *et al.*, 2010), so that distributions do not necessarily inform population viability.

We refer to 'pressure' as a human-induced perturbation that negatively affects a population and that may be transient (pulse), persistent (press), or monotonically changing in magnitude (ramping) over time. We synonymize pressure with 'stressor' and 'threat'. Pressures have causative effects on demographic rates (e.g. decreased seed germination, increased nest predation), while associations between pressures and changes in species richness, species occurrence and abundance are correlative. The close connection between a pressure and a demographic-rate response means that measuring the changes in demographic rates should offer a more accurate indication of the mechanisms through which anthropogenic pressures affect population viability (Fig. 1).

Here, we review the effects of some of the major anthropogenic pressures on population viability, and we present a conceptual model to describe these relationships. We focus on the processes through which climate change and changed landscape condition induced by human land use affect population viability in terrestrial plant and animal populations. Last, we quantify these relationships by synthesizing the results of empirical studies to provide a comparison of the effects of these major pressures on population viability. For tractability, this review concentrates on terrestrial systems; different sets of pressures may predominate for freshwater (Ficke, Myrick & Hansen, 2007) and marine (Halpern *et al.*, 2008) systems. There are other pressures on biodiversity such as direct harvest (including fisheries), pollution, invasive species and disease (Mace, Masundire & Baillie, 2005). These are vast topics, so we do not consider them further; instead we focus on the influence of landscape condition and climate change as the main pressures of interest, given their pervasive influence.




Fig. 1. A general representation of the linkages between the effects of a pressure, such as vegetation loss, and commonly used measures of populations.

(1) Factors affecting population viability

Population viability is a quantitative measure of the capacity of a population to persist, typically the probability of persistence for 100 years, which indicates the risk of extinction (Boyce, 1992). Population viability analyses often are used to quantify extinction risks for individual populations, which can include the identification of minimum viable population size (Reed *et al.*, 2003). The processes that lead to the extinction of a population arise from deleterious changes to demographic rates, which occur through changes in reproductive output, survival and dispersal of individuals in response to a pressure (Fig. 1). Population viability is based on likely changes in population size over time, with the component demographic rates contributing to changes in population size. Birth, death, immigration and emigration are the four fundamental demographic parameters that determine changes in population size (Begon, Mortimer & Thompson, 1996). The dynamics of a population can be represented by (Cohen, 1969):

$$N_{t+1} = N_t (1 + b + i - d - e),$$

where: N_t is the abundance of a population at time t , b and d are the *per capita* birth and death rates, and i and e are *per capita* immigration and emigration rates during time interval $(t + 1) - t$. The effective population size will be affected by the sex ratios of individuals contributing to these demographic rates (Frankham, 1995). If one or more of these demographic rates is affected by a proximal pressure, arising from a distal driver, then this will affect the size of the population, and may decrease its viability, unless offset by changes to another demographic rate (i.e. consistently have $N_{t+1} < N_t$, Fig. 2). Once populations become small, stochastic events, inbreeding depression and genetic erosion further affect demographic rates and steepen the rate of decline in population viability (Young *et al.*, 2000; Keller & Waller, 2002). Given the direct effects on population dynamics, measuring changes in demographic rates allows us to infer likely changes to a population's viability in response to human pressures.

II. CONCEPTUAL MODEL

(1) Overview of land-use change and climate change

Changes in human land use for food and resource production and urbanization affect landscape condition through the loss and fragmentation of native vegetation (Fahrig, 1997) and the degradation of remnant vegetation (Fischer & Lindenmayer, 2007). Climate change can further degrade vegetation condition through changes to the frequency and intensity of disturbances that can affect vegetation composition, structure and function (Cunningham *et al.*, 2009; Bennett *et al.*, 2013), decrease plant growth and cause disruptions to plant–pollinator interactions (Memmott *et al.*, 2007). In some locations, increased temperature or carbon dioxide levels may enhance plant growth (Reich & Oleksyn, 2008; Wigley, Bond & Hoffman, 2010).

Barriers to movement caused by vegetation loss and fragmentation affect the movement of individuals and propagules (Cunningham, 2000a; Schtickzelle & Baguette, 2003). Vegetation loss and degradation alter microclimates, habitat quality and habitat structure, affecting conditions for survival and reproduction and modify species interactions (Mac Nally, Bennett & Horrocks, 2000). Resources for survival and reproduction are diminished in degraded and fragmented vegetation (Zanette, Doyle & Tremont, 2000).

Changes to the global climate include increased global temperature and sea levels, decreased extent of snow and ice (both sea and ice-caps) and increased prevalence and intensity of drought (IPCC, 2013). Changes to climate alter demographic rates because of the physiological responses of organisms to environmental variables such as temperature, which affect survival and reproduction (Chown *et al.*, 2010). Climate conditions affect dispersal behaviour (Altermatt, Pajunen & Ebert, 2008) and pathways (Kuparinen *et al.*, 2009). Climate-induced changes to phenology are well documented (Parmesan, 2006), and these affect demographic rates through their effects on reproduction and survival (Lehikoinen, Kilpi & Öst, 2006; Briscoe *et al.*, 2012), through mismatches in trophic relationships and species interactions (Durant *et al.*, 2007; Miller-Rushing *et al.*, 2010).




Fig. 2. A general model of the how anthropogenic pressures (P_1 – P_4) impinge on demographic rates (i , e , b , d) in complex networks of effects. The quantification of the strengths of the relationships is key to managing population viability. ‘Pressure’ refers to a human-induced perturbation that negatively affects a population and that may be transient (pulse), persistent (press), or monotonically changing in magnitude (ramping) over time (Mac Nally *et al.*, 2011). A population may be exposed to multiple pressures and the one pressure may affect multiple syntopic populations of different species. We synonymize pressure with ‘stressor’ and ‘threat’.

Demographic rates are controlled by resource availability (Skogland, 1985), such as food, which depends on climate (Previtali *et al.*, 2009; Tian *et al.*, 2010). Some populations may benefit from climate change, perhaps through an increase in survival or growth with warmer temperatures (Reich & Oleksyn, 2008). Climate-induced changes to species interactions may benefit some populations by competitor or predator release, while others may be adversely affected by, for example, weakened mutualistic relationships (Tylianakis *et al.*, 2008).

Despite the numerous mechanisms through which land-use change and climate change affect demographic rates, there has been little attention to the relationships between these pressures and demographic responses. Identifying and quantifying the pathways through which anthropogenic pressures affect population viability is important for framing management actions to contribute to population persistence.

(2) Model description

Multiple pressures need to be considered together because pressures rarely occur singly and interactions among pressures may be multiplicative rather than additive (Dawson *et al.*, 2011; Mantyka-Pringle, Martin & Rhodes, 2012). The relationships among pressures and demographic rates are shown in Fig. 2.

Depending on biological characteristics such as longevity, sexual maturity, and propensity to disperse, changes in one or more demographic rates may have a greater influence on population viability than a proportionally similar change in others (Harper, Rittenhouse & Semlitsch, 2008). For

example, long-lived species are most affected by changes in death rates because adult survivorship contributes most to population persistence (Li *et al.*, 2009).

By populating the general model of Fig. 2 with empirical information, we show how the principal human pressures (Mace *et al.*, 2005) impinge on demographic rates in plant and animal populations (Fig. 3). The model emphasizes the large roles that land-use change and climate change play in affecting population viability, which we quantify in Section III.

The loss, fragmentation and degradation of native vegetation are proximal ecological pressures stemming from land-use change, which affect demographic rates and population viability through their effects on landscape condition and resource availability. We refer to ‘landscape condition’ as the degree to which a landscape resembles its natural condition prior to substantial human disturbance or alteration, consisting of native vegetation cover, connectivity and quality. Climate change and changed landscape condition decrease resource availability, such as food, shelter, soil, nutrients, water and other resources necessary for population survival.

III. QUANTIFYING THE EFFECTS OF HUMAN PRESSURES ON DEMOGRAPHIC RATES

Here, we parameterized the strength of the linkages in the conceptual model (Fig. 3) using a representative set of literature estimates. We quantified the effects of changed landscape condition and climate variation on




Fig. 3. Empirical application of the general model of Fig. 2 to effects on population dynamics in terrestrial landscapes; the same conventions apply. Grey arrows represent relationships reviewed here. Arrow width represents the mean effect size of the relationship for reviewed studies: low = $0.2 < r < 0.4$; medium $0.4 < r < 0.6$; high = $0.6 < r < 0.8$. No mean effect sizes were < 0.2 (very low) or > 0.8 (very high).

demographic rates, which provides an assessment of the relative importance of changed landscape condition and climate change on population viability.

(1) Literature search

We searched for papers published between 1970 and 2012 using search terms consisting of descriptors for these pressures and demographic rates under TOPIC (i.e. title, abstract and key words) in Thomson–ISI *Web of Science* (Science Citation Index Expanded) (see online supporting information, Table S1), which returned 206 papers. We examined the titles and abstracts of the papers and retained those that provided quantitative relationships between pressures arising from climate or landscape condition and demographic variables, resulting in the retention of 24 papers. A second search, including broader terms for demography (see online Table S2) was conducted to find other studies that measured variables related to demographic rates. Searching for these terms within TOPIC returned >75000 papers; a random selection of 300 of these revealed no studies that provided quantitative information on the effects of a pressure on a demographic rate. The search was restricted to titles, returning 2324 papers, of which 209 were retained. Another 60 studies were found by using the reference lists of the 233 papers found during both searches.

We examined the results of the 294 studies to obtain statistics that were appropriate for calculating the r correlation coefficient (Rosenthal, 1994) for relationships between demographic variables and landscape condition or climate; this was possible for 147 studies. We used the r correlation coefficient because of its generality and simplicity

of interpretation and consistency of meaning (Rosenthal, 1994). While r is most appropriate for relationships between continuous variables, it can also be calculated from pairwise comparisons (Rosenthal, 1994). We included empirical, field-based or experimental studies that directly measured the effects of variables of climate and landscape condition on demographic variables in native plant and animal populations. Only 19 studies looked specifically at *per capita* demographic rates, so we included studies that measured variables that were related to these rates, such as clutch size, fruit production, juvenile survival, and genetic differentiation.

(2) Quantification of effect sizes

Values of the correlation coefficient r (including linear and rank correlations: Pearson's r , Spearman's r , Kendall's τ , point-biserial r , and phi) range between -1 and $+1$, and indicate the strength of the association between variables; the sign indicates the direction of the monotonic association (De Veaux, Velleman & Bock, 2008). Where no correlation coefficient was presented, we calculated r following standard methods (Rosenthal, 1994; Nakagawa & Cuthill, 2007) from reported test statistics (t -statistic, F -statistic, χ^2 , Z -score, coefficient of determination R^2 , Hedge's d). Where the P -value was the only statistic reported, we transformed these to Z -scores using a standard normal variate (De Veaux *et al.*, 2008).

The correlation coefficient r for each documented relationship between a climate variable (e.g. rainfall, temperature) or landscape-condition variable (e.g. vegetation cover, patch size) and the demographic response was

obtained from all species in each study. If >1 variable related to a particular demographic rate was measured (e.g. number of eggs and number of fledglings, or number of seeds and number of seedlings), we used the variable that would contribute most to the number of adult individuals in that population, usually the more advanced life stage (e.g. number of fledglings or number of seedlings). If >1 variable related to climate (e.g. rainfall and temperature) or to landscape condition (e.g. fragment size and isolation) was measured, we included the variable that had the largest effect size on the demographic response variable. Details of included studies and their effect sizes are in Table S3. Thirty-six studies measured >1 species, demographic rate and/or driver, and so, contributed >1 datum to the analysis.

For landscape condition, values of r ranged between -1 and $+1$, with positive values being associated with a positive effect of measures such as vegetation cover or contiguity on a demographic rate. For example, if fragmentation had a negative effect on a measure of birth rates in a study, the correlation coefficient for landscape condition on birth rates for that relationship would be positive.

We did not estimate the direction of relationships between climate variables (e.g. temperature, rainfall) and demographic rates because there is difficulty in generalizing the effects of climate variables on population viability given that directional climate deviations do not uniformly affect demographic rates (Glenn *et al.*, 2011). Changes in climate depend on region, so that generalizations are not appropriate. For example, there may be increases in precipitation in some regions and decreases in others, so that decreased rainfall cannot be considered to be a consistent climate-change effect (IPCC, 2013). The effects of climate variables on demographic rates may differ among seasons (Reed & Slade, 2009) and many studies measured within-year climate measures (e.g. winter rainfall) making it inappropriate to extrapolate to general trends given the scope of this review. We considered the correlation coefficient to be an absolute value for climate variables on demographic rates when calculating an average effect size, with r ranging from 0 to 1. This provides an indication of the size of the effect that climate may have on demographic rates rather than generalizing the effects of climate variables.

We converted all r values to ζ using Fisher's transformation, which transforms r to a near-normal distribution, because the distribution of r values becomes skewed as r becomes absolutely larger (Rosenthal, 1994). We calculated the mean effect size and standard error for the effect of landscape condition and climate on demographic rates using the ζ values to gauge the size of the effect that climate and landscape condition have on demographic processes and, in the case of pressures arising from landscape condition, the direction of this effect. Means were calculated for plants and animals separately. The means and upper and lower confidence interval values (95% confidence interval) were then back-transformed to r , so that the effect size could be between 0 and 1 for the effect of climate, and between -1 and $+1$ for the effect of landscape condition (Rosenthal, 1994).

(3) Results

Most studies on climate and landscape condition were from North America and Europe (see online Table S4). Birds were the most studied animals, followed by mammals, with other groups poorly represented (see online Table S3). There were few studies on the effects of climate on plant demographic rates (see online Table S3).

Landscape condition had a mean positive effect on birth rates in plant ($\bar{r}=0.3$) and animal populations ($\bar{r}=0.5$), a negative effect on death rates in animal populations ($\bar{r}=-0.6$), and a positive effect on plant dispersal and animal immigration ($\bar{r}=0.6$ for both). Landscape condition had a mean negative effect on death rates in plant populations ($\bar{r}=-0.6$) and emigration in animal populations ($\bar{r}=-0.2$), but studies were few ($N=2$ and 5) and confidence intervals overlapped zero, indicating that these effects were not significantly different from zero (Harrison, 2011) (Fig. 4A). The mean absolute effect sizes of climate on demographic rates were similar, for birth rates in plants ($\bar{r}=0.7$) and animals ($\bar{r}=0.6$), and plant ($\bar{r}=0.7$) and animal death rates ($\bar{r}=0.6$) (Fig. 4B).

There was a small mean effect size on animal emigration ($\bar{r}=0.2$), but there were only three studies, each of which reported increased measures of emigration with higher temperatures. There was just one study on animal immigration ($\bar{r}=0.6$) (Fig. 4B). There were no studies that provided statistics for calculating the effect size of climate on plant dispersal.

Studies that measured the effects of temperature and rainfall used a wide variety of temporal measures of climate (e.g. week, month, season, year, life-cycle stage), so we cannot extrapolate to responses to climate change (see online Table S5). For studies that reported an effect of rainfall, most were lower birth rates (11 of 13 studies) and increased death rates (five of eight studies) with decreasing rainfall (see online Table S5). For those assessing temperature effects, most showed a negative effect on birth rates (13 of 17 studies) and survival (five of five studies) with increasing temperatures (see online Table S5).

Landscape condition and climate appear to have substantial effects on demographic rates in plant and animal populations, with absolute effect sizes of 0.5–0.7 for all demographic rates except animal emigration (Fig. 4B). Given the large number of studies, there is good support for the positive effect of landscape condition on plant and animal birth rates and animal immigration (Fig. 4A). There were ≤ 5 studies on the effect of landscape condition on plant death rates, animal emigration and plant dispersal, but the directions of the relationships from these studies supported the conceptual model (Fig. 3).

IV. MECHANISMS AFFECTING DEMOGRAPHIC RATES

Here, we qualitatively review the mechanisms through which demographic rates in plant and animal populations are affected by changed landscape condition and climate change.




Fig. 4. (A) Mean effect sizes (\bar{r}) of landscape condition parameters and (B) mean absolute effect sizes (\bar{r} ; values between 0 and 1 only) of climate parameters on measures of birth rates, death rates, immigration, emigration and dispersal in plant and animal populations. Error bars are 95% confidence intervals. The number of data points (study \times species) is shown by N .

(1) Birth rates

Our quantitative review shows strong evidence for a negative effect of changed landscape condition on birth rates in plant and animal populations. The most proximate effect on plant reproduction in changed landscapes is usually pollination limitation (Aguilar *et al.*, 2006). Changed landscape condition results in declines in native pollinator populations and reduced pollinator visitation due to isolation, which reduces fruit production and seed set (Wilcock & Neiland, 2002; Gómez *et al.*, 2010). Allee effects, including inbreeding and genetic erosion, affect mate availability and seed set and interact with pollen limitation to reduce population viability (Wagenius, Lonsdorf & Neuhauser, 2007; Young, Broadhurst & Thrall, 2012). Wind-pollination may be disrupted by fragmentation, possibly causing inbreeding (Jump & Peñuelas, 2006). Reduced seed dispersal or increased seed predation occur in modified landscapes (Benitez-Malvido, 1998; Tallmon *et al.*, 2003). Loss and degradation of vegetation alters the conditions for germination and seedling establishment, including light environments (Uriarte *et al.*, 2010), microclimatic conditions (Jacquemyn *et al.*, 2003; Werner & Gradstein, 2008), and wind erosion (Li *et al.*, 2009). Grazing by domestic stock causes trampling and herbivory of seedlings (Jansen & Robertson, 2001). These declines in plant recruitment have large effects on population viability (Bruna & Oli, 2005).

Vegetation loss and fragmentation influence birth rates in animal populations by affecting access to food resources (Mbora, Wiczowski & Munene, 2009) and by reducing food and resource levels in vegetation remnants (Zanette *et al.*, 2000). Increased nest predation and parasitism are

common in much-modified landscapes, particularly near vegetation boundaries (Lampila *et al.*, 2005). Decreased vegetation connectivity reduces mate availability (Cooper & Walters, 2002), including through inbreeding avoidance (Boudjemadi, Lecomte & Clobert, 1999; Stow & Sunnucks, 2004). Some plant and animal populations experience higher birth rates in changed landscapes, especially those with a preference for open or edge habitat (Mac Nally *et al.*, 2012), or through decreased competition for resources such as light (Neal, Hardner & Gross, 2010).

Climate had a strong effect on birth rates, affecting rates in several ways. Most studies reported decreased birth rates with increased temperature and decreased precipitation. Global temperatures have risen, and the frequency of hot days and of heat waves is likely increasing (IPCC, 2013). Increased annual temperatures and short-term heat waves may reduce germination of plants (Chidumayo, 2008; Shevtsova *et al.*, 2009). In animals, heat stress of parents may induce declines in neonatal survival (Griffin *et al.*, 2011) and decreased fecundity (Neveu, 2009). Higher temperatures may cause heat stress in young animals, leading to lower survival rates of young (Steenhof, Kochert & McDonald, 1997). Warm and dry conditions, such as those associated with El Niño events, may harm eggs and hatchlings by altering microclimate conditions in nests (Tomillo *et al.*, 2012), although warmer temperatures may increase hatching success (Beissinger, Cook & Arendt, 2005). Warmer temperatures may enhance the breeding success and survival of young and seedlings by reducing energy needs (Nielsen & Møller, 2006; Milbau *et al.*, 2009) and reducing the occurrence of severe winters that limit reproductive success (McIntyre & Schmidt, 2012). Warming may lengthen periods suitable for breeding and result in

increased birth rates and additional generations within an annual cycle (Jönsson *et al.*, 2009; Clarke & Zani, 2012).

Lower rainfall and increased drought frequency may affect plant birth rates through decreased fruit set (Ågren, Ehrlén & Solbreck, 2008) and seedling survival (Hallett, Standish & Hobbs, 2011). Reduced food-plant productivity and food availability during periods of low rainfall, such as in El Niño events, may depress fecundity (Dunham, Erhart & Wright, 2010), prevent reproductive maturation (Lima *et al.*, 2001), and lessen offspring survival (Silleet, Holmes & Sherry, 2000). Limited water availability for lactating females may affect juvenile survival (Dunham *et al.*, 2010). Lower water levels at aquatic breeding sites may result in increased ultraviolet radiation and heat exposure, which can affect hatching success (Blaustein *et al.*, 2012), increase the vulnerability of embryos to pathogens (Kiesecker, Blaustein & Belden, 2001), and desiccate tadpoles (Pechmann *et al.*, 1991). Heavy rains or snowfalls, which are expected to increase in frequency even in areas with decreased annual precipitation (IPCC, 2013), stress gestating females (Dunham *et al.*, 2010), and increase juvenile and egg mortality (Skagen & Adams, 2012).

Phenological changes triggered by climate changes such as earlier warming, may increase self-fertilization in monoecious plants or cause mistiming in the flowering of dioecious plants (Miller-Rushing *et al.*, 2010). While earlier breeding may benefit birth rates of some species (Nielsen & Møller, 2006), advances in breeding and flowering expose flower buds (Inouye, 2000, 2008), eggs and young (Lehikoinen *et al.*, 2009) to poor or more variable weather conditions (e.g. frosts or heavy rain) if seasonal climate patterns do not advance in concert. Changed climate conditions may delay breeding so that the young may experience adverse conditions later in the season (Waite & Strickland, 2006; Senapathi *et al.*, 2011), and may inhibit breeding altogether (Pankhurst & Munday, 2011). Phenological changes in plants can cause asynchrony with pollinators, increase exposure to florivores and granivores, and increase synchrony of flowering among species competing for pollinators (Miller-Rushing *et al.*, 2010). Phenological changes to a population or its biotic resource may affect birth rates if the two do not change in synchrony. Asynchrony between food needs during breeding and food availability arises from earlier breeding (Moss, Oswald & Baines, 2001), advancement of peak prey availability (Sanz *et al.*, 2003), advanced phenology of food and larval host plants (Parmesan, 2005; Post & Forchhammer, 2008), or changed timing of food peaks (Wolf *et al.*, 2009). Climate-induced asynchronies in resource availability and resource needs during breeding have caused population extinctions (McLaughlin *et al.*, 2002). For some species, earlier warming may increase synchrony with food resources, which can increase birth rates (Vatka, Orell & Rytönen, 2011).

Spring snow cover is decreasing in the northern hemisphere (Werner, 2011). Reduction of snow cover may decrease seedling survival by permitting increased herbivory (Brodie *et al.*, 2012) and by increasing exposure to frost (Bannister *et al.*, 2005). Sea levels are rising (IPCC, 2013), and this can affect birth rates through more frequent

flooding of coastal nesting sites (van de Pol *et al.*, 2010). Physiological stress from severe weather limits reproductive success of many animals (Dunham *et al.*, 2010).

Within species, the extents to which birth rates are affected by climate changes differ depending on the elevational (Munier *et al.*, 2010; Hargrove & Rotenberry, 2011) or latitudinal (Ontiveros & Pleguezuelos, 2003; Sanz, 2003) location of populations, with some populations experiencing opposite effects of climate on birth rates in different locations (Gaston, Gilchrist & Hipfner, 2005). Climate effects on other demographic characteristics, such as death rates or sex ratios can dampen or counter positive effects (Zani, 2008; Schwanz *et al.*, 2010).

The effects of both landscape change and climate are diverse, and it is possible that there will be interactions or additive effects of these pressures on birth rates. However, while studies on variables related to birth rates were the most numerous of the demographic rates, this does not necessarily reflect the proportional importance of birth rates to population viability. In many species, rates of adult survival have a greater influence on population growth rates than do birth rates (Sæther & Bakke, 2000; Bruna, Fiske & Trager, 2009).

(2) Death rates

Although rates of survivorship in established plants usually contribute more to plant population growth rates than reproduction and seedling dynamics, there has been more focus on the effects of landscape condition on plant reproduction (Bruna *et al.*, 2009). There have been few studies on the effects of landscape condition on plant death rates, but mortality increases in many species due to transformation of native forest to plantations (Jules, 1998), and increased wind turbulence and microclimate changes near vegetation boundaries with agricultural land (Laurance *et al.*, 1998; Werner, 2011).

Elevated death rates in changed landscapes may reduce population viability for animal species (Harper *et al.*, 2008; Li *et al.*, 2009). Diminished availability of resources can contribute to higher death rates in fragmented landscapes and in small vegetation remnants (Boudjemadi *et al.*, 1999; Doherty & Grubb, 2002). Death rates may be affected by higher predation and desiccation in degraded or cleared vegetation (Rothermel & Semlitsch, 2002; Harper *et al.*, 2008), including during dispersal (Cushman, 2006). Mortality during dispersal through much-modified landscapes affects sex ratios, birth rates (Banks *et al.*, 2005) and the persistence of populations (Brooker & Brooker, 2002).

High temperatures and heat waves (Jakalanemi, 2011; Andreello *et al.*, 2012) and low rainfall and drought (Toräng, Ehrlén & Ågren, 2010) increase plant death rates through physiological stress. Drought increases susceptibility and exposure to pest species that cause mortality (Kloeppel *et al.*, 2003). Mortality of trees from increased drought occurs in many forests around the world and is expected to become more frequent (Van Mantgem & Stephenson, 2007; Horner *et al.*, 2009).

Warmer temperatures and low rainfall can accelerate water loss and energy expenditure in animals, leading to chronic stress, desiccation or hyperthermia (Grafe *et al.*, 2004; Moses, Frey & Roemer, 2012), particularly if these climate changes occur during energetically demanding phases of a species annual cycle (Grosbois *et al.*, 2006), or if temperatures approach or exceed the upper lethal limit of a species (Bale & Hayward, 2010). High temperatures increase population death rates (Grosbois *et al.*, 2006; Griffiths, Sewell & McCrea, 2010) and the frequency of catastrophic mortality events (McKechnie & Wolf, 2010). While increased temperatures may improve survival rates in some animals that experience cold stress, earlier melting of protective snow layers increases death rates by exposing animals to deleterious weather conditions, such as freezing rain and cool air temperatures (Bale & Hayward, 2010; Fisher & Davis, 2011) and increases predation risk (Lindström & Hörnfeldt, 1994). In cooler climates, elevated temperatures may increase survival rates for organisms near their lower thermal limits (Walther *et al.*, 2002; Frenot *et al.*, 2005). Asynchronies in the life cycles of predator and prey may increase the survival of the prey species, particularly if the prey is limited by predation rather than by food availability (Miller-Rushing *et al.*, 2010).

Increased frequency of high-energy weather events, such as hurricanes, storms and heavy rainfall, increase death rates in plants (Van Mantgem & Stephenson, 2007) and animals (Langtimm & Beck, 2003). Severe rain, snow or wind events cause mass mortality events (Newton, 2008a). Death rates increase with fewer food and foraging resources in the aftermath of intense weather events (Wiley & Wunderle, 1993).

Drought and much reduced rainfall can increase death rates through decreased food availability for terrestrial animals (Sillett *et al.*, 2000; Frick, Reynolds & Kunz, 2010), particularly when these occur during crucial times of breeding and survival. Climate oscillations affect food availability, and therefore death rates (Sandvik *et al.*, 2005; Morrison *et al.*, 2011).

While we have detailed several predicted and observed effects of both landscape condition and climate change on mortality, there has been relatively little research that measures the effects of these processes on death rates, and their subsequent effect on population viability. A better understanding of the effects of major anthropogenic pressures on death rates will be particularly important for those species whose population viability is most acutely affected by death rates, such as long-lived species (Sæther & Bakke, 2000).

(3) Emigration and immigration

Given that adult terrestrial plants are sedentary, emigration and immigration mostly is through the transport of seeds, fruits or vegetative propagules by animals, wind or water (Raulings *et al.*, 2011) and does not constitute the loss of an adult from the donor population *per se*. Increased isolation of plant populations and declines in seed-disperser populations (Cordeiro & Howe, 2003) inhibit biotic and abiotic seed dispersal, particularly for heavy-seeded species (Hewitt &

Kellman, 2002; McEuen & Curran, 2004), with potentially substantial effects on population viability (Hewitt & Kellman, 2002). Gene flow of plants predominantly is through the dispersal of pollen by biotic vectors and physical transmission (Ellstrand, 1992), which can be impeded by declines in landscape condition and climate change (Section IV.1).

The loss, fragmentation and degradation of native vegetation increase emigration rates and decrease immigration rates in animal populations, which affect population size and hence population viability, but the evidence for these expectations is weak (Section III). Reduced immigration can lead to skewed sex ratios (Harrisson *et al.*, 2012), inbreeding (Daniels, Priddy & Walters, 2000), disruption of mating systems (Pavlova *et al.*, 2012) and mate limitations (Stow & Sunnucks, 2004), which decrease population viability.

Low emigration rates generally occur when habitat and resources are ample (Baguette, Petit & Queva, 2000). If a site is rich in resources, immigration is likely to be higher because the immigrants are attracted by the presence of numerous conspecifics (Buechner, 1987) and highly suitable habitat (for the species) increases the 'attractiveness' of sites for recolonizing individuals (Doerr, Doerr & Jenkins, 2006).

Populations in high-intensity human land-use areas or that are experiencing low resource availability are more likely to experience emigration, and, in extreme circumstances, this can cause extinction (Lin & Batzli, 2001; Mac Nally *et al.*, 2009). Individuals are more likely to emigrate if they experience low reproductive or pairing success (Bayne & Hobson, 2002; Zitske, Betts & Diamond, 2011).

Small and isolated vegetation remnants generally attract fewer immigrants (Wauters *et al.*, 1994; Holland & Bennett, 2010). Decreased dispersal success caused by death during dispersal or the inability to locate appropriate habitat in high-intensity land-use areas lowers immigration rates (Matthysen, 1999; Püttker *et al.*, 2011) and reduces population viability, even in mobile animals, such as birds (Cooper & Walters, 2002; Robles *et al.*, 2008). Measurements of genetic connectivity among populations suggest decreases in dispersal in fragmented landscapes (Vos *et al.*, 2001). These measures, when combined with direct measures of movement, have the potential to help tease out the effects of landscape condition and other pressures on immigration and emigration rates (Lowe & Allendorf, 2010).

Warmer temperatures can increase animal emigration rates (Pärn *et al.*, 2011; Franzén & Nilsson, 2012) and dispersal distances (Cormont *et al.*, 2011), but may cause disparities in dispersal between the sexes (Merckx, Karlsson & Van Dyck, 2006). Increased atmospheric instability caused by warmer temperatures induces long-distance wind dispersal of seeds (Kuparinen *et al.*, 2009) and small invertebrates (Coulson *et al.*, 2002) by increasing convective turbulent airflow. Warmer temperatures may discourage juvenile dispersal (Massot, Clobert & Ferrière, 2008) and increase dispersal mortality due to heat stress (Henry, Sim & Russello, 2012). Lower rainfall can decrease vegetation quality in high-intensity land-use areas, discouraging

emigration between fragments of native vegetation (Blaum *et al.*, 2012). Climatic events such as El Niño Southern Oscillation (ENSO) phases and consequent declines in food resources may trigger irruptive migrations of animals (Holmgren *et al.*, 2006; Lindén *et al.*, 2011).

Studies that use niche models to predict changes in species distributions predict elevational and latitudinal shifts in response to climate exposure, assuming colonization of newly suitable climate conditions (Fordham *et al.*, 2012). The structure and condition of many human-dominated landscapes are likely to impede colonization (Opdam & Wascher, 2004). Although organisms have responded to climate changes through migration and adaptation in the past, the barriers imposed by human land use and the unprecedented rate of climate change are unlikely to allow the predicted range shifts in many species to occur (Davis & Shaw, 2001). Range shifts are inhibited in much-modified landscapes, and may stall where the amount or cohesion of habitat is below thresholds necessary for population persistence (Opdam & Wascher, 2004). Fragmented vegetation may be disproportionately affected (higher mortality or die-back) by climate change (Bennett *et al.*, 2013), creating further barriers to climate-induced range shifts. Some species may be unimpeded by modified landscapes and this will affect species interactions in receiving habitats (Menéndez *et al.*, 2008). For example, landscape and climate change have increased the distribution and abundance of the despotic noisy miner (*Manorina melanocephala*) in eastern Australia. This has caused local emigration and a lack of immigration of small-bodied birds in fragmented vegetation where the species is present (Maron *et al.*, 2013).

To gauge the effects of climate change on species distributions, an understanding of the effects of climate on immigration and emigration rates and the processes of dispersal is vital, particularly in changed landscapes where these rates are already affected.

V. SYNTHESIS AND FUTURE WORK

Demographic rates are rarely the focus of studies on the effects of human pressures on native populations of plants and animals. However, these effects can be substantial and their identification enables a better understanding of the mechanisms through which pressures affect population viability. That vegetation loss, fragmentation and degradation affect demographic rates in plant and animal populations is not unexpected given the widespread declines in biodiversity that have been seen as a result of these pressures (Foley *et al.*, 2005; Butchart *et al.*, 2010). Our finding that the mean effect of climate on demographic rates is of comparable magnitude to changes in landscape condition is significant and supports recent assertions that climate change will become as, or more, important in species declines and extinctions in coming decades (e.g. Mantyka-Pringle *et al.*, 2012).

The relative effects of climate on demographic rates probably are underestimates. Most studies assess

relationships between general climate measures, such as annual temperature or seasonal precipitation within average ranges of year-to-year variation. The characteristics of relationships between demographic variables and climate variations are likely to change once changes in climate fall outside the average range. The effects of climate variables on demographic rates may become greater, new effects may emerge, or the direction of relationships may change. There are likely to be critical windows of climate effects on population parameters, where climate conditions at very specific times in species life cycles are disproportionately important to population viability (Lada *et al.*, 2013). Assessing general trends in climate and demographic rates may not detect the true size of the effects on population viability that will occur if changes in climate occur during critical windows. Critical thresholds may exist, such as where temperatures exceed lethal limits (Somero, 2010). Studies that measure demographic responses to climate conditions within the average range are unlikely to detect such responses. While the studies we reviewed assumed monotonic relationships between pressures and demographic variables, physiological responses to temperature are commonly asymmetric, such that a positive response to temperature may be reversed once an optimal level is reached (Sinclair & Chown, 2003).

Climate change may introduce new pressures to otherwise viable populations, or may cause the decline of populations in changed landscapes faster than otherwise expected. Decreases in rainfall and increases in temperature probably will have deleterious effects on many populations, although some taxa almost certainly will benefit. Small populations have less capacity to evolve rapidly to changed conditions (Willi, Buskirk & Hoffmann, 2006), so climate change may have a cascading effect on the viability of populations that have been affected by changed landscape condition. Some species will have increased population viability with the amelioration of limiting climate conditions. Changes in population viability in either direction will affect species interactions, with disruptions for communities (Sorte & White, 2013). A greater focus on the relationships between climate conditions and demographic rates is needed to produce better predictions for likely impacts of climate changes on animal and plant populations. A more complete understanding of the effects on immigration and emigration must improve predictions of range shifts. Identification of the demographic rates most affected by projected climate changes will assist with better planning for climate adaptation.

Populations in changed landscapes may decline faster than expected with the added pressures arising from climate change. This is important in making predictions about population size in response to pressures such as habitat loss, including considerations of critical thresholds (Swift & Hannon, 2010), which could be reached earlier than expected with the added imposts of climate pressures and their effects may be synergistic (Mantyka-Pringle *et al.*, 2012). Landscape modification may hinder or reverse the expected

population growth in response to changed climate conditions (Warren *et al.*, 2001). Whether the effects of landscape condition and climate on demographic rates are additive or multiplicative (or for some species, opposing), is a core question.

While our review highlights some mechanisms through which the major anthropogenic pressures affect population viability, there is a clear need for more data. A more comprehensive understanding of these relationships will contribute greatly to improving the effectiveness of conservation policies and management actions. Specifically, there is a need for expanding research beyond North America and Europe, and we suggest that the most important areas for conducting this research are those that are predicted to experience the greatest changes in climate conditions. Warming is likely to occur most rapidly in the polar regions, while mid-latitude and sub-tropical dry regions are likely to be most affected by decreased precipitation (IPCC, 2013). There is a dearth of research into the effects of climate on plant demographic rates despite climate change being the most commonly cited factor in the extinction and endangerment of plant species (Mora & Zapata, 2013).

VI. CONCLUSIONS

(1) Given their intimate connection with population viability, demographic responses provide a critical indication of likely changes in extinction risk in response to human pressures.

(2) Changes in landscape condition generally have a negative effect on birth and immigration rates in plant and animal populations, and increase death and emigration rates. We predict that climate change will have a negative effect on birth and immigration rates, and a positive effect on death and emigration rates, although we did not quantitatively assess this.

(3) Despite the recognition of landscape change as the major driver of biodiversity loss, the effects of climate on demographic rates in plant and animal populations are of equivalent magnitude. This supports consideration of climate change as a major driver of population viability, of similar importance to human land-use change.

(4) A more comprehensive understanding of the rate and size of the effects of pressures on demographic rates among taxa and regions will greatly assist management attempts to arrest species declines and extinctions.

VII. ACKNOWLEDGEMENTS

R.M. was supported partially by an Australian Research Council grant (DP120100797). J. Bennett contributed to development of Fig. 3. P. Sunnucks, H.A. Ford and J.D. Roberts provided valuable comments that much improved the manuscript.

VIII. REFERENCES

- *ADAMS, R. A. (2010). Bat reproduction declines when conditions mimic climate change projections for western North America. *Ecology* **91**, 2437–2445.
- ÅGREN, J., EHRLÉN, J. & SOLBRECK, C. (2008). Spatio-temporal variation in fruit production and seed predation in a perennial herb influenced by habitat quality and population size. *Journal of Ecology* **96**, 334–345.
- AGUILAR, R., ASHWORTH, L., GALETTO, L. & AIZEN, M. A. (2006). Plant reproductive susceptibility to habitat fragmentation: review and synthesis through a meta-analysis. *Ecology Letters* **9**, 968–980.
- *AGUILAR, R. & GALETTO, L. (2004). Effects of forest fragmentation on male and female reproductive success in *Cestrum parqui* (Solanaceae). *Oecologia* **138**, 213–220.
- *AIZEN, M. A. & FEINSINGER, P. (1994). Forest fragmentation, pollination, and plant reproduction in a Chaco dry forest, Argentina. *Ecology* **75**, 330–351.
- AKÇAKAYA, H. R., BUTCHART, S. H. M., MACE, G. M., STUART, S. N. & HILTON-TAYLOR, C. (2006). Use and misuse of the IUCN Red List Criteria in projecting climate change impacts on biodiversity. *Global Change Biology* **12**, 2037–2043.
- ALTERMATT, F., PAJUNEN, V. I. & EBERT, D. (2008). Climate change affects colonization dynamics in a metacommunity of three *Daphnia* species. *Global Change Biology* **14**, 1209–1220.
- ANDREILLO, M., BIZOUX, J.-P., BARBET-MASSIN, M., GAUDEUL, M., NICOLÉ, F. & TILL-BOTTRAUD, I. (2012). Effects of management regimes and extreme climatic events on plant population viability in *Eryngium alpinum*. *Biological Conservation* **147**, 99–106.
- BAGUETTE, M., PETIT, S. & QUEVA, F. (2000). Population spatial structure and migration of three butterfly species within the same habitat network: consequences for conservation. *Journal of Applied Ecology* **37**, 100–108.
- BALE, J. S. & HAYWARD, S. A. L. (2010). Insect overwintering in a changing climate. *Journal of Experimental Biology* **213**, 980–994.
- BANKS, S. C., FINLAYSON, G. R., LAWSON, S. J., LINDENMAYER, D. B., PAETKAU, D., WARD, S. J. & TAYLOR, A. C. (2005). The effects of habitat fragmentation due to forestry plantation establishment on the demography and genetic variation of a marsupial carnivore, *Antechinus agilis*. *Biological Conservation* **122**, 581–597.
- BANNISTER, P., MAEGLI, T., DICKINSON, K. J., HALLOY, S. R., KNIGHT, A., LORD, J. M., MARK, A. F. & SPENCER, K. L. (2005). Will loss of snow cover during climatic warming expose New Zealand alpine plants to increased frost damage? *Oecologia* **144**, 245–256.
- BAYNE, E. M. & HOBSON, K. A. (2002). Apparent survival of male ovenbirds in fragmented and forested boreal landscapes. *Ecology* **83**, 1307–1316.
- BEGON, M., MORTIMER, M. & THOMPSON, D. J. (1996). Single-species populations. In *Population Ecology: A Unified Study of Animals and Plants*, pp. 2–27. Blackwell Science, Oxford.
- BEISSINGER, S. R., COOK, M. I. & ARENDT, W. J. (2005). The shelf life of bird eggs: testing egg viability using a tropical climate gradient. *Ecology* **86**, 2164–2175.
- BELLARD, C., BERTELSMIEIER, C., LEADLEY, P., THUILLER, W. & COURCHAMP, F. (2012). Impacts of climate change on the future of biodiversity. *Ecology Letters* **15**, 365–377.
- BENITEZ-MALVIDO, J. (1998). Impact of forest fragmentation on seedling abundance in a tropical rain forest. *Conservation Biology* **12**, 380–389.
- BENNETT, J. M., CUNNINGHAM, S. C., CONNELLY, C. A., CLARKE, R. H., THOMSON, J. R. & MAC NALLY, R. (2013). The interaction between a drying climate and land use affects forest structure and above-ground carbon storage. *Global Ecology and Biogeography* **22**, 1238–1247.
- *BERVEN, K. A. (1990). Factors affecting population fluctuation in larval and adult stages of the wood frog (*Rana sylvatica*). *Ecology* **71**, 1599–1608.
- *BIGGER, D. S. (1999). Consequences of patch size and isolation for a rare plant: pollen limitation and seed predation. *Natural Areas Journal* **19**, 239–244.
- *BIONDA, R. & BRAMBILLA, M. (2012). Rainfall and landscape features affect productivity in an alpine population of Eagle Owl *Bubo bubo*. *Journal of Ornithology* **153**, 167–171.
- BLAUM, N., SCHWAGER, M., WICHMANN, M. C. & ROSSMANITH, E. (2012). Climate induced changes in matrix suitability explain gene flow in a fragmented landscape – the effect of interannual rainfall variability. *Ecography* **35**, 650–660.
- BLAUSTEIN, A. R., SEARLE, C., BANCROFT, B. A. & LAWLER, J. (2012). Amphibian population declines and climate change. In *Ecological Consequences of Climate Change: Mechanisms, Conservation, and Management* (eds E. A. BEEVER and J. L. BELANT), pp. 30–46. CRC Press, Boca Raton.
- *BOSCH, M., SIMON, J., ROVIRA, A. M., MOLERO, J. & BLANCHE, C. (2002). Pollination ecology of the pre-Pyrenean endemic *Petrocoptis montsicciana* (Caryophyllaceae): effects of population size. *Biological Journal of the Linnean Society* **76**, 79–90.
- BOUDJEMADI, K., LECOMTE, J. & CLOBERT, J. (1999). Influence of connectivity on demography and dispersal in two contrasting habitats: an experimental approach. *Journal of Animal Ecology* **68**, 1207–1224.

- *BOWERS, M. A., GREGARIO, K., BRAME, C. J., MATTER, S. F. & DOOLEY, J. L. (1996). Use of space and habitats by meadow voles at the home range, patch and landscape scales. *Oecologia* **105**, 107–115.
- BOYCE, M. S. (1992). Population viability analysis. *Annual Review of Ecology and Systematics* **23**, 481–506.
- BRISCOE, N. J., PORTER, W. P., SUNNUCKS, P. & KEARNEY, M. R. (2012). Stage-dependent physiological responses in a butterfly cause non-additive effects on phenology. *Oikos* **121**, 1464–1472.
- BRODIE, J., POST, E., WATSON, F. & BERGER, J. (2012). Climate change intensification of herbivore impacts on tree recruitment. *Proceedings of the Royal Society of London Series B: Biological Sciences* **279**, 1366–1370.
- *BROOKER, M. & BROOKER, L. (2001). Breeding biology, reproductive success and survival of blue-breasted fairy-wrens in fragmented habitat in the Western Australian wheatbelt. *Wildlife Research* **28**, 205–214.
- BROOKER, L. C. & BROOKER, M. G. (2012). Dispersal and population dynamics of the blue-breasted fairy-wren, *Malurus pulcherrimus*, in fragmented habitat in the Western Australian wheatbelt. *Wildlife Research* **29**, 225–233.
- BRUNA, E. M., FISKE, I. J. & TRAGER, M. D. (2009). Habitat fragmentation and plant populations: is what we know demographically irrelevant? *Journal of Vegetation Science* **20**, 569–576.
- *BRUNA, E. M. & KRESS, W. J. (2002). Habitat fragmentation and the demographic structure of an Amazonian understory herb (*Heliconia acuminata*). *Conservation Biology* **16**, 1256–1266.
- BRUNA, E. M. & OLI, M. K. (2005). Demographic effects of habitat fragmentation on a tropical herb: life-table response experiments. *Ecology* **86**, 1816–1824.
- *BRYN, R., JACQUEMYN, H., ENDELS, P., VAN ROSSUM, F., HERMY, M., TRIEST, L., DE BRUYN, L. & BLUST, G. D. E. (2004). Reduced reproductive success in small populations of the self-incompatible. *Journal of Ecology* **92**, 5–14.
- BUCKLEY, L. B., NUÑO, C. R. & KINGSOLVER, J. G. (2013). Phenotypic clines, energy balances and ecological responses to climate change. *Journal of Animal Ecology* **82**(4), 894–902.
- BUECHNER, M. (1987). Conservation in insular parks: Simulation models of factors affecting the movement of animals across park boundaries. *Biological Conservation* **41**, 57–76.
- *BURHANS, D. E. & THOMPSON, F. R. III (1999). Habitat patch size and nesting success of yellow-breasted chats. *Wilson Bulletin* **111**, 210–215.
- *BURKE, D. M. & NOL, E. (2000). Landscape and fragment size effects on reproductive success of forest-breeding birds in Ontario. *Ecological Applications* **10**, 1749–1761.
- BUTCHART, S. H. M., WALPOLE, M., COLLEN, B., VAN STRIEN, A., SCHARLEMANN, J. P. W., ALMOND, R. E. A., BAILLIE, J. E. M., BOMHARD, B., BROWN, C., BRUNO, J., CARPENTER, K. E., CARR, G. M., CHANSON, J., CHENERY, A. M., CSIRKE, J., DAVIDSON, N. C., DENTENER, F., FOSTER, M., GALLI, A., GALLOWAY, J. N., GENOVESI, P., GREGORY, R. D., HOCKINGS, M., KAPOV, V., LAMARQUE, J.-F., LEVERINGTON, F., LOH, J., MCGEOCH, M. A., MCRAE, L., MINASYAN, A., MORCILLO, M. H., OLDFIELD, T. E. E., PAULY, D., QUADER, S., REVENGA, C., SAUER, J. R., SKOLNIK, B., SPEAR, D., STANWELL-SMITH, D., STUART, S. N., SYMES, A., TIERNEY, M., TYRRELL, T. D., VIÉ, J.-C. & WATSON, R. (2010). Global biodiversity: indicators of recent declines. *Science* **328**, 1164–1168.
- *CAMARERO, J. J. & GUTIÉRREZ, E. (2007). Response of *Pinus uncinata* recruitment to climate warming and changes in grazing pressure in an isolated population of the Iberian system (NE Spain). *Arctic, Antarctic, and Alpine Research* **39**, 210–217.
- *CAMERON, M. (2009). The influence of climate on Glossy Black-cockatoo reproduction. *Pacific Conservation Biology* **15**, 65–71.
- CAMPBELL, R. D., NOUVELLET, P., NEWMAN, C., MACDONALD, D. W. & ROSELL, F. (2012). The influence of mean climate trends and climate variance on beaver survival and recruitment dynamics. *Global Change Biology* **18**, 2730–2742.
- *CASCANTE, A., QUESADA, M., LOBO, J. J. & FUCHS, E. A. (2002). Effects of dry tropical forest fragmentation on the reproductive success and genetic structure of the tree *Samanea saman*. *Conservation Biology* **16**, 137–147.
- CHAMBERS, L. E. & KEATLEY, M. R. (2010). Phenology and climate—early Australian botanical records. *Australian Journal of Botany* **58**, 473–484.
- *CHAMBERS, L. E., QUIN, B. R., MENKHORST, P., FRANKLIN, D. C. & SMALES, I. (2008). The effects of climate on breeding in the Helmeted Honeyeater. *Emu* **108**, 15–22.
- CHEN, I.-C., HILL, J. K., OHLEMÜLLER, R., ROY, D. B. & THOMAS, C. D. (2011). Rapid range shifts of species associated with high levels of climate warming. *Science* **333**, 1024–1026.
- CHIDUMAYO, E. N. (2008). Implications of climate warming on seedling emergence and mortality of African savanna woody plants. *Plant Ecology* **198**, 61–71.
- CHOWN, S., HOFFMANN, A., KRISTENSEN, T., ANGILETTA, M. J., STENSETH, N. & PERTOLDI, C. (2010). Adapting to climate change: a perspective from evolutionary physiology. *Climate Research* **43**, 3–15.
- CLARKE, D. N. & ZANI, P. A. (2012). Effects of night-time warming on temperate ectotherm reproduction: potential fitness benefits of climate change for side-blotched lizards. *Journal of Experimental Biology* **215**, 1117–1127.
- COHEN, J. E. (1969). Natural primate troops and a stochastic population model. *American Naturalist* **103**, 455–477.
- *CONDIT, R., HUBBELL, S. P. & FOSTER, R. B. (1995). Mortality rates of 205 neotropical tree and shrub species and the impact of a severe drought. *Ecological Monographs* **65**, 419–439.
- COOPER, C. B. & WALTERS, J. R. (2002). Experimental evidence of disrupted dispersal causing decline of an Australian passerine in fragmented habitat. *Conservation Biology* **16**, 471–478.
- CORDEIRO, N. J. & HOWE, H. F. (2003). Forest fragmentation severs mutualism between seed dispersers and an endemic African tree. *Proceedings of the National Academy of Sciences of the United States of America* **100**, 14052–14056.
- CORMONT, A., MALINOWSKA, A. H., KOSTENKO, O., RADCHUK, V., HEMERIK, L., WALLISDEVRIES, M. F. & VERBOOM, J. (2011). Effect of local weather on butterfly flight behaviour, movement, and colonization: significance for dispersal under climate change. *Biodiversity and Conservation* **20**, 483–503.
- *COSTIN, B. J., MORGAN, J. W. & YOUNG, A. G. (2001). Reproductive success does not decline in fragmented populations of *Leucochrysum albicans* subsp. *albicans* var. *tricolor* (Asteraceae). *Biological Conservation* **98**, 273–284.
- COULSON, S. J., HODKINSON, I. D., WEBB, N. R., MIKKOLA, K., HARRISON, J. A. & PEDGLEY, D. E. (2002). Aerial colonization of high Arctic islands by invertebrates: the diamondback moth *Plutella xylostella* (Lepidoptera: Yponomeutidae) as a potential indicator species. *Diversity and Distributions* **8**, 327–334.
- CUNNINGHAM, S. A. (2000a). Depressed pollination in habitat fragments causes low fruit set. *Proceedings of the Royal Society of London Series B: Biological Sciences* **267**, 1149–1152.
- *CUNNINGHAM, S. A. (2000b). Effects of habitat fragmentation on the reproductive ecology of four plant species in mallee woodland. *Conservation Biology* **14**, 758–768.
- CUNNINGHAM, S. C., MACNALLY, R., READ, J., BAKER, P. J., WHITE, M., THOMSON, J. R. & GRIFFOEN, P. (2009). A robust technique for mapping vegetation condition across a major river system. *Ecosystems* **12**, 207–219.
- CUSHMAN, S. A. (2006). Effects of habitat loss and fragmentation on amphibians: a review and prospectus. *Biological Conservation* **128**, 231–240.
- *DALE, S. (2004). Effects of a golf course on population dynamics of the endangered ortolan bunting. *Journal of Wildlife Management* **68**, 719–724.
- DANIELS, S. J., PRIDDY, J. A. & WALTERS, J. R. (2000). Inbreeding in small populations of red-cockaded woodpeckers: insights from a spatially explicit individual-based model. In *Genetics, Demography and Viability of Fragmented Populations* (eds A. G. YOUNG and G. M. CLARKE), pp. 129–148. Cambridge University Press, Cambridge.
- DAVIS, M. B. & SHAW, R. G. (2001). Range shifts and adaptive responses to quaternary climate change. *Science* **292**, 673–679.
- DAWSON, T. P., JACKSON, S. T., HOUSE, J. I., PRENTICE, I. C. & MACE, G. M. (2011). Beyond predictions: biodiversity conservation in a changing climate. *Science* **332**, 53–58.
- DEBINSKI, D. M. & HOLT, R. D. (2000). Review: a survey and overview of habitat fragmentation experiments. *Conservation Biology* **14**, 342–355.
- DE VEAUX, R. D., VELLEMAN, P. F. & BOCK, D. E. (2008). *Stats: Data and Models*. Pearson Education, Boston.
- *DICK, C. W. (2001). Genetic rescue of remnant tropical trees by an alien pollinator. *Proceedings of the Royal Society of London Series B: Biological Sciences* **268**, 2391–2396.
- DOERR, V. A. J., DOERR, E. D. & JENKINS, S. H. (2006). Habitat selection in two Australasian treecreepers: what cues should they use? *Emu* **106**, 93–103.
- DOHERTY, P. F. & GRUBB, T. C. (2002). Survivorship of permanent-resident birds in a fragmented forested landscape. *Ecology* **83**, 844–857.
- *DONALDSON, J. S., NANNI, I., ZACHARIADES, C. & KEMPER, J. (2002). Effects of habitat fragmentation on pollinator diversity and plant reproductive success in renosterveld shrublands of South Africa. *Conservation Biology* **16**, 1267–1276.
- *DONELSON, J. M., MUNDAY, P. L., MCCORMICK, M. I., PANKHURST, N. W. & PANKHURST, P. M. (2010). Effects of elevated water temperature and food availability on the reproductive performance of a coral reef fish. *Marine Ecology Progress Series* **401**, 233–243.
- *DONOVAN, T. M., THOMPSON, F. R. III, FAABORG, J. & PROBST, J. R. (1995). Reproductive success of migratory birds in habitat sources and sinks. *Conservation Biology* **9**, 1380–1395.
- *DOOLEY, J. L. & BOWERS, M. A. (1998). Demographic responses to habitat fragmentation: experimental tests at the landscape and patch scale. *Ecology* **79**, 969–980.
- DUNHAM, A. E., ERHART, E. M. & WRIGHT, P. C. (2010). Global climate cycles and cyclones: consequences for rainfall patterns and lemur reproduction in southeastern Madagascar. *Global Change Biology* **17**, 219–227.
- DURANT, J. M., HJERMANN, D. Å., OTTERSEN, G. & STENSETH, N. C. (2007). Climate and the match or mismatch between predator requirements and resource availability. *Climate Research* **33**, 271–283.
- ELLSTRAND, N. C. (1992). Gene flow by pollen: implications for plant conservation genetics. *Oikos* **63**, 77–86.
- FAHRIG, L. (1997). Relative effects of habitat loss and fragmentation on population extinction. *Journal of Wildlife Management* **61**, 603–610.
- *FERGUSON, S. H., STIRLING, I. & MCLOUGHLIN, P. (2005). Climate change and ringed seal (*Phoca hispida*) recruitment in western Hudson Bay. *Marine Mammal Science* **21**, 121–135.

- FICKE, A. D., MYRICK, C. A. & HANSEN, L. J. (2007). Potential impacts of global climate change on freshwater fisheries. *Reviews in Fish Biology and Fisheries* **17**, 581–613.
- FISCHER, J. & LINDENMAYER, D. B. (2007). Landscape modification and habitat fragmentation: a synthesis. *Global Ecology and Biogeography* **16**, 265–280.
- FISHER, R. J. & DAVIS, S. K. (2011). Post-fledging dispersal, habitat use, and survival of Sprague's pipits: are planted grasslands a good substitute for native? *Biological Conservation* **144**, 263–271.
- FODEN, W. B., BUTCHART, S. H. M., STUART, S. N., VIÉ, J.-C., AKÇAKAYA, H. R., ANGULO, A., DEVANTIER, L. M., GUTSCHE, A., TURAK, E., CAO, L., DONNER, S. D., KATARIYA, V., BERNARD, R., HOLLAND, R. A., HUGHES, A. F., O'HANLON, S. E., GARNETT, S. T., ŞEKERCIOĞLU, Ç. H. & MACE, G. M. (2013). Identifying the world's most climate change vulnerable species: a systematic trait-based assessment of all birds, amphibians and corals. *PLoS ONE* **8**, e65427.
- FOLEY, J. A., DEFRIES, R., ASNER, G. P., BARFORD, C., BONAN, G., CARPENTER, S. R., CHAPIN, F. S., COE, M. T., DAILY, G. C., GIBBS, H. K., HELKOWSKI, J. H., HOLLOWAY, T., HOWARD, E. A., KUCHARIK, C. J., MONFREDA, C., PATZ, J. A., PRENTICE, I. C., RAMANKUTTY, N. & SNYDER, P. K. (2005). Global consequences of land use. *Science* **309**, 570–574.
- FORDHAM, D. A., RESIT AKÇAKAYA, H., ARAÚJO, M. B., ELITH, J., KEITH, D. A., PEARSON, R., AULD, T. D., MELLIN, C., MORGAN, J. W., REGAN, T. J., TOZER, M., WATTS, M. J., WHITE, M., WINTLE, B. A., YATES, C. & BROOK, B. W. (2012). Plant extinction risk under climate change: are forecast range shifts alone a good indicator of species vulnerability to global warming? *Global Change Biology* **18**, 1357–1371.
- FRANKHAM, R. (1995). Effective population size/adult population size ratios in wildlife: a review. *Genetics Research* **66**, 95–107.
- FRANZEN, M. & NILSSON, S. G. (2012). Climate-dependent dispersal rates in metapopulations of burnet moths. *Journal of Insect Conservation*, **16**(6), 1–7.
- FRENOT, Y., CHOWN, S. L., WHINAM, J., SELKIRK, P. M., CONVEY, P., SKOTNICKI, M. & BERGSTROM, D. M. (2005). Biological invasions in the Antarctic: extent, impacts and implications. *Biological Reviews* **80**, 45–72.
- FRICK, W. F., REYNOLDS, D. S. & KUNZ, T. H. (2010). Influence of climate and reproductive timing on demography of little brown myotis *Myotis lucifugus*. *Journal of Animal Ecology* **79**, 128–136.
- *FUCHS, E. A., LOBO, J. A. & QUESADA, M. (2003). Effects of forest fragmentation and flowering phenology on the reproductive success and mating patterns of the tropical dry forest tree *Pachira quinata*. *Conservation Biology* **17**(1), 149–157.
- GASTON, A. J., GILCHRIST, H. G. & HIPFNER, J. M. (2005). Climate change, ice conditions and reproduction in an Arctic nesting marine bird: Brunnich's guillemot (*Uria lomvia* L.). *Journal of Animal Ecology* **74**, 832–841.
- *GATES, J. E. & GYSEL, L. W. (1978). Avian nest dispersion and fledging success in field-forest ecotones. *Ecology* **59**, 871–883.
- *GHAZOUL, J. & MCLEISH, M. (2001). Reproductive ecology of tropical forest trees in logged and fragmented habitats in Thailand and Costa Rica. *Plant Ecology* **153**, 335–345.
- *GIBBS, J. P. & FAABORG, J. (1990). Estimating the viability of ovenbird and Kentucky warbler populations in forest fragments. *Conservation Biology* **4**, 193–196.
- *GIGORD, L., PICOT, F. & SHYKOFF, J. A. (1999). Effects of habitat fragmentation on *Dombeya acutangula* (Sterculiaceae), a native tree on La Réunion (Indian Ocean). *Biological Conservation* **88**, 43–51.
- *GJERDRUM, C., VALLÉE, A. M. G., ST. CLAIR, C. C., BERTRAM, D. F., RYDER, J. L. & BLACKBURN, G. S. (2003). Tufted Puffin reproduction reveals ocean climate variability. *Proceedings of the National Academy of Sciences of the United States of America* **100**, 9377–9382.
- GLENN, E. M., ANTHONY, R. G., FORSMAN, E. D. & OLSON, G. S. (2011). Reproduction of northern spotted owls: the role of local weather and regional climate. *Journal of Wildlife Management* **75**, 1279–1294.
- GÓMEZ, J. M., ABDELAZIZ, M., LORITE, J., JESÚS MUÑOZ-PAJARES, A. & PERFECTTI, F. (2010). Changes in pollinator fauna cause spatial variation in pollen limitation. *Journal of Ecology* **98**, 1243–1252.
- GRAFE, T. U., KAMINSKY, S. K., BITZ, J. H., LÜSSOW, H. & LINSINMAIR, K. E. (2004). Demographic dynamics of the afro-tropical pig-nosed frog, *Hemisus marmoratus*: effects of climate and predation on survival and recruitment. *Oecologia* **141**, 40–46.
- GRIFFIN, K. A., HEBBLEWHITE, M., ROBINSON, H. S., ZAGER, P., BARBER-MEYER, S. M., CHRISTIANSON, D., CREEL, S., HARRIS, N. C., HURLEY, M. A., JACKSON, D. H., JOHNSON, B. K., MYERS, W. L., RAITHEL, J. D., SCHLEGEL, M., SMITH, B. L., WHITE, C. & WHITE, P. J. (2011). Neonatal mortality of elk driven by climate, predator phenology and predator community composition. *Journal of Animal Ecology* **80**, 1246–1257.
- GRIFFITHS, R. A., SEWELL, D. & MCCREA, R. S. (2010). Dynamics of a declining amphibian metapopulation: survival, dispersal and the impact of climate. *Biological Conservation* **143**, 485–491.
- *GROOM, M. J. (2001). Consequences of subpopulation isolation for pollination, herbivory, and population growth in *Clarkia concinna concinna* (Onagraceae). *Biological Conservation* **100**, 55–63.
- GROSBOIS, V., HENRY, P. Y., BLONDEL, J., PERRET, P., LEBRETON, J. D., THOMAS, D. W. & LAMBRECHTS, M. M. (2006). Climate impacts on Mediterranean blue tit survival: an investigation across seasons and spatial scales. *Global Change Biology* **12**, 2235–2249.
- HALLETT, L. M., STANDISH, R. J. & HOBBS, R. J. (2011). Seed mass and summer drought survival in a Mediterranean-climate ecosystem. *Plant Ecology* **212**, 1479–1489.
- HALPERN, B. S., WALBRIDGE, S., SELKOE, K. A., KAPPEL, C. V., MICHELI, F., D'AGROSA, C., BRUNO, J. F., CASEY, K. S., EBERT, C. & FOX, H. E. (2008). A global map of human impact on marine ecosystems. *Science* **319**, 948–952.
- HARGROVE, L. & ROTENBERRY, J. T. (2011). Breeding success at the range margin of a desert species: implications for a climate-induced elevational shift. *Oikos* **120**, 1568–1576.
- HARPER, E. B., RITTENHOUSE, T. A. G. & SEMLITSCH, R. D. (2008). Demographic consequences of terrestrial habitat loss for pool-breeding amphibians: predicting extinction risks associated with inadequate size of buffer zones. *Conservation Biology* **22**, 1205–1215.
- HARRISON, F. (2011). Getting started with meta-analysis. *Methods in Ecology and Evolution* **2**, 1–10.
- HARRISSON, K., PAVLOVA, A., AMOS, J. N., TAKEUCHI, N., LILL, A., RADFORD, J. & SUNNUCKS, P. (2012). Fine-scale effects of habitat loss and fragmentation despite large-scale gene flow for some regionally declining woodland bird species. *Landscape Ecology* **27**, 813–827.
- HELLER, N. E. & ZAVALA, E. S. (2009). Biodiversity management in the face of climate change: a review of 22 years of recommendations. *Biological Conservation* **142**, 14–32.
- HENRY, P., SIM, Z. & RUSSELLO, M. A. (2012). Genetic evidence for restricted dispersal along continuous altitudinal gradients in a climate change-sensitive mammal: the American Pika. *PLoS ONE* **7**, e39077.
- HEWITT, N. & KELLMAN, M. (2002). Tree seed dispersal among forest fragments: II. Dispersal abilities and biogeographical controls. *Journal of Biogeography* **29**, 351–363.
- *HINSLEY, S. A., ROTHERY, P. & BELLAMY, P. E. (1999). Influence of woodland area on breeding success in great tits *Parus major* and blue tits *Parus caeruleus*. *Journal of Avian Biology* **30**, 271–281.
- HOLLAND, G. J. & BENNETT, A. F. (2010). Habitat fragmentation disrupts the demography of a widespread native mammal. *Ecography* **33**, 841–853.
- HOLMGREN, M., STAPP, P., DICKMAN, C. R., GRACIA, C., GRAHAM, S., GUTIÉRREZ, J. R., HICE, C., JAKSIC, F., KELT, D. A., LETNIC, M., LIMA, M., LÓPEZ, B. C., MESERVE, P. L., MILSTEAD, W. B., POLIS, G. A., PREVITALI, M. A., RICHTER, M., SABATÉ, S. & SQUEO, F. A. (2006). Extreme climatic events shape arid and semiarid ecosystems. *Frontiers in Ecology and the Environment* **4**, 87–95.
- *HOOVER, J. P., BRITTINGHAM, M. C. & GOODRICH, L. J. (1995). Effects of forest patch size on nesting success of wood thrushes. *Auk* **112**, 146–155.
- HORNER, G. J., BAKER, P. J., MAC NALLY, R., CUNNINGHAM, S. C., THOMSON, J. R. & HAMILTON, F. (2009). Mortality of developing floodplain forests subjected to a drying climate and water extraction. *Global Change Biology* **15**, 2176–2186.
- INOUE, D. W. (2000). The ecological and evolutionary significance of frost in the context of climate change. *Ecology Letters* **3**, 457–463.
- INOUE, D. W. (2008). Effects of climate change on phenology, frost damage, and floral abundance of montane wildflowers. *Ecology* **89**, 353–362.
- IPCC (2013). Summary for policy makers. In *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (eds T. F. STOCKER, D. QIN, G.-K. PLATTNER, M. TIGNOR, S. K. ALLEN, J. BOSCHUNG, A. NAUJES, Y. XIA, V. BEX and P. M. MIDGLEY), Cambridge University Press, Cambridge and New York.
- *JACQUEMYN, H., BRYNS, R. & HERMY, M. (2002). Patch occupancy, population size and reproductive success of a forest herb (*Primula elatior*) in a fragmented landscape. *Oecologia* **130**, 617–625.
- JACQUEMYN, H., VAN ROSSUM, F., BRYNS, R., ENDELS, P., HERMY, M., TRIEST, L. & DE BLUST, G. (2003). Effects of agricultural land use and fragmentation on genetics, demography and population persistence of the rare *Primula vulgaris*, and implications for conservation. *Belgian Journal of Botany* **136**, 5–22.
- JAKALANIEMI, A. (2011). Narrow climate and habitat envelope affect the survival of relict populations of a northern *Arnica angustifolia*. *Environmental and Experimental Botany* **72**, 415–421.
- JANSEN, A. & ROBERTSON, A. I. (2001). Relationships between livestock management and the ecological condition of riparian habitats along an Australian floodplain river. *Journal of Applied Ecology* **38**, 63–75.
- *JENNERSTEN, O. L. A. (1988). Pollination in *Dianthus deltoides* (Caryophyllaceae): effects of habitat fragmentation on visitation and seed set. *Conservation Biology* **2**, 359–366.
- *JENOUVRIER, S., BARBRAUD, C. & WEIMERSKIRCH, H. (2003). Effects of climate variability on the temporal population dynamics of southern fulmars. *Journal of Animal Ecology* **72**, 576–587.
- *JOHNSON, S. D., NEAL, P. R., PETER, C. I. & EDWARDS, T. J. (2004). Fruiting failure and limited recruitment in remnant populations of the hawkmoth-pollinated tree *Oxyanthus pyriflorus* subsp. *pyriflorus* (Rubiaceae). *Biological Conservation* **120**, 31–39.
- JÖNSSON, A. M., APPELBERG, G., HARDING, S. & BÄRRING, L. (2009). Spatio-temporal impact of climate change on the activity and voltinism of the spruce bark beetle, *Ips typographus*. *Global Change Biology* **15**, 486–499.

- *JULES, E. S. (1998). Habitat fragmentation and demographic change for a common plant: Trillium in old-growth forest. *Ecology* **79**, 1645–1656.
- JULES, E. S. & RATHCKE, B. J. (1999). Mechanisms of reduced Trillium recruitment along edges of old-growth forest fragments. *Conservation Biology* **13**, 784–793.
- JUMP, A. S. & PEÑUELAS, J. (2006). Genetic effects of chronic habitat fragmentation in a wind-pollinated tree. *Proceedings of the National Academy of Sciences of the United States of America* **103**, 8096–8100.
- *KARREBERG, S. & JENSEN, K. (2000). Effects of pollination and pollen source on the seed set of *Pedicularis palustris*. *Folia Geobotanica* **35**, 191–202.
- KELLER, L. F. & WALLER, D. M. (2002). Inbreeding effects in wild populations. *Trends in Ecology and Evolution* **17**, 230–241.
- *KELLY, D., LADLEY, J. J., ROBERTSON, A. W. & NORTON, D. A. (2000). Limited forest fragmentation improves reproduction in the declining New Zealand mistletoe *Peraxilla tetrapetala* (Loranthaceae). In *Genetics, Demography and Viability of Fragmented Populations* (eds A. G. YOUNG and G. CLARKE), pp. 241–252. Cambridge University Press, Cambridge.
- *KÉRY, M., MADSEN, J. & LEBRETON, J. D. (2006). Survival of Svalbard pink-footed geese *Anser brachyrhynchus* in relation to winter climate, density and land-use. *Journal of Animal Ecology* **75**, 1172–1181.
- KIESECKER, J. M., BLAUSTEIN, A. R. & BELDEN, L. K. (2001). Complex causes of amphibian population declines. *Nature* **410**, 681–684.
- KLOEPPEL, B. D., CLINTON, B. D., VOSE, J. M. & COOPER, A. R. (2003). Drought impacts on tree growth and mortality of Southern Appalachian Forests. In *Climate Variability and Ecosystem Response at Long-Term Ecological Research Sites* (eds D. GREENLAND, D. G. GOODIN and R. C. SMITH), Oxford University Press, Oxford.
- *KOLB, A. (2005). Reduced reproductive success and offspring survival in fragmented populations of the forest herb *Phyteuma spicatum*. *Journal of Ecology* **93**, 1226–1237.
- KUPARINEN, A., KATUL, G., NATHAN, R. & SCHURR, F. M. (2009). Increases in air temperature can promote wind-driven dispersal and spread of plants. *Proceedings of the Royal Society of London Series B: Biological Sciences* **276**, 3081–3087.
- LADA, H., THOMSON, J. R., CUNNINGHAM, S. C. & MAC NALLY, R. (2013). Rainfall in prior breeding seasons influences population size of a small marsupial. *Austral Ecology* **38**, 581–591.
- *LAHTI, K., ORELL, M., RYTKÖNEN, S. & KOIVULA, K. (1998). Time and food dependence in willow tit winter survival. *Ecology* **79**, 2904–2916.
- *LAMONT, B. B., KLINKHAMER, P. G. L. & WITKOWSKI, E. T. F. (1993). Population fragmentation may reduce fertility to zero in *Banksia goodii* – a demonstration of the Allee effect. *Oecologia* **94**, 446–450.
- LAMPILA, P., MÖNKÖNEN, M. & DESROCHERS, A. (2005). Demographic responses by birds to forest fragmentation. *Conservation Biology* **19**, 1537–1546.
- LANGTIMM, C. A. & BECK, C. A. (2003). Lower survival probabilities for adult Florida manatees in years with intense coastal storms. *Ecological Applications* **13**, 257–268.
- LAURANCE, W. F., FERREIRA, L. V., MERONA, J. M. R.-D., LAURANCE, S. G., HUTCHINGS, R. W. & LOVEJOY, T. E. (1998). Effects of forest fragmentation on recruitment patterns in Amazonian tree communities. *Conservation Biology* **12**, 460–464.
- LEHIKONEN, A., BYHOLM, P., RANTA, E., SAUROLA, P., VALKAMA, J., KORPIMÄKI, E., PIETIÄINEN, H. & HENTTONEN, H. (2009). Reproduction of the common buzzard at its northern range margin under climatic change. *Oikos* **118**, 829–836.
- LEHIKONEN, A., KILPI, M. & ÖST, M. (2006). Winter climate affects subsequent breeding success of common eiders. *Global Change Biology* **12**, 1355–1365.
- *LEIMU, R. & SYRJÄNEN, K. (2002). Effects of population size, seed predation and plant size on male and female reproductive success in *Vincetoxicum hirsutaria* (Asclepiadaceae). *Oikos* **98**, 229–238.
- *LENNARTSSON, T. (2002). Extinction thresholds and disrupted plant-pollinator interactions in fragmented plant populations. *Ecology* **83**, 3060–3072.
- LI, F. R., ZHAO, L. Y., ZHANG, H., LIU, J. L., LU, H. Y. & KANG, L. F. (2009). Habitat degradation, topography and rainfall variability interact to determine seed distribution and recruitment in a sand dune grassland. *Journal of Vegetation Science* **20**, 847–859.
- *LIENERT, J. & FISCHER, M. (2003). Habitat fragmentation affects the common wetland specialist *Primula farinosa* in north-east Switzerland. *Journal of Ecology* **91**, 587–599.
- LIMA, M., JULLIARD, R., STENSETH, N. C. & JAKSIC, F. M. (2001). Demographic dynamics of a neotropical small rodent (*Phyllotis darwini*): feedback structure, predation and climatic factors. *Journal of Animal Ecology* **70**, 761–775.
- LIN, Y. T. K. & BATZLI, G. O. (2001). The influence of habitat quality on dispersal, demography, and population dynamics of voles. *Ecological Monographs* **71**, 245–275.
- LINDÉN, A., LEHIKONEN, A., HOKKANEN, T. & VÄISÄNEN, R. A. (2011). Modelling irruptions and population dynamics of the great spotted woodpecker – joint effects of density and cone crops. *Oikos* **120**, 1065–1075.
- LINDSTRÖM, E. R. & HÖRNELDT, B. (1994). Vole cycles, snow depth and fox predation. *Oikos* **70**, 156–160.
- *LLOYD, P., MARTIN, T. E., REDMOND, R. L., LANGNER, U. & HART, M. M. (2005). Linking demographic effects of habitat fragmentation across landscapes to continental source-sink dynamics. *Ecological Applications* **15**, 1504–1514.
- LOWE, W. H. & ALLENDORF, F. W. (2010). What can genetics tell us about population connectivity? *Molecular Ecology* **19**, 3038–3051.
- *LUIJTEN, S. H., DIERICK, A., OOSTERMEIJER, J. G. B., RAIJMAN, L. E. & DEN NIJS, H. C. M. (2000). Population size, genetic variation, and reproductive success in a rapidly declining, self-incompatible perennial (*Arnica montana*) in the Netherlands. *Conservation Biology* **14**, 1776–1787.
- MAC NALLY, R., BENNETT, A. F. & HORROCKS, G. (2000). Forecasting the impacts of habitat fragmentation. Evaluation of species-specific predictions of the impact of habitat fragmentation on birds in the box-ironbark forests of central Victoria, Australia. *Biological Conservation* **95**, 7–29.
- MAC NALLY, R., BENNETT, A. F., THOMSON, J. R., RADFORD, J. Q., UNMACK, G., HORROCKS, G. & VESK, P. A. (2009). Collapse of an avifauna: climate change appears to exacerbate habitat loss and degradation. *Diversity and Distributions* **15**, 720–730.
- MAC NALLY, R., BOWEN, M., HOWES, A., MCALPINE, C. A. & MARON, M. (2012). Despotic, high-impact species and subcontinental scale control of avian assemblage structure. *Ecology* **93**, 668–678.
- MAC NALLY, R., CUNNINGHAM, S. C., BAKER, P. J., HORNER, G. J. & THOMSON, J. R. (2011). Dynamics of Murray-Darling floodplain forests under multiple stressors: the past, present, and future of an Australian icon. *Water Resources Research* **47**(12), 1–11.
- MACE, G., MASUNDIRE, H. & BAILLIE, J. (2005). Biodiversity. In *Ecosystems and Human Well-Being: Current State and Trends: Findings of the Condition and Trends Working Group. The Millennium Ecosystem Assessment Series*, Chapter 4 (eds R. HASSAN, R. J. SCHOLES and N. ASH). Island Press, Washington, DC.
- MANTYKA-PRINGLE, C. S., MARTIN, T. G. & RHODES, J. R. (2012). Interactions between climate and habitat loss effects on biodiversity: a systematic review and meta-analysis. *Global Change Biology* **18**, 1239–1252.
- MARON, M., GREY, M. J., CATTERALL, C. P., MAJOR, R. E., OLIVER, D. L., CLARKE, M. F., LOYN, R. H., MAC NALLY, R., DAVIDSON, I. & THOMSON, J. R. (2013). Avifaunal array due to a single despotic species. *Diversity and Distributions* **19**, 1468–1479.
- MARTINEZ-MEYER, E. (2012). Advances, limitation, and synergies in predicting changes in species' distribution and abundance under contemporary climate change. In *Ecological Consequences of Climate Change: Mechanisms, Conservation and Management* (eds E. A. BEEVER and J. L. BELANT), pp. 67–84. CRC Press, Boca Raton.
- MASSOT, M., CLOBERT, J. & FERRIERE, R. (2008). Climate warming, dispersal inhibition and extinction risk. *Global Change Biology* **14**, 461–469.
- MATTHYSEN, E. (1999). Nuthatches (*Sitta europaea*: Aves) in forest fragments: demography of a patchy population. *Oecologia* **119**, 501–509.
- *MAVRAGANIS, K. & ECKERT, C. G. (2001). Effect of population size and isolation on reproductive output in *Aquilegia canadensis* (Ranunculaceae). *Oikos* **95**, 300–310.
- MBORA, D. N. M., WIECZKOWSKI, J. & MUNENE, E. (2009). Links between habitat degradation, and social group size, ranging, fecundity, and parasite prevalence in the Tana River mangabey (*Cercocebus galensis*). *American Journal of Physical Anthropology* **140**, 562–571.
- MCÉUEN, A. B. & CURRAN, L. M. (2004). Seed dispersal and recruitment limitation across spatial scales in temperate forest fragments. *Ecology* **85**, 507–518.
- MCGARIGAL, K. & CUSHMAN, S. A. (2002). Comparative evaluation of experimental approaches to the study of habitat fragmentation effects. *Ecological Applications* **12**, 335–345.
- MCINTYRE, C. L. & SCHMIDT, J. H. (2012). Ecological and environmental correlates of territory occupancy and breeding performance of migratory Golden Eagles *Aquila chrysaetos* in interior Alaska. *Ibis* **154**, 124–135.
- MCKECHNIE, A. E. & WOLF, B. O. (2010). Climate change increases the likelihood of catastrophic avian mortality events during extreme heat waves. *Biology Letters* **6**, 253–256.
- MCLAUGHLIN, J. F., HELLMANN, J. J., BOGGS, C. L. & EHRLICH, P. R. (2002). Climate change hastens population extinctions. *Proceedings of the National Academy of Sciences of the United States of America* **99**, 6070–6074.
- MEMMOTT, J., CRAZE, P. G., WASER, N. M. & PRICE, M. V. (2007). Global warming and the disruption of plant–pollinator interactions. *Ecology Letters* **10**, 710–717.
- MENÉNDEZ, R., GONZÁLEZ-MEGÍAS, A., LEWIS, O. T., SHAW, M. R. & THOMAS, C. D. (2008). Escape from natural enemies during climate-driven range expansion: a case study. *Ecological Entomology* **33**, 413–421.
- MERCKX, T., KARLSSON, B. & VAN DYCK, H. (2006). Sex- and landscape-related differences in flight ability under suboptimal temperatures in a woodland butterfly. *Functional Ecology* **20**, 436–441.
- MILBAU, A., GRAAE, B. J., SHEVTSOVA, A. & NIJS, I. (2009). Effects of a warmer climate on seed germination in the subarctic. *Annals of Botany* **104**, 287–296.
- MILLER-RUSHING, A. J., HØYE, T. T., INOUE, D. W. & POST, E. (2010). The effects of phenological mismatches on demography. *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences* **365**, 3177–3186.
- *MØLLER, A. P. (1988). Nest predation and nest site choice in passerine birds in habitat patches of different size: a study of magpies and blackbirds. *Oikos* **53**, 215–221.
- *MOODY-WEIS, J. M. & HEYWOOD, J. S. (2001). Pollination limitation to reproductive success in the Missouri evening primrose *Oenothera macrocarpa* (Onagraceae). *American Journal of Botany* **88**, 1615–1622.

- MORA, C. & ZAPATA, F. A. (2013). Anthropogenic footprints on biodiversity. In *The Balance of Nature and Human Impact* (ed. K. ROHDE), pp. 239–258. Cambridge University Press, Cambridge.
- *MORGAN, J. W. (1999). Effects of population size on seed production and germinability in an endangered, fragmented grassland plant. *Conservation Biology* **13**, 266–273.
- *MORRISON, S. F. & HIK, D. S. (2007). Demographic analysis of a declining pika *Ochotona collaris* population: Linking survival to broad-scale climate patterns via spring snowmelt patterns. *Journal of Animal Ecology* **76**, 899–907.
- MORRISON, K. W., HIPFNER, J. M., BLACKBURN, G. S. & GREEN, D. J. (2011). Effects of extreme climate events on adult survival of three pacific auks. *Auk* **128**, 707–715.
- *MORSE, S. F. & ROBINSON, S. K. (1999). Nesting success of a neotropical migrant in a multiple-use, forested landscape. *Conservation Biology* **13**, 327–337.
- MOSES, M. R., FREY, J. K. & ROEMER, G. W. (2012). Elevated surface temperature depresses survival of banner-tailed kangaroo rats: will climate change cook a desert icon? *Oecologia* **168**, 257–268.
- MOSS, R., OSWALD, J. & BAINES, D. (2001). Climate change and breeding success: decline of the capercaillie in Scotland. *Journal of Animal Ecology* **70**, 47–61.
- MUNIER, A., HERMANUTZ, L., JACOBS, J. D. & LEWIS, K. (2010). The interacting effects of temperature, ground disturbance, and herbivory on seedling establishment: implications for treeline advance with climate warming. *Plant Ecology* **210**, 19–30.
- *MUSTAJARVI, K., SIKAMAKI, P., RYTKÖNEN, S. & LAMMI, A. (2001). Consequences of plant population size and density for plant–pollinator interactions and plant performance. *Journal of Ecology* **89**, 80–87.
- NAKAGAWA, S. & CUTHILL, I. C. (2007). Effect size, confidence interval and statistical significance: a practical guide for biologists. *Biological Reviews* **82**, 591–605.
- NEAL, J. M., HARDNER, C. M. & GROSS, C. L. (2010). Population demography and fecundity do not decline with habitat fragmentation in the rainforest tree *Macadamia integrifolia* (Proteaceae). *Biological Conservation* **143**, 2591–2600.
- NEVEU, A. (2009). Incidence of climate on common frog breeding: long-term and short-term changes. *Acta Oecologica* **35**, 671–678.
- NEWTON, I. (2008a). Mass mortality of migrants. In *Migration Ecology of Birds* (ed. I. NEWTON). Academic Press, London.
- NEWTON, I. (2008b). The Palaearctic-Afrotropical migration system. In *Migration Ecology of Birds* (ed. I. NEWTON). Academic Press, London.
- NIELSEN, J. T. & MÖLLER, A. P. (2006). Effects of food abundance, density and climate change on reproduction in the sparrowhawk *Accipiter nisus*. *Oecologia* **149**, 505–518.
- *OLSSON, M., SCHWARTZ, T., WAPSTRA, E., ULLER, T., UJVARI, B., MADSEN, T. & SHINE, R. (2011). Climate change, multiple paternity and offspring survival in lizards. *Evolution* **65**, 3323–3326.
- ONTIVEROS, D. & PLEGUEZUELOS, J. M. (2003). Influence of climate on Bonelli's eagle's (*Hieraetus fasciatus* V. 1822) breeding success through the Western Mediterranean. *Journal of Biogeography* **30**, 755–760.
- *OOSTERMEIJER, J. G. B., LUIJTEN, S. H., KRENOVA, Z. V. & DENNIJS, H. C. M. (1998). Relationships between population and habitat characteristics and reproduction of the rare *Gentiana pneumonanthe* L. *Conservation Biology* **12**, 1042–1053.
- OPDAM, P. & WASCHER, D. (2004). Climate change meets habitat fragmentation: linking landscape and biogeographical scale levels in research and conservation. *Biological Conservation* **117**, 285–297.
- PANKHURST, N. W. & MUNDAY, P. L. (2011). Effects of climate change on fish reproduction and early life history stages. *Marine and Freshwater Research* **62**, 1015–1026.
- PARMESAN, C. (2005). Detection at multiple levels: *Euphydryas editha* and climate change. In *Climate Change and Biodiversity* (eds T. E. LOVEJOY and L. HANNAH), pp. 56–60. Yale University Press, New Haven, CT and London.
- PARMESAN, C. (2006). Ecological and evolutionary responses to recent climate Change. *Annual Review of Ecology, Evolution, and Systematics* **37**, 637–669.
- PÄRN, H., RINGSBY, T. H., JENSEN, H. & SÆTHER, B. E. (2011). Spatial heterogeneity in the effects of climate and density-dependence on dispersal in a house sparrow metapopulation. *Proceedings of the Royal Society of London Series B: Biological Sciences* **279**, 144–152.
- *PARRA-TABLA, V. C., VARGAS, C. F., MAGAÑA-RUEDA, S. & NAVARRO, J. (2000). Female and male pollination success of *Oncidium ascendens* Lindey (Orchidaceae) in two contrasting habitat patches: forest vs agricultural field. *Biological Conservation* **94**, 335–340.
- *PASCHKE, M., ABS, C. & SCHMID, B. (2002). Effects of population size and pollen diversity on reproductive success and offspring size in the narrow endemic *Cochlearia bavarica* (Brassicaceae). *American Journal of Botany* **89**, 1250–1259.
- PAVLOVA, A., AMOS, J. N., GORETSKAIA, M. I., BEME, I. R., BUCHANAN, K. L., TAKEUCHI, N., RADFORD, J. Q. & SUNNUCKS, P. (2012). Genes and song: genetic and social connections in fragmented habitat in a woodland bird with limited dispersal. *Ecology* **93**, 1717–1727.
- PECHMANN, J. H. K., SCOTT, D. E., SEMLITSCH, R. D., CALDWELL, J. P., VITT, L. J. & GIBBONS, J. W. (1991). Declining amphibian populations: the problem of separating human impacts from natural fluctuations. *Science* **253**, 892–895.
- VAN DE POL, M., ENS, B. J., HEG, D., BROUWER, L., KROL, J., MAIER, M., EXO, K.-M., OOSTERBEEK, K., LOK, T., EISING, C. M. & KOFFIJBERG, K. (2010). Do changes in the frequency, magnitude and timing of extreme climatic events threaten the population viability of coastal birds? *Journal of Applied Ecology* **47**, 720–730.
- *PORNELUZI, P. A. & FAABORG, J. (2001). Season-long fecundity, survival, and viability of ovenbirds in fragmented and unfragmented landscapes. *Conservation Biology* **13**, 1151–1161.
- POST, E. & FORCHHAMMER, M. C. (2008). Climate change reduces reproductive success of an Arctic herbivore through trophic mismatch. *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences* **363**, 2369–2375.
- *POST, E. & STENSETH, N. C. (1999). Climatic variability, plant phenology, and northern ungulates. *Ecology* **80**, 1322–1339.
- PREVITALI, M. A., LIMA, M., MESERVE, P. L., KELT, D. A. & GUTIÉRREZ, J. R. (2009). Population dynamics of two sympatric rodents in a variable environment: rainfall, resource availability, and predation. *Ecology* **90**, 1996–2006.
- PÜTTKER, T., BUENO, A. A., SANTOS, D., DE BARROS, C., SOMMER, S. & PARDINI, R. (2011). Immigration rates in fragmented landscapes: empirical evidence for the importance of habitat amount for species persistence. *PLoS ONE* **6**, e27963.
- *QUESADA, M., STONER, K. E., LOBO, J. A., HERRERIAS-DIEGO, Y., PALACIOS-GUEVERA, C., MUNGIA-ROSAS, M. A., SALAZAR, K. A. & ROSAS-GUERRERO, V. (2004). Effects of forest fragmentation on pollinator activity and consequences for plant reproductive success and mating patterns in bat pollinated bombacaceous trees. *Biotropica* **36**, 400–406.
- *QUESADA, M., STONER, K. E., ROSAS-GUERRERO, V., PALACIOS-GUEVERA, C. & LOBO, J. A. (2003). Effects of habitat disruption on the activity of nectarivorous bats (Chiroptera: Phyllostomidae) in a dry tropical forest: implications for the reproductive success of the neotropical tree *Cecropia grandifolia*. *Oecologia* **135**, 400–406.
- RAULINGS, E., MORRIS, K., THOMPSON, R. & MAC NALLY, R. (2011). Do birds of a feather disperse plants together? *Freshwater Biology* **56**, 1390–1402.
- *REDPATH, S. M. (1995). Habitat fragmentation and the individual: tawny owls *Strix aluco* in woodland patches. *Journal of Animal Ecology* **64**, 652–661.
- REED, D. H., O'GRADY, J. J., BROOK, B. W., BALLOU, J. D. & FRANKHAM, R. (2003). Estimates of minimum viable population sizes for vertebrates and factors influencing those estimates. *Biological Conservation* **113**, 23–34.
- REED, A. W. & SLADE, N. A. (2009). Environmental correlates of survival and reproduction in old-field rodents. *Journal of Mammalogy* **90**, 680–685.
- REICH, P. B. & OLEKSYN, J. (2008). Climate warming will reduce growth and survival of Scots pine except in the far north. *Ecology Letters* **11**, 588–597.
- *ROBINSON, S. K., FRANK, R. T. III, DONOVAN, T. M., WHITEHEAD, D. R. & FAABORG, J. (1995). Regional forest fragmentation and the nesting success of migratory birds. *Science* **267**, 1987–1990.
- ROBLES, H., CIUDAD, C., VERA, R., OLEA, P. P. & MATTYSEN, E. (2008). Demographic responses of middle spotted woodpeckers (*Dendrocopos medius*) to habitat fragmentation. *Auk* **125**, 131–139.
- *ROCHA, O. J. & AGUILAR, G. (2001). Reproductive biology of the dry forest tree *Enterolobium cyclocarpum* (Guanacaste) in Costa Rica: a comparison between trees left in pastures and trees in continuous forest. *American Journal of Botany* **88**, 1607–1614.
- *RODRÍGUEZ, C. & BUSTAMANTE, J. (2003). The effect of weather on lesser kestrel breeding success: can climate change explain historical population declines? *Journal of Animal Ecology* **72**, 793–810.
- ROSENTHAL, R. (1994). Parametric measures of effect size. In *The Handbook of Research Synthesis* (eds H. COOPER and L. V. HEDGES), pp. 231–244. Sage, New York.
- *ROSSETTO, M., GROSS, C. L., JONES, R. & HUNTER, J. (2004). The impact of clonality on an endangered tree (*Elaeocarpus williamsianus*) in a fragmented rainforest. *Biological Conservation* **117**, 33–39.
- ROTHERMEL, B. B. & SEMLITSCH, R. D. (2002). An experimental investigation of landscape resistance of forest versus old-field habitats to emigrating juvenile amphibians. *Conservation Biology* **16**, 1324–1332.
- SALA, O. E., CHAPIN, F. S. III, ARMESTO, J. J., BERLOW, E., BLOOMFIELD, J., DIRZO, R., HUBER-SANWALD, E., HUENNEKE, L. F., JACKSON, R. B., KINZIG, A., LEEMANS, R., LODGE, D. M., MOONEY, H. A., OESTERHELD, M. N., POFF, N. L., SYKES, M. T., WALKER, B. H., WALKER, M. & WALL, D. H. (2000). Global biodiversity scenarios for the year 2100. *Science* **287**, 1770–1774.
- SÆTHER, B.-E. & BAKKE, Ø. (2000). Avian life history variation and contribution of demographic traits to the population growth rate. *Ecology* **81**, 642–653.
- *SALEWSKI, V., HOCHACHKA, W. M. & FIEDLER, W. (2013). Multiple weather factors affect apparent survival of European passerine birds. *PLoS ONE* **8**, e59110.
- *SANDERS, N. J. & GORDON, D. M. (2004). The interactive effects of climate, life history, and interspecific neighbours on mortality in a population of seed harvester ants. *Ecological Entomology* **29**, 632–637.
- SANDERSON, F. J., DONALD, P. F., PAIN, D. J., BURFIELD, I. J. & VAN BOMMEL, F. P. J. (2006). Long-term population declines in Afro-Palaearctic migrant birds. *Biological Conservation* **131**, 93–105.
- SANDVIK, H., ERIKSTAD, K. E., BARRETT, R. T. & YOCOZ, N. G. (2005). The effect of climate on adult survival in five species of North Atlantic seabirds. *Journal of Animal Ecology* **74**, 817–831.
- SANZ, J. J. (2003). Large-scale effect of climate change on breeding parameters of pied flycatchers in western Europe. *Ecography* **26**, 45–50.

- SANZ, J. J., POTTI, J., MORENO, J., MERINO, S. & FRÍAS, O. (2003). Climate change and fitness components of a migratory bird breeding in the Mediterranean region. *Global Change Biology* **9**, 461–472.
- *SCHMIDT, K. & JENSEN, K. (2000). Genetic structure and AFLP variation of remnant populations in the rare plant *Pedicularis palustris* (Scrophulariaceae) and its relation to population size and reproductive components. *American Journal of Botany* **87**, 678–689.
- SCHTICKZELLE, N. & BAGUETTE, M. (2003). Behavioural responses to habitat patch boundaries restrict dispersal and generate emigration–patch area relationships in fragmented landscapes. *Journal of Animal Ecology* **72**, 533–545.
- SCHWANZ, L. E., SPENCER, R. J., BOWDEN, R. M. & JANZEN, F. J. (2010). Climate and predation dominate juvenile and adult recruitment in a turtle with temperature-dependent sex determination. *Ecology* **91**, 3016–3026.
- SENPATHI, D., NICOLL, M. A. C., TEPLITSKY, C., JONES, C. G. & NORRIS, K. (2011). Climate change and the risks associated with delayed breeding in a tropical wild bird population. *Proceedings of the Royal Society of London Series B: Biological Sciences* **278**, 3184–3190.
- *SEVERNS, P. (2003). Inbreeding and small population size reduce seed set in threatened and fragmented plant species, *Lupinus sulphureus* ssp. *kincaidii* (Fabaceae). *Biological Conservation* **110**, 221–229.
- SEVTSOVA, A., GRAAE, B. J., JOCHUM, T., MILBAU, A., KOCKELBERGH, F., BEYENS, L. & NIJS, I. (2009). Critical periods for impact of climate warming on early seedling establishment in subarctic tundra. *Global Change Biology* **15**, 2662–2680.
- SILLETT, T. S., HOLMES, R. T. & SHERRY, T. W. (2000). Impacts of a global climate cycle on population dynamics of a migratory songbird. *Science* **288**, 2040–2043.
- SINCLAIR, B. J. & CHOWN, S. L. (2003). Rapid responses to high temperature and desiccation but not to low temperature in the freeze tolerant sub-Antarctic caterpillar *Pringleophaga marioni* (Lepidoptera, Tineidae). *Journal of Insect Physiology* **49**, 45–52.
- SKAGEN, S. K. & ADAMS, A. A. Y. (2012). Weather effects on avian breeding performance and implications of climate change. *Ecological Applications* **22**, 1131–1145.
- SKOGLAND, T. (1985). The effects of density-dependent resource limitations on the demography of wild reindeer. *Journal of Animal Ecology* **54**, 359–374.
- *SMITH-RAMIREZ, C. & ARMESTO, J. J. (2003). Foraging behaviour of bird pollinators on *Embothrium coccineum* (Proteaceae) trees in forest fragments and pastures in southern Chile. *Austral Ecology* **28**, 53–60.
- *SOMANATHAN, H. & BORGES, R. M. (2000). Influence of exploitation on population structure, spatial distribution, and reproductive success of dioecious species in a fragmented cloud forest in India. *Biological Conservation* **94**, 243–256.
- SOMERO, G. N. (2010). The physiology of climate change: how potentials for acclimatization and genetic adaptation will determine 'winners' and 'losers'. *Journal of Experimental Biology* **213**, 912–920.
- SORTE, C. J. B. & WHITE, J. W. (2013). Competitive and demographic leverage points of community shifts under climate warming. *Proceedings of the Royal Society of London Series B: Biological Sciences* **280**, 20130572.
- STEENHOF, K., KOCHERT, M. N. & McDONALD, T. L. (1997). Interactive effects of prey and weather on Golden eagle reproduction. *Journal of Animal Ecology* **66**, 350–362.
- *STEFFAN-DEWENTER, I. & TSCHARNTKE, T. (1999). Effects of habitat isolation on pollinator communities and seed set. *Oecologia* **121**, 432–440.
- STOW, A. J. & SUNNUCKS, P. (2004). Inbreeding avoidance in Cunningham's skink (*Egernia cunninghami*) in natural and fragmented habitat. *Molecular Ecology* **13**, 443–447.
- SWIFT, T. L. & HANNON, S. J. (2010). Critical thresholds associated with habitat loss: a review of the concepts, evidence, and applications. *Biological Reviews* **85**, 35–53.
- TALLMON, D. A., JULES, E. S., RADKE, N. J. & MILLS, L. S. (2003). Of mice and men and trillium: cascading effects of forest fragmentation. *Ecological Applications* **13**, 1193–1203.
- *TEWKSURY, J. J., HEJL, S. J. & MARTIN, T. E. (2005). Breeding productivity does not decline with increasing fragmentation in a western landscape. *Ecology* **79**, 2890–2903.
- *TEWKSURY, J. J., LEVEY, D. J., HADDAD, N. M., SARGENT, S., ORROCK, J. L., WELDON, A., DANIELSON, B. J., BRINKERHOFF, J., DAMSCHEN, E. I. & TOWNSEND, P. (2002). Corridors affect plants, animals, and their interactions in fragmented landscapes. *Proceedings of the National Academy of Sciences of the United States of America* **99**, 12923–12926.
- THOMAS, C. D., FRANCO, A. M. A. & HILL, J. K. (2006). Range retractions and extinction in the face of climate warming. *Trends in Ecology and Evolution* **21**, 415–416.
- TIAN, H., CHEN, G., LIU, M., ZHANG, C., SUN, G., LU, C., XU, X., REN, W., PAN, S. & CHAPPELKA, A. (2010). Model estimates of net primary productivity, evapotranspiration, and water use efficiency in the terrestrial ecosystems of the southern United States during 1895–2007. *Forest Ecology and Management* **259**, 1311–1327.
- TOMILLO, P. S., SABA, V. S., BLANCO, G. S., STOCK, C. A., PALADINO, F. V. & SPOTILA, J. R. (2012). Climate driven egg and hatchling mortality threatens survival of Eastern Pacific leatherback turtles. *PLoS ONE* **7**, e37602.
- *TOMIMATSU, H. & OHARA, M. (2002). Effects of forest fragmentation on seed production of the understory herb *Trillium camschatcense*. *Conservation Biology* **16**, 1277–1285.
- TORÄNG, P., EHRLÉN, J. & ÅGREN, J. (2010). Linking environmental and demographic data to predict future population viability of a perennial herb. *Oecologia* **163**, 99–109.
- TYLIANAKIS, J. M., DIDHAM, R. K., BASCOMPTE, J. & WARDLE, D. A. (2008). Global change and species interactions in terrestrial ecosystems. *Ecology Letters* **11**, 1351–1363.
- *UMAPATHY, G., HUSSAIN, S. & SHIVAJI, S. (2011). Impact of habitat fragmentation on the demography of lion-tailed macaque (*Macaca silenus*) populations in the rainforests of Anamalai Hills, Western Ghats, India. *International Journal of Primatology* **32**, 889–900.
- *UMAPATHY, G. & KUMAR, A. (2000). The Demography of the lion-tailed macaque (*Macaca silenus*) in rain forest fragments in the Anamalai Hills, South India. *Primates* **41**, 119–126.
- URIARTE, M., BRUNA, E. M., RUBIM, P., ANCIÃES, M. & JONCKHEERE, I. (2010). Effects of forest fragmentation on the seedling recruitment of a tropical herb: assessing seed vs. safe-site limitation. *Ecology* **91**, 1317–1328.
- *VALDIVIA, C. E., SIMONETTI, J. A. & HENRIQUEZ, C. A. (2006). Depressed pollination of *Lapageria rosea* Ruiz et Pav. (Philesiaceae) in the fragmented temperate rainforest of southern South America. *Biodiversity and Conservation* **15**, 1845–1856.
- *VALIMAKI, P. & ITAMIES, J. (2003). Migration of the clouded Apollo butterfly *Parnassius mnemosyne* in a network of suitable habitats: effects of patch characteristics. *Ecography* **26**, 679–691.
- VAN MANTGEM, P. J. & STEPHENSON, N. L. (2007). Apparent climatically induced increase of tree mortality rates in a temperate forest. *Ecology Letters* **10**, 909–916.
- *VAN ROSSUM, F., ECHCHAGGADA, G., SZABADI, I. & TRIEST, L. (2002). Commonness and long-term survival in fragmented habitats: *Primula elatior* as a study case. *Conservation Biology* **16**, 1286–1295.
- VATKA, E., ORELL, M. & RYTKÖNEN, S. (2011). Warming climate advances breeding and improves synchrony of food demand and food availability in a boreal passerine. *Global Change Biology* **17**, 3002–3009.
- VIÉ, J.-C., HILTON-TAYLOR, C. & STUART, S. N. (2009). *Wildlife in a Changing World: An Analysis of the 2008 IUCN Red List of Threatened Species*. IUCN, Gland.
- *VILLARD, M. A., MARTIN, P. R. & DRUMMOND, C. G. (1993). Habitat fragmentation and pairing success in the ovenbird (*Seiurus aurocapillus*). *Auk* **110**, 759–769.
- VOS, C. C., ANTONISSE-DE JONG, A. G., GOEDHART, P. W. & SMULDERS, M. J. M. (2001). Genetic similarity as a measure for connectivity between fragmented populations of the moor frog (*Rana arvalis*). *Heredity* **86**, 598–608.
- WAGENIUS, S., LONSDORF, E. & NEUHAUSER, C. (2007). Patch aging and the S-Allee effect: breeding system effects on the demographic response of plants to habitat fragmentation. *American Naturalist* **169**, 383–397.
- WAITE, T. A. & STRICKLAND, D. (2006). Climate change and the demographic demise of a hoarding bird living on the edge. *Proceedings of the Royal Society of London Series B: Biological Sciences* **273**, 2809–2813.
- WALTHER, G.-R., POST, E., CONVEY, P., MENZEL, A., PARMESAN, C., BEEBEE, T. J. C., FROMENTIN, J.-M., HOEGH-GULDBERG, O. & BAIRLEIN, F. (2002). Ecological responses to recent climate change. *Nature* **416**, 389–395.
- *WANG, H., SORK, V. L., WU, J. & GE, J. (2010). Effect of patch size and isolation on mating patterns and seed production in an urban population of Chinese pine (*Pinus tabulaeformis* Carr.). *Forest Ecology and Management* **260**, 965–974.
- *WARBURTON, C. L., JAMES, E. A., FRIPP, Y. J., TRUEMAN, S. J. & WALLACE, H. M. (2000). Clonality and sexual reproductive failure in remnant populations of *Santalum lanceolatum* (Santalaceae). *Conservation Biology* **96**, 45–54.
- *WARD, M. & JOHNSON, S. D. (2005). Pollen limitation and demographic structure in small fragmented populations of *Brunsvigia radulosa* (Amaryllidaceae). *Oikos* **108**, 253–262.
- WARREN, M., HILL, J., THOMAS, J., ASHER, J., FOX, R., HUNTLEY, B., ROY, D., TELFER, M., JEFFCOATE, S. & HARDING, P. (2001). Rapid responses of British butterflies to opposing forces of climate and habitat change. *Nature* **414**, 65–69.
- *WATANABE, A., GOKA, K. & WASHITANI, I. (2003). Effects of population spatial structure on the quantity and quality of seeds set by *Primula sieboldii* (Primulaceae). *Plant Species Biology* **18**, 107–121.
- WATERS, L. A., HUTCHINSON, Y., PARKIN, D. T. & DHONDT, A. A. (1994). The effects of habitat fragmentation on demography and on the loss of genetic variation in the red squirrel. *Proceedings of the Royal Society of London Series B: Biological Sciences* **255**, 107–111.
- WERNER, F. A. (2011). Reduced growth and survival of vascular epiphytes on isolated remnant trees in a recent tropical montane forest clear-cut. *Basic and Applied Ecology* **12**, 172–181.
- WERNER, F. & GRADSTEIN, S. (2008). Seedling establishment of vascular epiphytes on isolated and enclosed forest trees in an Andean landscape, Ecuador. *Biodiversity and Conservation* **17**, 3195–3207.
- *WESTLIN, L. M. (2000). Do pouched mice alter litter size through resorption in response to resource availability? *South African Journal of Wildlife Research* **30**, 118–121.
- WIGLEY, B. J., BOND, W. J. & HOFFMAN, M. T. (2010). Thicket expansion in a South African savanna under divergent land use: local vs. global drivers? *Global Change Biology* **16**, 964–976.
- WILCOCK, C. & NEILAND, R. (2002). Pollination failure in plants: why it happens and when it matters. *Trends in Plant Science* **7**, 270–277.

- WILEY, J. W. & WUNDERLE, J. M. JR. (1993). The effects of hurricanes on birds, with special reference to Caribbean islands. *Bird Conservation International* **3**, 319–349.
- WILLI, Y., BUSKIRK, J. V. & HOFFMANN, A. A. (2006). Limits to the adaptive potential of small populations. *Annual Review of Ecology, Evolution, and Systematics* **37**, 433–458.
- *WOLF, A. T. & HARRISON, S. P. (2001). Effects of habitat size and patch isolation on reproductive success of the serpentine morning glory. *Conservation Biology* **15**, 111–121.
- WOLF, S. G., SYDEMAN, W. J., HIPFNER, J. M., ABRAHAM, C. L., TERSHY, B. R. & CROLL, D. A. (2009). Range-wide reproductive consequences of ocean climate variability for the seabird Cassin's Auklet. *Ecology* **90**, 742–753.
- *YATES, C. J. & LADD, P. G. (2005). Effects of population spatial structure on the quantity and quality of seeds set by *Primula sieboldii* (Primulaceae). *Conservation Biology* **19**, 239–249.
- YOUNG, A. G., BROADHURST, L. M. & THRALL, P. H. (2012). Non-additive effects of pollen limitation and self-incompatibility reduce plant reproductive success and population viability. *Annals of Botany* **109**, 643–653.
- YOUNG, A. G., BROWN, A. D. H., MURRAY, B. G., THRALL, P. H. & MILLER, C. H. (2000). Genetic erosion, restricted mating and reduced viability in fragmented populations of the endangered grassland herb *Rutidosia leptorrhynchoidea*. In *Genetics, Demography and Viability of Fragmented Populations* (eds A. G. YOUNG and G. M. CLARKE), pp. 335–360. Cambridge University Press, Cambridge.
- ZANETTE, L., DOYLE, P. & TREMONT, S. M. (2000). Food shortage in small fragments: evidence from an area-sensitive passerine. *Ecology* **81**, 1654–1666.
- ZANI, P. A. (2008). Climate change trade-offs in the side-blotched lizard (*Uta stansburiana*): effects of growing-season length and mild temperatures on winter survival. *Physiological and Biochemical Zoology* **81**, 797–809.
- ZITSKE, B. P., BETTS, M. G. & DIAMOND, A. W. (2011). Negative effects of habitat loss on survival of migrant warblers in a forest mosaic. *Conservation Biology* **25**, 993–1001.

IX. SUPPORTING INFORMATION

Additional supporting information may be found in the online version of this article.

Table S1. Search terms used to locate studies that measured the effect of climate and landscape condition on demographic rates.

Table S2. Search terms used to locate studies that measured the effect of climate and landscape condition on variables related to demographic rates.

Table S3. List of species used for the calculation of mean effect sizes for climate and landscape condition on population vital rates.

Table S4. Breakdown of individual studies (December 2012 and earlier) that measured demographic responses to landscape condition and climate by region and taxonomic group.

Table S5. Subset of studies (from online Table S3) that showed effects of temperature and rainfall variables on birth or death rates.

(Received 4 December 2013; revised 8 July 2014; accepted 14 July 2014; published online 25 August 2014)