

Fitness Analytics Report

Fat loss · 75 day tracking period

Cohort Profile

AGE

23 years

SEX

MALE

HEIGHT

174 cm

STARTING WEIGHT

76 kg

GOAL

FAT_LOSS

PARTICIPANTS

75

Overall Outcome

18 / 75

Participants meeting the primary outcome

57 participants did not meet the primary outcome.

Metric Performance

Ending Weight

How each participant's ending body weight compared to the target range for their goal.

PASSED

18

FAILED ABOVE IDEAL

57

FAILED BELOW IDEAL

0

Participants lost weight too slowly, resulting in minimal or insignificant fat loss over the tracked period.

Daily Calorie Target

Average daily calorie intake compared to the target range for the stated goal.

PASSED

8

FAILED ABOVE IDEAL

67

FAILED BELOW IDEAL

0

Average intake exceeded the target range, undermining the intended deficit or overshooting the intended surplus.

Daily Protein Goal

Average daily protein intake compared to the recommended range.

PASSED	FAILED ABOVE IDEAL	FAILED BELOW IDEAL
75	0	0

Daily Fat Intake

Average daily fat intake compared to the recommended range.

PASSED	FAILED ABOVE IDEAL	FAILED BELOW IDEAL
13	62	0

Fat intake exceeded the recommended range, contributing excess calories beyond the target.

Weekly Cardio Minutes

Average weekly cardio minutes compared to the recommended range.

PASSED	FAILED ABOVE IDEAL	FAILED BELOW IDEAL
75	0	0

Daily Steps

Average daily step count compared to the recommended range.

PASSED	FAILED ABOVE IDEAL	FAILED BELOW IDEAL
75	0	0

Upper Body Training Frequency

Upper body sessions per week compared to the recommended range.

PASSED	FAILED ABOVE IDEAL	FAILED BELOW IDEAL
0	75	0

Upper body frequency exceeded the recommended range, raising the risk of inadequate recovery between sessions.

Lower Body Training Frequency

Lower body sessions per week compared to the recommended range.

PASSED	FAILED ABOVE IDEAL	FAILED BELOW IDEAL
28	0	47

Lower body frequency fell short of the recommended range, limiting stimulus for growth and strength adaptation.

Total Training Frequency

Total training sessions per week compared to the recommended range.

PASSED	FAILED ABOVE IDEAL	FAILED BELOW IDEAL
61	14	0

Total training frequency exceeded the recommended range, raising the risk of overtraining or inadequate recovery.

Training Intensity (RIR)

Average reps-in-reserve compared to the recommended range for effective training.

PASSED	FAILED ABOVE IDEAL	FAILED BELOW IDEAL
75	0	0

Average Sleep

Average nightly sleep compared to the recommended range.

PASSED	FAILED ABOVE IDEAL	FAILED BELOW IDEAL
75	0	0

Analytical Findings

Ending weight results showed a clear split across the cohort, with 18 participants successfully landing within the ideal target range of 67.86 kg to 71.93 kg to support their fat loss goal. However, 57 participants finished above this ideal range with an average deviation of 3.15 kg, indicating that the majority of the cohort lost weight too slowly to achieve significant fat loss over the 75-day tracking period.

Daily calorie intake exceeded the ideal target range of 1628.48 to 2029.69 kcal for 67 participants, who averaged an intake above the maximum threshold by 365.91 kcal. Only 8 participants successfully maintained their average intake at 1971.21 kcal within the recommended range, directly contributing to the difficulties many faced in reaching their ideal ending weight.

Daily fat intake averaged 55.44 g across the cohort, with 62 participants exceeding the ideal range of 43.77 to 58.36 g by an average of 8.70 g. Meanwhile, 13 participants successfully kept their average fat consumption within the prescribed limits, aligning better with the specific nutritional constraints required for the cohort's fat loss objective.

Daily protein intake averaged 141.89 g, successfully meeting the ideal target range of 131.30 to 160.48 g across all 75 participants. This consistent protein consumption effectively supported muscle repair and retention while participants worked toward their body composition goals.

Weekly cardio volume averaged 206.65 minutes, successfully falling entirely within the ideal target range of 150.0 to 300.0 minutes for all 75 participants. This volume met general cardiovascular health guidelines and provided a reliable baseline of activity to support the cohort's energy expenditure objectives.

Daily step counts averaged 10,502.60 steps, successfully placing all 75 participants within the ideal range of 7,000.0 to 15,000.0 steps. This consistent daily movement effectively supported general activity levels and overall calorie expenditure requirements.

Upper body training frequency exceeded the ideal range of 2.0 to 3.0 sessions per week for all 75 participants, raising the risk of inadequate recovery between sessions. This elevated frequency deviated above the maximum target by an average of 1.24 sessions per week, potentially impacting the cohort's overall recovery capacity.

Lower body training frequency fell below the ideal range of 2.0 to 3.0 sessions per week for 47 participants, who experienced a deviation below the minimum target by an average of 1.0 session. Conversely, 28 participants successfully maintained a lower body frequency of 2.0 sessions per week, supporting adequate muscle stimulus for their goals.

Total training frequency averaged 5.59 sessions per week, with 61 participants successfully landing within the ideal range of 4.0 to 6.0 sessions. However, 14 participants exceeded this ideal upper limit by an average of 0.14 sessions per week, slightly increasing their risk of overtraining relative to the rest of the cohort.

Training intensity, measured by reps-in-reserve, averaged 2.65 across all 75 participants, successfully staying

within the ideal target range of 0.0 to 3.0. This intensity ensured that training stayed close enough to failure to provide an effective stimulus while participants pursued their physical development goals.

Average sleep duration was 7.24 hours, successfully placing all 75 participants within the ideal target range of 7.0 to 9.0 hours. This consistent rest duration effectively supported recovery and hormone regulation throughout the 75-day tracking period.