

Function Operations

Objectives

- 1 Perform arithmetic operations to functions
- 2 Find the domain of the sum, difference, product, or quotient of two functions
- 3 Find the difference quotient of a function

Notation

We can add, subtract, multiply, and divide functions much like we can with real numbers:

Notation

We can add, subtract, multiply, and divide functions much like we can with real numbers:

- $(f + g)(x) = f(x) + g(x)$

Notation

We can add, subtract, multiply, and divide functions much like we can with real numbers:

- $(f + g)(x) = f(x) + g(x)$
 - add corresponding y -coordinates

Notation

We can add, subtract, multiply, and divide functions much like we can with real numbers:

- $(f + g)(x) = f(x) + g(x)$
 - add corresponding y -coordinates
- $(f - g)(x) = f(x) - g(x)$

Notation

We can add, subtract, multiply, and divide functions much like we can with real numbers:

- $(f + g)(x) = f(x) + g(x)$
 - add corresponding y -coordinates
- $(f - g)(x) = f(x) - g(x)$
 - subtract corresponding y -coordinates

Notation

- $(fg)(x) = f(x) \cdot g(x)$

Notation

- $(fg)(x) = f(x) \cdot g(x)$
 - multiply corresponding y -coordinates

Notation

- $(fg)(x) = f(x) \cdot g(x)$
 - multiply corresponding y -coordinates
- $\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)}, \quad g(x) \neq 0$

Notation

- $(fg)(x) = f(x) \cdot g(x)$
 - multiply corresponding y -coordinates
- $\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)}, \quad g(x) \neq 0$
 - divide corresponding y -coordinates

Example 1

For $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(a) Simplify $(f + g)(x)$

Example 1

For $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(a) Simplify $(f + g)(x)$

$$(f + g)(x) = f(x) + g(x)$$

Example 1

For $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(a) Simplify $(f + g)(x)$

$$(f + g)(x) = f(x) + g(x)$$

$$= 6x^2 - 2x + 3 - \frac{1}{x}$$

Example 1 $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(b) Evaluate $(f + g)(-1)$

Example 1 $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(b) Evaluate $(f + g)(-1)$

$$(f + g)(-1) = f(-1) + g(-1)$$

Example 1 $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(b) Evaluate $(f + g)(-1)$

$$(f + g)(-1) = f(-1) + g(-1)$$

$$= 8 + 4$$

Example 1 $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(b) Evaluate $(f + g)(-1)$

$$(f + g)(-1) = f(-1) + g(-1)$$

$$= 8 + 4$$

$$= 12$$

Example 1 $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(b) Evaluate $(f + g)(-1)$

$$(f + g)(-1) = f(-1) + g(-1)$$

$$= 8 + 4$$

$$= 12$$

$$(f + g)(-1) = 6(-1)^2 - 2(-1) + 3 - \left(\frac{1}{-1}\right)$$

Example 1 $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(b) Evaluate $(f + g)(-1)$

$$(f + g)(-1) = f(-1) + g(-1)$$

$$= 8 + 4$$

$$= 12$$

$$(f + g)(-1) = 6(-1)^2 - 2(-1) + 3 - \left(\frac{1}{-1}\right)$$

$$= 12$$

Example 1 $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(c) Simplify $(f - g)(x)$

Example 1 $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(c) Simplify $(f - g)(x)$

$$(f - g)(x) = f(x) - g(x)$$

Example 1 $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(c) Simplify $(f - g)(x)$

$$(f - g)(x) = f(x) - g(x)$$

$$= (6x^2 - 2x) - \left(3 - \frac{1}{x}\right)$$

Example 1 $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(c) Simplify $(f - g)(x)$

$$(f - g)(x) = f(x) - g(x)$$

$$= (6x^2 - 2x) - \left(3 - \frac{1}{x}\right)$$

$$= 6x^2 - 2x - 3 + \frac{1}{x}$$

Example 1 $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(d) Simplify $(g - f)(x)$

Example 1 $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(d) Simplify $(g - f)(x)$

$$(g - f)(x) = g(x) - f(x)$$

Example 1 $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(d) Simplify $(g - f)(x)$

$$(g - f)(x) = g(x) - f(x)$$

$$= 3 - \frac{1}{x} - (6x^2 - 2x)$$

Example 1 $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(d) Simplify $(g - f)(x)$

$$(g - f)(x) = g(x) - f(x)$$

$$= 3 - \frac{1}{x} - (6x^2 - 2x)$$

$$= 3 - \frac{1}{x} - 6x^2 + 2x$$

Example 1 $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(e) Simplify $(fg)(x)$

Example 1 $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(e) Simplify $(fg)(x)$

$$(fg)(x) = f(x) \cdot g(x)$$

Example 1 $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(e) Simplify $(fg)(x)$

$$(fg)(x) = f(x) \cdot g(x)$$

$$= (6x^2 - 2x) \left(3 - \frac{1}{x} \right)$$

Example 1 $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(e) Simplify $(fg)(x)$

$$(fg)(x) = f(x) \cdot g(x)$$

$$= (6x^2 - 2x) \left(3 - \frac{1}{x} \right)$$

$$= 18x^2 - 6x - 6x + 2$$

Example 1 $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(e) Simplify $(fg)(x)$

$$(fg)(x) = f(x) \cdot g(x)$$

$$= (6x^2 - 2x) \left(3 - \frac{1}{x} \right)$$

$$= 18x^2 - 6x - 6x + 2$$

$$= 18x^2 - 12x + 2$$

Example 1 $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(f) Simplify $\left(\frac{f}{g}\right)(x)$

Example 1 $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(f) Simplify $\left(\frac{f}{g}\right)(x)$

$$\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)}$$

Example 1 $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(f) Simplify $\left(\frac{f}{g}\right)(x)$

$$\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)}$$

$$= \frac{6x^2 - 2x}{3 - \frac{1}{x}}$$

Example 1 $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(f) Simplify $\left(\frac{f}{g}\right)(x)$

$$\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)}$$

$$= \frac{6x^2 - 2x}{3 - \frac{1}{x}} \left(\frac{x}{x}\right)$$

Example 1 $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(f) Simplify $\left(\frac{f}{g}\right)(x)$

$$\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)}$$

$$= \frac{6x^2 - 2x}{3 - \frac{1}{x}} \left(\frac{x}{x}\right)$$

$$= \frac{6x^3 - 2x^2}{3x - 1}$$

Example 1 $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(f)

$$\left(\frac{f}{g}\right)(x) = \frac{6x^3 - 2x^2}{3x - 1}$$

Example 1 $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(f)

$$\begin{aligned}\left(\frac{f}{g}\right)(x) &= \frac{6x^3 - 2x^2}{3x - 1} \\ &= \frac{2x^2(3x - 1)}{3x - 1}\end{aligned}$$

Example 1 $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

(f)

$$\begin{aligned}\left(\frac{f}{g}\right)(x) &= \frac{6x^3 - 2x^2}{3x - 1} \\ &= \frac{2x^2(3x - 1)}{3x - 1} \\ &= 2x^2\end{aligned}$$

Objectives

- 1 Perform arithmetic operations to functions
- 2 Find the domain of the sum, difference, product, or quotient of two functions
- 3 Find the difference quotient of a function

Finding Domain

When finding the domain of the sum, difference, product, or quotient of two functions, one method is to analyze the sum/difference/product/quotient before simplifying.

Example 2

(a) Find the domain of $(f + g)(x)$ if $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

Example 2

(a) Find the domain of $(f + g)(x)$ if $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

$$(f + g)(x) = 6x^2 - 2x + 3 - \frac{1}{x}$$

Example 2

(a) Find the domain of $(f + g)(x)$ if $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

$$(f + g)(x) = 6x^2 - 2x + 3 - \frac{1}{x}$$

$$x \neq 0$$

Example 2

(a) Find the domain of $(f + g)(x)$ if $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

$$(f + g)(x) = 6x^2 - 2x + 3 - \frac{1}{x}$$

$$x \neq 0$$

$$(-\infty, 0) \cup (0, \infty)$$

Example 2

(b) Find the domain of $\left(\frac{f}{g}\right)(x)$ if $f(x) = 6x^2 - 2x$ and $g(x) = 3 - \frac{1}{x}$

Example 2

(b) Find the domain of $\left(\frac{f}{g}\right)(x)$ if $f(x) = 6x^2 - 2x$ and

$$g(x) = 3 - \frac{1}{x}$$

For $g(x)$, $x \neq 0$

Example 2

For $\frac{6x^2 - 2x}{3 - \frac{1}{x}}$:

Example 2

For $\frac{6x^2 - 2x}{3 - \frac{1}{x}}$:

$$3 - \frac{1}{x} \neq 0$$

Example 2

$$\text{For } \frac{6x^2 - 2x}{3 - \frac{1}{x}}:$$

$$3 - \frac{1}{x} \neq 0$$

$$3 \neq \frac{1}{x}$$

Example 2

For $\frac{6x^2 - 2x}{3 - \frac{1}{x}}$:

$$3 - \frac{1}{x} \neq 0$$

$$3 \neq \frac{1}{x}$$

$$3x \neq 1$$

$$x \neq \frac{1}{3}$$

Example 2

$$x \neq 0, \frac{1}{3}$$

Example 2

$$x \neq 0, \frac{1}{3}$$

$$(-\infty, 0) \cup \left(0, \frac{1}{3}\right) \cup \left(\frac{1}{3}, \infty\right)$$

Objectives

- 1 Perform arithmetic operations to functions
- 2 Find the domain of the sum, difference, product, or quotient of two functions
- 3 Find the difference quotient of a function

Difference Quotient

The **difference quotient** is fundamental to the idea of the **derivative** in Calculus.

Difference Quotient

The **difference quotient** is fundamental to the idea of the **derivative** in Calculus.

For a given function, f , the difference quotient of f is

$$\frac{f(x + h) - f(x)}{h}$$

Difference Quotient

The **difference quotient** is fundamental to the idea of the **derivative** in Calculus.

For a given function, f , the difference quotient of f is

$$\frac{f(x + h) - f(x)}{h}$$

We will break it down into 3 steps:

- 1 Evaluate $f(x + h)$

Difference Quotient

The **difference quotient** is fundamental to the idea of the **derivative** in Calculus.

For a given function, f , the difference quotient of f is

$$\frac{f(x + h) - f(x)}{h}$$

We will break it down into 3 steps:

- 1 Evaluate $f(x + h)$
- 2 Subtract original function from that

Difference Quotient

The **difference quotient** is fundamental to the idea of the **derivative** in Calculus.

For a given function, f , the difference quotient of f is

$$\frac{f(x + h) - f(x)}{h}$$

We will break it down into 3 steps:

- 1 Evaluate $f(x + h)$
- 2 Subtract original function from that
- 3 Divide that result by h

Example 3

Find and simplify the difference quotient for each of the following.

(a) $f(x) = x^2 - x - 2$

Example 3

Find and simplify the difference quotient for each of the following.

(a) $f(x) = x^2 - x - 2$

$$f(x + h) = (x + h)^2 - (x + h) - 2$$

Example 3

Find and simplify the difference quotient for each of the following.

(a) $f(x) = x^2 - x - 2$

$$f(x + h) = (x + h)^2 - (x + h) - 2$$

$$= x^2 + 2hx + h^2 - x - h - 2$$

Example 3

Find and simplify the difference quotient for each of the following.

(a) $f(x) = x^2 - x - 2$

$$f(x + h) = (x + h)^2 - (x + h) - 2$$

$$= x^2 + 2hx + h^2 - x - h - 2$$

$$f(x + h) - f(x) = x^2 + 2hx + h^2 - x - h - 2 - (x^2 - x - 2)$$

Example 3

Find and simplify the difference quotient for each of the following.

(a) $f(x) = x^2 - x - 2$

$$f(x + h) = (x + h)^2 - (x + h) - 2$$

$$= x^2 + 2hx + h^2 - x - h - 2$$

$$f(x + h) - f(x) = x^2 + 2hx + h^2 - x - h - 2 - (x^2 - x - 2)$$

$$= x^2 + 2hx + h^2 - x - h - 2 - x^2 + x + 2$$

Example 3

Find and simplify the difference quotient for each of the following.

(a) $f(x) = x^2 - x - 2$

$$f(x + h) = (x + h)^2 - (x + h) - 2$$

$$= x^2 + 2hx + h^2 - x - h - 2$$

$$f(x + h) - f(x) = x^2 + 2hx + h^2 - x - h - 2 - (x^2 - x - 2)$$

$$= x^2 + 2hx + h^2 - x - h - 2 - x^2 + x + 2$$

$$= 2hx + h^2 - h$$

Example 3

$$\frac{f(x+h) - f(x)}{h} = \frac{2hx + h^2 - h}{h}$$

Example 3

$$\begin{aligned}\frac{f(x+h) - f(x)}{h} &= \frac{2hx + h^2 - h}{h} \\ &= 2x + h - 1\end{aligned}$$

Example 3

$$(b) \quad g(x) = \frac{3}{2x+1}$$

Example 3

$$(b) \quad g(x) = \frac{3}{2x+1}$$

$$g(x+h) = \frac{3}{2(x+h)+1}$$

Example 3

$$(b) \quad g(x) = \frac{3}{2x+1}$$

$$g(x+h) = \frac{3}{2(x+h)+1}$$

$$= \frac{3}{2x+2h+1}$$

Example 3

$$(b) \quad g(x) = \frac{3}{2x+1}$$

$$g(x+h) = \frac{3}{2(x+h)+1}$$

$$= \frac{3}{2x+2h+1}$$

$$g(x+h) - g(x) = \frac{3}{2x+2h+1} - \frac{3}{2x+1}$$

Example 3

$$(b) \quad g(x) = \frac{3}{2x+1}$$

$$g(x+h) = \frac{3}{2(x+h)+1}$$

$$= \frac{3}{2x+2h+1}$$

$$g(x+h) - g(x) = \frac{3}{2x+2h+1} - \frac{3}{2x+1}$$

$$\frac{g(x+h) - g(x)}{h} = \frac{\frac{3}{2x+2h+1} - \frac{3}{2x+1}}{h}$$

Example 3

$$\frac{g(x+h) - g(x)}{h} = \frac{\frac{3}{2x+2h+1}}{h} - \frac{\frac{3}{2x+1}}{h}$$

Example 3

$$\frac{g(x+h) - g(x)}{h} = \frac{\frac{3}{2x+2h+1}}{h} - \frac{\frac{3}{2x+1}}{h} \left(\frac{(2x+2h+1)(2x+1)}{(2x+2h+1)(2x+1)} \right)$$

Example 3

$$\begin{aligned}\frac{g(x+h) - g(x)}{h} &= \frac{\frac{3}{2x+2h+1} - \frac{3}{2x+1}}{h} \left(\frac{(2x+2h+1)(2x+1)}{(2x+2h+1)(2x+1)} \right) \\ &= \frac{3(2x+1) - 3(2x+2h+1)}{h(2x+2h+1)(2x+1)}\end{aligned}$$

Example 3

$$\begin{aligned}\frac{g(x+h) - g(x)}{h} &= \frac{\frac{3}{2x+2h+1}}{h} - \frac{\frac{3}{2x+1}}{h} \left(\frac{(2x+2h+1)(2x+1)}{(2x+2h+1)(2x+1)} \right) \\&= \frac{3(2x+1) - 3(2x+2h+1)}{h(2x+2h+1)(2x+1)} \\&= \frac{6x+3-6x-6h-3}{h(2x+2h+1)(2x+1)}\end{aligned}$$

Example 3

$$\begin{aligned}\frac{g(x+h) - g(x)}{h} &= \frac{\frac{3}{2x+2h+1} - \frac{3}{2x+1}}{h} \left(\frac{(2x+2h+1)(2x+1)}{(2x+2h+1)(2x+1)} \right) \\&= \frac{3(2x+1) - 3(2x+2h+1)}{h(2x+2h+1)(2x+1)} \\&= \frac{6x+3-6x-6h-3}{h(2x+2h+1)(2x+1)} \\&= \frac{-6h}{h(2x+2h+1)(2x+1)}\end{aligned}$$

Example 3

$$\frac{g(x+h) - g(x)}{h} = \frac{-6h}{h(2x+2h+1)(2x+1)}$$

Example 3

$$\begin{aligned}\frac{g(x+h) - g(x)}{h} &= \frac{-6h}{h(2x+2h+1)(2x+1)} \\ &= \frac{-6}{(2x+2h+1)(2x+1)}\end{aligned}$$

Example 3

$$(c) \quad r(x) = \sqrt{x}$$

Example 3

$$(c) \quad r(x) = \sqrt{x}$$

$$r(x + h) = \sqrt{x + h}$$

Example 3

$$(c) \quad r(x) = \sqrt{x}$$

$$r(x + h) = \sqrt{x + h}$$

$$r(x + h) - r(x) = \sqrt{x + h} - \sqrt{x}$$

Example 3

$$(c) \quad r(x) = \sqrt{x}$$

$$r(x + h) = \sqrt{x + h}$$

$$r(x + h) - r(x) = \sqrt{x + h} - \sqrt{x}$$

$$\frac{r(x + h) - r(x)}{h} = \frac{\sqrt{x + h} - \sqrt{x}}{h}$$

Example 3

$$(c) \quad r(x) = \sqrt{x}$$

$$r(x+h) = \sqrt{x+h}$$

$$r(x+h) - r(x) = \sqrt{x+h} - \sqrt{x}$$

$$\frac{r(x+h) - r(x)}{h} = \frac{\sqrt{x+h} - \sqrt{x}}{h} \left(\frac{\sqrt{x+h} + \sqrt{x}}{\sqrt{x+h} + \sqrt{x}} \right)$$

Example 3

$$(c) \quad r(x) = \sqrt{x}$$

$$r(x+h) = \sqrt{x+h}$$

$$r(x+h) - r(x) = \sqrt{x+h} - \sqrt{x}$$

$$\frac{r(x+h) - r(x)}{h} = \frac{\sqrt{x+h} - \sqrt{x}}{h} \left(\frac{\sqrt{x+h} + \sqrt{x}}{\sqrt{x+h} + \sqrt{x}} \right)$$

$$= \frac{(\sqrt{x+h})^2 - (\sqrt{x})^2}{h(\sqrt{x+h} + \sqrt{x})}$$

Example 3

$$(c) \quad r(x) = \sqrt{x}$$

$$r(x+h) = \sqrt{x+h}$$

$$r(x+h) - r(x) = \sqrt{x+h} - \sqrt{x}$$

$$\frac{r(x+h) - r(x)}{h} = \frac{\sqrt{x+h} - \sqrt{x}}{h} \left(\frac{\sqrt{x+h} + \sqrt{x}}{\sqrt{x+h} + \sqrt{x}} \right)$$

$$= \frac{(\sqrt{x+h})^2 - (\sqrt{x})^2}{h(\sqrt{x+h} + \sqrt{x})}$$

$$= \frac{x+h-x}{h(\sqrt{x+h} + \sqrt{x})}$$

Example 3

$$\frac{r(x+h) - r(x)}{h} = \frac{x+h-x}{h(\sqrt{x+h} + \sqrt{x})}$$

Example 3

$$\begin{aligned}\frac{r(x+h) - r(x)}{h} &= \frac{x+h-x}{h(\sqrt{x+h} + \sqrt{x})} \\ &= \frac{h}{h(\sqrt{x+h} + \sqrt{x})}\end{aligned}$$

Example 3

$$\begin{aligned}\frac{r(x+h) - r(x)}{h} &= \frac{x+h-x}{h(\sqrt{x+h} + \sqrt{x})} \\ &= \frac{h}{h(\sqrt{x+h} + \sqrt{x})} \\ &= \frac{1}{\sqrt{x+h} + \sqrt{x}}\end{aligned}$$