

# Sequences

# Objectives

- 1 List the terms of an explicit sequence.
- 2 List the terms of a recursive sequence.
- 3 Understand factorial notation
- 4 Find terms of an arithmetic sequence
- 5 Find terms of a geometric sequence

# Sequences

An **sequence**  $\{a_n\}$  is a function whose domain is the set of non-negative integers (*i.e.*  $0, 1, 2, \dots$ ).

We will mostly stick with positive integers ( $1, 2, 3, \dots$ ).

# Sequences

An **sequence**  $\{a_n\}$  is a function whose domain is the set of non-negative integers (*i.e.*  $0, 1, 2, \dots$ ).

We will mostly stick with positive integers ( $1, 2, 3, \dots$ ).

The function values (**terms**) are  $a_1, a_2, a_3, a_4, \dots, a_n$ ; or  $a_0, a_1, a_2, a_3, \dots, a_n$  if using a starting value of 0.

# Sequences

The value  $a(n)$  is often written as  $a_n$  and is called the  $n^{\text{th}}$  term of the sequence.

The sequence is denoted by  $a_n$  or  $\{a_n\}_{n=1}^{\infty}$ .

# Sequences

The value  $a(n)$  is often written as  $a_n$  and is called the  $n^{\text{th}}$  term of the sequence.

The sequence is denoted by  $a_n$  or  $\{a_n\}_{n=1}^{\infty}$ .

Sequences that consist of only the first  $n$  positive integers are **finite sequences**.

# Sequences

The value  $a(n)$  is often written as  $a_n$  and is called the  $n^{\text{th}}$  term of the sequence.

The sequence is denoted by  $a_n$  or  $\{a_n\}_{n=1}^{\infty}$ .

Sequences that consist of only the first  $n$  positive integers are **finite sequences**.

Sequences that are not finite are infinite.

# Sequences

The value  $a(n)$  is often written as  $a_n$  and is called the  $n^{\text{th}}$  term of the sequence.

The sequence is denoted by  $a_n$  or  $\{a_n\}_{n=1}^{\infty}$ .

Sequences that consist of only the first  $n$  positive integers are **finite sequences**.

Sequences that are not finite are infinite.

We list out the terms of a sequence by substituting values of  $n$  into the sequence's definition.



# Sequences

The value  $a(n)$  is often written as  $a_n$  and is called the  $n^{\text{th}}$  term of the sequence.

The sequence is denoted by  $a_n$  or  $\{a_n\}_{n=1}^{\infty}$ .

Sequences that consist of only the first  $n$  positive integers are **finite sequences**.

Sequences that are not finite are infinite.

We list out the terms of a sequence by substituting values of  $n$  into the sequence's definition.

*Note:* We can use any variable to name a sequence  $a_n$ ,  $b_n$ ,  $c_n$ , etc. We can also use any letter we want instead of  $n$ .

## Example 1

Write the first 4 terms of the sequence whose explicit rule is given.

(a)  $a_n = \frac{5^{n-1}}{3^n}$

## Example 1

Write the first 4 terms of the sequence whose explicit rule is given.

$$(a) \quad a_n = \frac{5^{n-1}}{3^n}$$

$$a(1) = \frac{5^{1-1}}{3^1} = \frac{1}{3}$$

## Example 1

Write the first 4 terms of the sequence whose explicit rule is given.

$$(a) \quad a_n = \frac{5^{n-1}}{3^n}$$

$$a(1) = \frac{5^{1-1}}{3^1} = \frac{1}{3}$$

$$a(2) = \frac{5^{2-1}}{3^2} = \frac{5}{9}$$

## Example 1

Write the first 4 terms of the sequence whose explicit rule is given.

$$(a) \quad a_n = \frac{5^{n-1}}{3^n}$$

$$a(1) = \frac{5^{1-1}}{3^1} = \frac{1}{3}$$

$$a(2) = \frac{5^{2-1}}{3^2} = \frac{5}{9}$$

$$a(3) = \frac{5^{3-1}}{3^3} = \frac{25}{27}$$

## Example 1

Write the first 4 terms of the sequence whose explicit rule is given.

$$(a) \quad a_n = \frac{5^{n-1}}{3^n}$$

$$a(1) = \frac{5^{1-1}}{3^1} = \frac{1}{3}$$

$$a(2) = \frac{5^{2-1}}{3^2} = \frac{5}{9}$$

$$a(3) = \frac{5^{3-1}}{3^3} = \frac{25}{27}$$

$$a(4) = \frac{5^{4-1}}{3^4} = \frac{125}{81}$$

## Example 1

$$(b) \quad b_k = \frac{(-1)^k}{2k+1}, \quad k \geq 0$$

## Example 1

$$(b) \quad b_k = \frac{(-1)^k}{2k+1}, \quad k \geq 0$$

$$b(0) = \frac{(-1)^0}{2(0)+1} = 1$$



## Example 1

$$(b) \quad b_k = \frac{(-1)^k}{2k+1}, \quad k \geq 0$$

$$b(0) = \frac{(-1)^0}{2(0)+1} = 1$$

$$b(1) = \frac{(-1)^1}{2(1)+1} = -\frac{1}{3}$$

## Example 1

$$(b) \quad b_k = \frac{(-1)^k}{2k+1}, \quad k \geq 0$$

$$b(0) = \frac{(-1)^0}{2(0)+1} = 1$$

$$b(1) = \frac{(-1)^1}{2(1)+1} = -\frac{1}{3}$$

$$b(2) = \frac{(-1)^2}{2(2)+1} = \frac{1}{5}$$

## Example 1

$$(b) \quad b_k = \frac{(-1)^k}{2k+1}, \quad k \geq 0$$

$$b(0) = \frac{(-1)^0}{2(0)+1} = 1$$

$$b(1) = \frac{(-1)^1}{2(1)+1} = -\frac{1}{3}$$

$$b(2) = \frac{(-1)^2}{2(2)+1} = \frac{1}{5}$$

$$b(3) = \frac{(-1)^3}{2(3)+1} = -\frac{1}{7}$$

## Example 1

$$(c) \quad \{2n - 1\}_{n=1}^{\infty}$$

## Example 1

$$(c) \quad \{2n - 1\}_{n=1}^{\infty}$$

$$c(1) = 2(1) - 1 = 1$$

## Example 1

$$(c) \quad \{2n - 1\}_{n=1}^{\infty}$$

$$c(1) = 2(1) - 1 = 1$$

$$c(2) = 2(2) - 1 = 3$$

## Example 1

$$(c) \quad \{2n - 1\}_{n=1}^{\infty}$$

$$c(1) = 2(1) - 1 = 1$$

$$c(2) = 2(2) - 1 = 3$$

$$c(3) = 2(3) - 1 = 5$$

## Example 1

$$(c) \quad \{2n - 1\}_{n=1}^{\infty}$$

$$c(1) = 2(1) - 1 = 1$$

$$c(2) = 2(2) - 1 = 3$$

$$c(3) = 2(3) - 1 = 5$$

$$c(4) = 2(4) - 1 = 7$$



## Example 1

$$(d) \quad \left\{ \frac{1 + (-1)^i}{i} \right\}_{i=2}^{\infty}$$

## Example 1

$$(d) \quad \left\{ \frac{1 + (-1)^i}{i} \right\}_{i=2}^{\infty}$$

$$c(2) = \frac{1 + (-1)^2}{2} = 1$$

## Example 1

$$(d) \quad \left\{ \frac{1 + (-1)^i}{i} \right\}_{i=2}^{\infty}$$

$$c(2) = \frac{1 + (-1)^2}{2} = 1$$

$$c(3) = \frac{1 + (-1)^3}{3} = 0$$

## Example 1

$$(d) \quad \left\{ \frac{1 + (-1)^i}{i} \right\}_{i=2}^{\infty}$$

$$c(2) = \frac{1 + (-1)^2}{2} = 1$$

$$c(3) = \frac{1 + (-1)^3}{3} = 0$$

$$c(4) = \frac{1 + (-1)^4}{4} = \frac{1}{2}$$

## Example 1

$$(d) \quad \left\{ \frac{1 + (-1)^i}{i} \right\}_{i=2}^{\infty}$$

$$c(2) = \frac{1 + (-1)^2}{2} = 1$$

$$c(3) = \frac{1 + (-1)^3}{3} = 0$$


$$c(4) = \frac{1 + (-1)^4}{4} = \frac{1}{2}$$

$$c(5) = \frac{1 + (-1)^5}{5} = 0$$

# Graphs of Sequences

We can graph a sequence by using coordinates.

$$a_n = \frac{1}{n}$$



# Objectives

- 1 List the terms of an explicit sequence.
- 2 List the terms of a recursive sequence.
- 3 Understand factorial notation
- 4 Find terms of an arithmetic sequence
- 5 Find terms of a geometric sequence

# Recursive Sequences

A **recursive sequence** is one that is defined using previous terms.



# Recursive Sequences

A **recursive sequence** is one that is defined using previous terms.

You are given a starting term value and then a rule for generating additional terms.

# Recursive Sequences

A **recursive sequence** is one that is defined using previous terms.

You are given a starting term value and then a rule for generating additional terms.

The notations for successive terms are either

$$a_n \text{ and } a_{n+1}$$

or

$$a_{n-1} \text{ and } a_n$$

## Example 2

Write the first four terms of each sequence.

(a)  $a_1 = 5$  and  $a_n = 3a_{n-1} + 2$  for  $n \geq 2$

## Example 2

Write the first four terms of each sequence.

(a)  $a_1 = 5$  and  $a_n = 3a_{n-1} + 2$  for  $n \geq 2$

Each new term =  $(3 \times \text{previous term}) + 2$

## Example 2

Write the first four terms of each sequence.

(a)  $a_1 = 5$  and  $a_n = 3a_{n-1} + 2$  for  $n \geq 2$

Each new term =  $(3 \times \text{previous term}) + 2$

$$a_1 = 5$$

## Example 2

Write the first four terms of each sequence.

(a)  $a_1 = 5$  and  $a_n = 3a_{n-1} + 2$  for  $n \geq 2$

Each new term =  $(3 \times \text{previous term}) + 2$

$$a_1 = 5$$

$$a_2 = 3(5) + 2 = 17$$

## Example 2

Write the first four terms of each sequence.

(a)  $a_1 = 5$  and  $a_n = 3a_{n-1} + 2$  for  $n \geq 2$

Each new term =  $(3 \times \text{previous term}) + 2$

$$a_1 = 5$$

$$a_2 = 3(5) + 2 = 17$$

$$a_3 = 3(17) + 2 = 53$$

## Example 2

Write the first four terms of each sequence.

(a)  $a_1 = 5$  and  $a_n = 3a_{n-1} + 2$  for  $n \geq 2$

Each new term =  $(3 \times \text{previous term}) + 2$

$$a_1 = 5$$

$$a_2 = 3(5) + 2 = 17$$

$$a_3 = 3(17) + 2 = 53$$

$$a_4 = 3(53) + 2 = 161$$



## Example 2

(b)  $a_1 = 7$  and  $a_{n+1} = 2 - a_n$  for  $n \geq 1$

## Example 2

(b)  $a_1 = 7$  and  $a_{n+1} = 2 - a_n$  for  $n \geq 1$

Each new term =  $2 -$  previous term

## Example 2

(b)  $a_1 = 7$  and  $a_{n+1} = 2 - a_n$  for  $n \geq 1$

Each new term =  $2 -$  previous term

$$a_1 = 7$$

## Example 2

(b)  $a_1 = 7$  and  $a_{n+1} = 2 - a_n$  for  $n \geq 1$

Each new term =  $2 -$  previous term

$$a_1 = 7$$

$$a_2 = 2 - 7 = -5$$

## Example 2

(b)  $a_1 = 7$  and  $a_{n+1} = 2 - a_n$  for  $n \geq 1$

Each new term =  $2 -$  previous term

$$a_1 = 7$$

$$a_2 = 2 - 7 = -5$$

$$a_3 = 2 - (-5) = 7$$

## Example 2

(b)  $a_1 = 7$  and  $a_{n+1} = 2 - a_n$  for  $n \geq 1$

Each new term =  $2 -$  previous term

$$a_1 = 7$$

$$a_2 = 2 - 7 = -5$$

$$a_3 = 2 - (-5) = 7$$

$$a_4 = 2 - 7 = -5$$

## Example 2

(c)  $f_0 = 1$  and  $f_n = n \cdot f_{n-1}$  for  $n \geq 1$

## Example 2

(c)  $f_0 = 1$  and  $f_n = n \cdot f_{n-1}$  for  $n \geq 1$

Each new term = (term number)  $\times$  (previous term)



## Example 2

(c)  $f_0 = 1$  and  $f_n = n \cdot f_{n-1}$  for  $n \geq 1$

Each new term = (term number)  $\times$  (previous term)

$$f_0 = 1$$

## Example 2

(c)  $f_0 = 1$  and  $f_n = n \cdot f_{n-1}$  for  $n \geq 1$

Each new term = (term number)  $\times$  (previous term)

$$f_0 = 1$$

$$f_1 = 1(1) = 1$$

## Example 2

(c)  $f_0 = 1$  and  $f_n = n \cdot f_{n-1}$  for  $n \geq 1$

Each new term = (term number)  $\times$  (previous term)

$$f_0 = 1$$

$$f_1 = 1(1) = 1$$

$$f_2 = 2(1) = 2$$

## Example 2

(c)  $f_0 = 1$  and  $f_n = n \cdot f_{n-1}$  for  $n \geq 1$

Each new term = (term number)  $\times$  (previous term)

$$f_0 = 1$$

$$f_1 = 1(1) = 1$$

$$f_2 = 2(1) = 2$$

$$f_3 = 3(2) = 6$$

# Objectives

- 1 List the terms of an explicit sequence.
- 2 List the terms of a recursive sequence.
- 3 Understand factorial notation**
- 4 Find terms of an arithmetic sequence
- 5 Find terms of a geometric sequence

# Factorial Notation

Products of consecutive positive integers can be expressed using factorial notation.

# Factorial Notation

Products of consecutive positive integers can be expressed using factorial notation.

If  $n$  is a positive integer, then

$$n! = n(n-1)(n-2) \cdots (3)(2)(1)$$

By definition,  $0! = 1$ .

# Factorial Values

Factorial values grow very quickly.

$$1! = 1$$

$$2! = 2 \cdot 1 = 2$$

$$3! = 3 \cdot 2 \cdot 1 = 6$$

$$4! = 4 \cdot 3 \cdot 2 \cdot 1 = 24$$

$$5! = 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 = 120$$

$$6! = 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 = 720$$



# Grouping Symbols

Unless grouping symbols, like  $()$  are involved, factorials only affect the number or variable they follow.

$$2 \cdot 3! = 2(3 \cdot 2 \cdot 1) = 12, \text{ but } (2 \cdot 3)! = 6! = 720.$$

## Example 3

Write the first 4 terms of each sequence.

(a) 
$$a_n = \frac{2^n}{(n-1)!}$$

## Example 3

Write the first 4 terms of each sequence.

$$(a) \quad a_n = \frac{2^n}{(n-1)!}$$

$$a_1 = \frac{2^1}{(1-1)!} = 2$$

## Example 3

Write the first 4 terms of each sequence.

$$(a) \quad a_n = \frac{2^n}{(n-1)!}$$

$$a_1 = \frac{2^1}{(1-1)!} = 2$$

$$a_2 = \frac{2^2}{(2-1)!} = 4$$

## Example 3

Write the first 4 terms of each sequence.

$$(a) \quad a_n = \frac{2^n}{(n-1)!}$$

$$a_1 = \frac{2^1}{(1-1)!} = 2$$

$$a_2 = \frac{2^2}{(2-1)!} = 4$$

$$a_3 = \frac{2^3}{(3-1)!} = 4$$

## Example 3

Write the first 4 terms of each sequence.

$$(a) \quad a_n = \frac{2^n}{(n-1)!}$$

$$a_1 = \frac{2^1}{(1-1)!} = 2$$

$$a_2 = \frac{2^2}{(2-1)!} = 4$$

$$a_3 = \frac{2^3}{(3-1)!} = 4$$

$$a_4 = \frac{2^4}{(4-1)!} = \frac{8}{3}$$

## Example 3

Write the first 4 terms of each sequence.

$$(b) \quad b_n = \frac{20}{(n+1)!}$$

## Example 3

Write the first 4 terms of each sequence.

$$(b) \quad b_n = \frac{20}{(n+1)!}$$

$$b_1 = \frac{20}{(1+1)!} = 10$$



## Example 3

Write the first 4 terms of each sequence.

$$(b) \quad b_n = \frac{20}{(n+1)!}$$

$$b_1 = \frac{20}{(1+1)!} = 10$$

$$b_2 = \frac{20}{(2+1)!} = \frac{10}{3}$$

## Example 3

Write the first 4 terms of each sequence.

$$(b) \quad b_n = \frac{20}{(n+1)!}$$

$$b_1 = \frac{20}{(1+1)!} = 10$$

$$b_2 = \frac{20}{(2+1)!} = \frac{10}{3}$$

$$b_3 = \frac{20}{(3+1)!} = \frac{5}{6}$$

## Example 3

Write the first 4 terms of each sequence.

$$(b) \quad b_n = \frac{20}{(n+1)!}$$

$$b_1 = \frac{20}{(1+1)!} = 10$$

$$b_2 = \frac{20}{(2+1)!} = \frac{10}{3}$$

$$b_3 = \frac{20}{(3+1)!} = \frac{5}{6}$$

$$b_4 = \frac{20}{(4+1)!} = \frac{1}{6}$$

# Simplifying Factorial Expressions

When simplifying factorial expressions, it helps to remember that

$$n! = n(n-1)(n-2) \cdots (3)(2)(1)$$

## Example 4

Simplify each.

(a)  $\frac{(n+1)!}{n!}$

## Example 4

Simplify each.

$$(a) \quad \frac{(n+1)!}{n!}$$

$$\frac{(n+1)!}{n!} = \frac{(n+1)(n)(n-1)(n-2) \cdots (2)(1)}{n(n-1)(n-2) \cdots (2)(1)}$$

## Example 4

Simplify each.

$$(a) \quad \frac{(n+1)!}{n!}$$

$$\frac{(n+1)!}{n!} = \frac{(n+1)(n)(n-1)(n-2) \cdots (2)(1)}{n(n-1)(n-2) \cdots (2)(1)}$$

$$= \frac{(n+1)\cancel{(n)}\cancel{(n-1)}\cancel{(n-2)} \cdots \cancel{(2)}\cancel{(1)}}{\cancel{(n)}\cancel{(n-1)}\cancel{(n-2)} \cdots \cancel{(2)}\cancel{(1)}}$$

## Example 4

Simplify each.

$$(a) \quad \frac{(n+1)!}{n!}$$

$$\frac{(n+1)!}{n!} = \frac{(n+1)(n)(n-1)(n-2) \cdots (2)(1)}{n(n-1)(n-2) \cdots (2)(1)}$$

$$= \frac{(n+1)\cancel{(n)}\cancel{(n-1)}\cancel{(n-2)} \cdots \cancel{(2)}\cancel{(1)}}{\cancel{(n)}\cancel{(n-1)}\cancel{(n-2)} \cdots \cancel{(2)}\cancel{(1)}}$$

$$= n+1$$



## Example 4

Simplify each.

(b)  $\frac{n!}{(n-1)!}$

## Example 4

Simplify each.

$$(b) \quad \frac{n!}{(n-1)!}$$

$$\frac{n!}{(n-1)!} = \frac{n(n-1)(n-2)\cdots(2)(1)}{(n-1)(n-2)\cdots(2)(1)}$$

## Example 4

Simplify each.

$$(b) \quad \frac{n!}{(n-1)!}$$

$$\frac{n!}{(n-1)!} = \frac{n(n-1)(n-2)\cdots(2)(1)}{(n-1)(n-2)\cdots(2)(1)}$$

$$= \frac{n\cancel{(n-1)}\cancel{(n-2)}\cdots\cancel{(2)}\cancel{(1)}}{\cancel{(n-1)}\cancel{(n-2)}\cdots\cancel{(2)}\cancel{(1)}}$$

## Example 4

Simplify each.

$$(b) \quad \frac{n!}{(n-1)!}$$

$$\frac{n!}{(n-1)!} = \frac{n(n-1)(n-2) \cdots (2)(1)}{(n-1)(n-2) \cdots (2)(1)}$$

$$= \frac{n \cancel{(n-1)} \cancel{(n-2)} \cdots \cancel{(2)} \cancel{(1)}}{\cancel{(n-1)} \cancel{(n-2)} \cdots \cancel{(2)} \cancel{(1)}}$$

$$= n$$

# Objectives

- 1 List the terms of an explicit sequence.
- 2 List the terms of a recursive sequence.
- 3 Understand factorial notation
- 4 Find terms of an arithmetic sequence**
- 5 Find terms of a geometric sequence

# Arithmetic Sequences

When the difference between consecutive terms of a sequence is always the same, such as 8, 11, 14, 17,  $\dots$ , that sequence is an **arithmetic sequence**.

# Arithmetic Sequences

When the difference between consecutive terms of a sequence is always the same, such as 8, 11, 14, 17,  $\dots$ , that sequence is an **arithmetic sequence**.

The difference between those consecutive terms is called the **common difference**.

# Arithmetic Sequences

When the difference between consecutive terms of a sequence is always the same, such as 8, 11, 14, 17,  $\dots$ , that sequence is an **arithmetic sequence**.

The difference between those consecutive terms is called the **common difference**.

To find the common difference of an arithmetic sequence, **subtract** any two consecutive terms.




# Arithmetic Sequences and Linear Functions

Arithmetic sequences model linear functions.

# Arithmetic Sequences and Linear Functions

Arithmetic sequences model linear functions.

The following graph shows the first 4 terms of the sequence 8, 11, 14, 17:



# Find the Rule for the Sequence

Notice that the common difference between terms (3 in this case) is the same as the slope of the line.

# Find the Rule for the Sequence

Notice that the common difference between terms (3 in this case) is the same as the slope of the line.

If we subtract that 3 from the first term, we get 5; which is the  $y$ -intercept of the graph.

# Find the Rule for the Sequence

Notice that the common difference between terms (3 in this case) is the same as the slope of the line.

If we subtract that 3 from the first term, we get 5; which is the  $y$ -intercept of the graph.

Thus, a sequence rule for this sequence is

$$y = 3x + 5 \quad \text{or} \quad a_n = 3n + 5 \text{ for } n \geq 1$$

## Example 5

Find an explicit formula for the  $n^{\text{th}}$  term of each of the following sequences.

(a)  $-31, -24, -17, -10, \dots$

## Example 5

Find an explicit formula for the  $n^{\text{th}}$  term of each of the following sequences.

(a)  $-31, -24, -17, -10, \dots$

Common Difference:  $-24 - (-31) = 7$

## Example 5

Find an explicit formula for the  $n^{\text{th}}$  term of each of the following sequences.

(a)  $-31, -24, -17, -10, \dots$

Common Difference:  $-24 - (-31) = 7$

y-intercept:  $-31 - 7 = -38$



## Example 5

Find an explicit formula for the  $n^{\text{th}}$  term of each of the following sequences.

(a)  $-31, -24, -17, -10, \dots$

Common Difference:  $-24 - (-31) = 7$

y-intercept:  $-31 - 7 = -38$

$$a_n = 7n - 38$$

## Example 5

(b) 37, 67, 97, 127, ...

## Example 5

(b) 37, 67, 97, 127, ...

Common Difference:  $67 - 37 = 30$

## Example 5

(b) 37, 67, 97, 127, ...

Common Difference:  $67 - 37 = 30$

y-intercept:  $37 - 30 = 7$

## Example 5

(b) 37, 67, 97, 127, ...

Common Difference:  $67 - 37 = 30$

y-intercept:  $37 - 30 = 7$

$$b_n = 30n + 7$$

# Objectives

- 1 List the terms of an explicit sequence.
- 2 List the terms of a recursive sequence.
- 3 Understand factorial notation
- 4 Find terms of an arithmetic sequence
- 5 Find terms of a geometric sequence

# Geometric Sequences

A **geometric sequence** is one in which each term after the first is obtained by multiplying the preceding term by a fixed nonzero constant  $r$  (called the **common ratio**).

# Geometric Sequences


A **geometric sequence** is one in which each term after the first is obtained by multiplying the preceding term by a fixed nonzero constant  $r$  (called the **common ratio**).

Geometric sequences model exponential growth or decay.



# Model of a Geometric Sequence

The following graph shows the first 4 terms of the sequence 3, 6, 12, 18:



# Finding Rule for Geometric Sequences

Notice each point's  $y$ -coordinate is twice the previous one. This indicates that the common ratio,  $r$ , is the base of the exponential function, or 2.

# Finding Rule for Geometric Sequences

Notice each point's  $y$ -coordinate is twice the previous one. This indicates that the common ratio,  $r$ , is the base of the exponential function, or 2.

To find the  $y$ -intercept, divide the first term by the base of 2 to get 1.5.

## Finding Rule for Geometric Sequences

Notice each point's  $y$ -coordinate is twice the previous one. This indicates that the common ratio,  $r$ , is the base of the exponential function, or 2.

To find the  $y$ -intercept, divide the first term by the base of 2 to get 1.5.

Thus, the function can be modeled as

$$y = 1.5(2)^x \quad \text{or} \quad a_n = 1.5(2)^n \text{ for } n \geq 1$$

# Finding Rule for Geometric Sequences

Notice each point's  $y$ -coordinate is twice the previous one. This indicates that the common ratio,  $r$ , is the base of the exponential function, or 2.

To find the  $y$ -intercept, divide the first term by the base of 2 to get 1.5.

Thus, the function can be modeled as

$$y = 1.5(2)^x \quad \text{or} \quad a_n = 1.5(2)^n \text{ for } n \geq 1$$

**\*\*Note:** Unlike exponential functions, geometric sequences can have negative common ratios.

## Example 6

Find the 8th term of each geometric sequence.

(a)  $a_1 = -4$   $r = -2$

## Example 6

Find the 8th term of each geometric sequence.

(a)  $a_1 = -4$   $r = -2$

$y$ -intercept:  $-2$

## Example 6

Find the 8th term of each geometric sequence.

(a)  $a_1 = -4$   $r = -2$

y-intercept:  $-2$

$$a_n = -2(-2)^n$$



## Example 6

Find the 8th term of each geometric sequence.

(a)  $a_1 = -4$   $r = -2$

y-intercept:  $-2$

$$a_n = -2(-2)^n$$

$$a_8 = -2(-2)^8$$

## Example 6

Find the 8th term of each geometric sequence.

(a)  $a_1 = -4$   $r = -2$

y-intercept:  $-2$

$$a_n = -2(-2)^n$$

$$a_8 = -2(-2)^8$$

$$a_8 = -512$$

## Example 6

Find the 8th term of each geometric sequence.

(b)  $a_1 = 80$   $r = 0.5$

## Example 6

Find the 8th term of each geometric sequence.

(b)  $a_1 = 80$   $r = 0.5$

y-intercept: 160

## Example 6

Find the 8th term of each geometric sequence.

(b)  $a_1 = 80$   $r = 0.5$

y-intercept: 160

$$a_n = 160(0.5)^n$$

## Example 6

Find the 8th term of each geometric sequence.

(b)  $a_1 = 80$   $r = 0.5$

y-intercept: 160

$$a_n = 160 (0.5)^n$$

$$a_8 = 160 (0.5)^8$$

## Example 6

Find the 8th term of each geometric sequence.

(b)  $a_1 = 80$   $r = 0.5$

y-intercept: 160

$$a_n = 160 (0.5)^n$$

$$a_8 = 160 (0.5)^8$$

$$a_8 = \frac{5}{8}$$

## Example 6

$$(c) \quad a_3 = -48 \quad a_5 = -768$$




## Example 6

(c)  $a_3 = -48$   $a_5 = -768$

$$\underline{\hspace{1cm}} \quad \underline{\hspace{1cm}} \quad \underline{\hspace{1cm}} \quad \frac{-48}{a_3} \quad \underline{\hspace{1cm}} \quad \frac{-768}{a_5}$$


## Example 6

(c)  $a_3 = -48$   $a_5 = -768$



## Example 6


(c)  $a_3 = -48$   $a_5 = -768$



$$-48r^2 = -768$$

## Example 6

(c)  $a_3 = -48$   $a_5 = -768$




$$-48r^2 = -768$$

$$r^2 = 16$$

## Example 6

(c)  $a_3 = -48$   $a_5 = -768$




$$-48r^2 = -768$$

$$r^2 = 16$$

$$r = \pm 4$$

## Example 6

(c)  $a_3 = -48$   $a_5 = -768$




$$-48r^2 = -768$$

$$r^2 = 16$$

$$r = \pm 4$$

## Example 6

(c)  $a_3 = -48$   $a_5 = -768$



$$-48r^2 = -768$$

$$r^2 = 16$$

$$r = \pm 4$$

Example 6     $a_3 = -48$      $a_5 = -768$

$$64a_0 = -48$$



Example 6     $a_3 = -48$      $a_5 = -768$

$$64a_0 = -48$$

$$a_0 = -0.75$$

## Example 6 $a_3 = -48$ $a_5 = -768$

$$64a_0 = -48$$

$$a_0 = -0.75$$

$$a_n = -0.75(4)^n$$

## Example 6 $a_3 = -48$ $a_5 = -768$

$$64a_0 = -48$$

$$a_0 = -0.75$$

$$a_n = -0.75(4)^n$$

$$a_8 = -0.75(4)^8$$

## Example 6 $a_3 = -48$ $a_5 = -768$

$$64a_0 = -48$$

$$a_0 = -0.75$$

$$a_n = -0.75(4)^n$$

$$a_8 = -0.75(4)^8$$

$$= -49,152$$

## Example 6

$$(d) \quad a_1 = 5 \quad a_4 = 16.875$$

## Example 6

$$(d) \quad a_1 = 5 \quad a_4 = 16.875$$

$$5r^3 = 16.875$$

## Example 6

$$(d) \quad a_1 = 5 \quad a_4 = 16.875$$

$$5r^3 = 16.875$$

$$r^3 = 3.375$$

## Example 6

$$(d) \quad a_1 = 5 \quad a_4 = 16.875$$

$$5r^3 = 16.875$$

$$r^3 = 3.375$$

$$r = 1.5$$



## Example 6

$$(d) \quad a_1 = 5 \quad a_4 = 16.875$$

$$5r^3 = 16.875$$

$$r^3 = 3.375$$

$$r = 1.5$$

$$a_0 = 5/1.5$$

## Example 6

$$(d) \quad a_1 = 5 \quad a_4 = 16.875$$

$$5r^3 = 16.875$$

$$r^3 = 3.375$$

$$r = 1.5$$

$$a_0 = 5/1.5$$

$$a_0 = \frac{10}{3}$$

## Example 6

$$a_n = \frac{10}{3} (1.5)^n$$

## Example 6

$$a_n = \frac{10}{3} (1.5)^n$$

$$a_8 = \frac{10}{3} (1.5)^8$$

## Example 6

$$a_n = \frac{10}{3} (1.5)^n$$

$$a_8 = \frac{10}{3} (1.5)^8$$

$$= \frac{10,935}{128}$$

## Example 7

Write an explicit rule for each geometric sequence.

(a) 3, 6, 12, 24, 48, ...

## Example 7

Write an explicit rule for each geometric sequence.

(a) 3, 6, 12, 24, 48, ...

Common ratio =  $6/3 = 2$

## Example 7

Write an explicit rule for each geometric sequence.

(a) 3, 6, 12, 24, 48, ...

Common ratio =  $6/3 = 2$

y-intercept = 1.5



## Example 7

Write an explicit rule for each geometric sequence.

(a) 3, 6, 12, 24, 48, ...

Common ratio =  $6/3 = 2$

y-intercept = 1.5

$$a_n = 1.5(2)^n$$

## Example 7

(b)  $2, -10, 50, -250, 1250, \dots$

## Example 7

(b)  $2, -10, 50, -250, 1250, \dots$

Common ratio  $= -10/2 = -5$

## Example 7

$$(b) \quad 2, -10, 50, -250, 1250, \dots$$

$$\text{Common ratio} = -10/2 = -5$$

$$y\text{-intercept} = -\frac{2}{5}$$

## Example 7

$$(b) \quad 2, -10, 50, -250, 1250, \dots$$

$$\text{Common ratio} = -10/2 = -5$$

$$y\text{-intercept} = -\frac{2}{5}$$

$$b_n = -\frac{2}{5}(-5)^n$$