

# Properties of Functions

# Objectives

- 1 Determine increasing, decreasing, and constant intervals of a function
- 2 Determine relative (local) maximum and minimum coordinates
- 3 Determine if a function is even or odd
- 4 Evaluate piecewise-defined functions

# Increasing Intervals

A function is **increasing** in an interval if the  $y$ -coordinates increase in value as the  $x$ -coordinates increase in value.

# Increasing Intervals

A function is **increasing** in an interval if the  $y$ -coordinates increase in value as the  $x$ -coordinates increase in value.

- Visually, the graph is moving “upward” from left to right.

# Increasing Intervals

A function is **increasing** in an interval if the  $y$ -coordinates increase in value as the  $x$ -coordinates increase in value.

- Visually, the graph is moving “upward” from left to right.
- “Machine-wise,” the outputs go up in value as the inputs go up in value.

# Increasing Intervals

A function is **increasing** in an interval if the  $y$ -coordinates increase in value as the  $x$ -coordinates increase in value.

- Visually, the graph is moving “upward” from left to right.
- “Machine-wise,” the outputs go up in value as the inputs go up in value.
- Mathematically, if  $a < b$  then  $f(a) < f(b)$ .

# Increasing Intervals

A function is **increasing** in an interval if the  $y$ -coordinates increase in value as the  $x$ -coordinates increase in value.

- Visually, the graph is moving “upward” from left to right.
- “Machine-wise,” the outputs go up in value as the inputs go up in value.
- Mathematically, if  $a < b$  then  $f(a) < f(b)$ .

**\*\*\*FOCUS ON THE X-COORDINATES\*\*\***

# Decreasing Intervals

A function is **decreasing** in an interval if the  $y$ -coordinates decrease in value as the  $x$ -coordinates increase in value.



# Decreasing Intervals

A function is **decreasing** in an interval if the  $y$ -coordinates decrease in value as the  $x$ -coordinates increase in value.

- Visually, the graph is moving “downward” from left to right.

# Decreasing Intervals

A function is **decreasing** in an interval if the  $y$ -coordinates decrease in value as the  $x$ -coordinates increase in value.

- Visually, the graph is moving “downward” from left to right.
- “Machine-wise,” the outputs go down in value as the inputs go up in value.

# Decreasing Intervals

A function is **decreasing** in an interval if the  $y$ -coordinates decrease in value as the  $x$ -coordinates increase in value.

- Visually, the graph is moving “downward” from left to right.
- “Machine-wise,” the outputs go down in value as the inputs go up in value.
- Mathematically, if  $a < b$  then  $f(a) > f(b)$ .

# Decreasing Intervals

A function is **decreasing** in an interval if the  $y$ -coordinates decrease in value as the  $x$ -coordinates increase in value.

- Visually, the graph is moving “downward” from left to right.
- “Machine-wise,” the outputs go down in value as the inputs go up in value.
- Mathematically, if  $a < b$  then  $f(a) > f(b)$ .

**\*\*\*FOCUS ON THE X-COORDINATES\*\*\***

# Constant Intervals

A function is **constant** in an interval if the  $y$ -coordinates do not change in value as the  $x$ -coordinates increase in value.

# Constant Intervals

A function is **constant** in an interval if the  $y$ -coordinates do not change in value as the  $x$ -coordinates increase in value.

- Visually, the graph is a horizontal line.

# Constant Intervals

A function is **constant** in an interval if the  $y$ -coordinates do not change in value as the  $x$ -coordinates increase in value.

- Visually, the graph is a horizontal line.
- “Machine-wise,” the outputs don’t change as the inputs go up in value.

# Constant Intervals

A function is **constant** in an interval if the  $y$ -coordinates do not change in value as the  $x$ -coordinates increase in value.

- Visually, the graph is a horizontal line.
- “Machine-wise,” the outputs don’t change as the inputs go up in value.
- Mathematically,  $f(a) = f(b)$  for all  $a$  and  $b$  in the interval.



# Constant Intervals


A function is **constant** in an interval if the  $y$ -coordinates do not change in value as the  $x$ -coordinates increase in value.

- Visually, the graph is a horizontal line.
- “Machine-wise,” the outputs don’t change as the inputs go up in value.
- Mathematically,  $f(a) = f(b)$  for all  $a$  and  $b$  in the interval.

**\*\*\*FOCUS ON THE X-COORDINATES\*\*\***


# Example 1

Determine the intervals in which the function is increasing, decreasing, and constant.



## Example 1


Determine the intervals in which the function is increasing, decreasing, and constant.



**Increasing:**

# Example 1

Determine the intervals in which the function is increasing, decreasing, and constant.




**Increasing:**

$$(-4, -2) \cup (3, 4) \cup (5, 6)$$

# Example 1

Determine the intervals in which the function is increasing, decreasing, and constant.




**Increasing:**

$$(-4, -2) \cup (3, 4) \cup (5, 6)$$

**Decreasing:**

# Example 1

Determine the intervals in which the function is increasing, decreasing, and constant.



**Increasing:**


$$(-4, -2) \cup (3, 4) \cup (5, 6)$$

**Decreasing:**

$$(-2, 3)$$

# Example 1

Determine the intervals in which the function is increasing, decreasing, and constant.



**Increasing:**

$$(-4, -2) \cup (3, 4) \cup (5, 6)$$


**Decreasing:**

$$(-2, 3)$$

**Constant:**

## Example 1

Determine the intervals in which the function is increasing, decreasing, and constant.



**Increasing:**

$$(-4, -2) \cup (3, 4) \cup (5, 6)$$

**Decreasing:**

$$(-2, 3)$$

**Constant:**

$$(4, 5)$$



# Objectives

- 1 Determine increasing, decreasing, and constant intervals of a function
- 2 Determine relative (local) maximum and minimum coordinates
- 3 Determine if a function is even or odd
- 4 Evaluate piecewise-defined functions

# Relative Maximum

A **relative (or local) maximum** is the highest point in some interval.

# Relative Maximum

A **relative (or local) maximum** is the highest point in some interval.

- It is a “top of the hill” for many functions.

# Relative Maximum

A **relative (or local) maximum** is the highest point in some interval.

- It is a “top of the hill” for many functions.
- Where the graph changes from increasing to decreasing.

# Relative Maximum

A **relative (or local) maximum** is the highest point in some interval.

- It is a “top of the hill” for many functions.
- Where the graph changes from increasing to decreasing.

An absolute, or global, maximum is the highest point in the entire domain of the function.

# Relative Maximum

A **relative (or local) maximum** is the highest point in some interval.

- It is a “top of the hill” for many functions.
- Where the graph changes from increasing to decreasing.

An absolute, or global, maximum is the highest point in the entire domain of the function.

**\*\*\*GIVE YOUR ANSWER AS AN ORDERED PAIR\*\*\***

# Relative Minimum

A **relative (or local) minimum** is the lowest point in some interval.

# Relative Minimum

A **relative (or local) minimum** is the lowest point in some interval.

- It is a “bottom of the valley” for many functions.



# Relative Minimum

A **relative (or local) minimum** is the lowest point in some interval.

- It is a “bottom of the valley” for many functions.
- Where the graph changes from decreasing to increasing.

# Relative Minimum

A **relative (or local) minimum** is the lowest point in some interval.

- It is a “bottom of the valley” for many functions.
- Where the graph changes from decreasing to increasing.

An absolute, or global, minimum is the lowest point in the entire domain of the function.

# Relative Minimum

A **relative (or local) minimum** is the lowest point in some interval.


- It is a “bottom of the valley” for many functions.
- Where the graph changes from decreasing to increasing.

An absolute, or global, minimum is the lowest point in the entire domain of the function.

**\*\*\*GIVE YOUR ANSWER AS AN ORDERED PAIR\*\*\***


## Example 2

Determine the relative minimum and relative maximum for each.  
Then determine the global minimum and global maximum.



## Example 2


Determine the relative minimum and relative maximum for each.  
Then determine the global minimum and global maximum.



**Relative Minimum**

## Example 2


Determine the relative minimum and relative maximum for each.  
Then determine the global minimum and global maximum.



**Relative Minimum**  
 $(3, -8)$

## Example 2

Determine the relative minimum and relative maximum for each.  
Then determine the global minimum and global maximum.




**Relative Minimum**  
 $(3, -8)$

**Relative Maximum:**

## Example 2

Determine the relative minimum and relative maximum for each.  
Then determine the global minimum and global maximum.




**Relative Minimum**  
 $(3, -8)$

**Relative Maximum:**  
 $(-2, 4.5)$



## Example 2

Determine the relative minimum and relative maximum for each.  
Then determine the global minimum and global maximum.




**Relative Minimum**  
 $(3, -8)$

**Relative Maximum:**  
 $(-2, 4.5)$

**Global Minimum:**

## Example 2

Determine the relative minimum and relative maximum for each.  
Then determine the global minimum and global maximum.




**Relative Minimum**  
 $(3, -8)$

**Relative Maximum:**  
 $(-2, 4.5)$

**Global Minimum:**  
 $(3, -8)$

## Example 2

Determine the relative minimum and relative maximum for each.  
Then determine the global minimum and global maximum.



**Relative Minimum**  
 $(3, -8)$


**Relative Maximum:**  
 $(-2, 4.5)$

**Global Minimum:**  
 $(3, -8)$

**Global Maximum:**

## Example 2

Determine the relative minimum and relative maximum for each.  
Then determine the global minimum and global maximum.



**Relative Minimum**  
 $(3, -8)$

**Relative Maximum:**  
 $(-2, 4.5)$

**Global Minimum:**  
 $(3, -8)$

**Global Maximum:**  
 $(6, 5.5)$

# Objectives

- 1 Determine increasing, decreasing, and constant intervals of a function
- 2 Determine relative (local) maximum and minimum coordinates
- 3 Determine if a function is even or odd
- 4 Evaluate piecewise-defined functions

# Even Functions

A function  $f$  is **even** if

$$f(-x) = f(x)$$

# Even Functions

A function  $f$  is **even** if

$$f(-x) = f(x)$$

Negative input values give you the same outputs as their positive opposites.

# Even Functions

A function  $f$  is **even** if

$$f(-x) = f(x)$$

Negative input values give you the same outputs as their positive opposites.

Even functions are symmetric with respect to the  $y$ -axis.



# Odd Functions

A function  $f$  is **odd** if

$$f(-x) = -f(x)$$

# Odd Functions

A function  $f$  is **odd** if

$$f(-x) = -f(x)$$

Negative input values give you the opposite outputs as their positive opposites.

# Odd Functions

A function  $f$  is **odd** if

$$f(-x) = -f(x)$$

Negative input values give you the opposite outputs as their positive opposites.

Odd functions are symmetric with respect to the origin.

## Example 3

Determine whether each of the following functions is even, odd, or neither.

(a)  $f(x) = \frac{5}{2 - x^2}$

## Example 3

Determine whether each of the following functions is even, odd, or neither.

$$(a) \quad f(x) = \frac{5}{2 - x^2}$$

$$f(-x) = \frac{5}{2 - (-x)^2}$$

## Example 3

Determine whether each of the following functions is even, odd, or neither.

$$(a) \quad f(x) = \frac{5}{2 - x^2}$$

$$f(-x) = \frac{5}{2 - (-x)^2}$$

$$= \frac{5}{2 - x^2}$$

## Example 3

Determine whether each of the following functions is even, odd, or neither.

$$(a) \quad f(x) = \frac{5}{2 - x^2}$$

$$f(-x) = \frac{5}{2 - (-x)^2}$$

$$= \frac{5}{2 - x^2}$$

$$= f(x)$$

## Example 3

Determine whether each of the following functions is even, odd, or neither.

$$(a) \quad f(x) = \frac{5}{2 - x^2}$$

$$f(-x) = \frac{5}{2 - (-x)^2}$$

$$= \frac{5}{2 - x^2}$$

$$= f(x)$$

$$f(x) = \frac{5}{2 - x^2} \text{ is even}$$



## Example 3

$$(b) \quad g(x) = \frac{5x}{2 - x^2}$$

## Example 3

$$(b) \quad g(x) = \frac{5x}{2 - x^2}$$

$$g(-x) = \frac{5(-x)}{2 - (-x)^2}$$

## Example 3

$$(b) \quad g(x) = \frac{5x}{2 - x^2}$$

$$g(-x) = \frac{5(-x)}{2 - (-x)^2}$$

$$= \frac{-5x}{2 - x^2}$$

## Example 3

$$(b) \quad g(x) = \frac{5x}{2 - x^2}$$

$$g(-x) = \frac{5(-x)}{2 - (-x)^2}$$

$$= \frac{-5x}{2 - x^2}$$

$$= - \left( \frac{5x}{2 - x^2} \right)$$

## Example 3

$$(b) \quad g(x) = \frac{5x}{2-x^2}$$

$$g(-x) = \frac{5(-x)}{2-(-x)^2}$$

$$= \frac{-5x}{2-x^2}$$

$$= -\left(\frac{5x}{2-x^2}\right)$$

$$= -g(x)$$

## Example 3

$$(b) \quad g(x) = \frac{5x}{2-x^2}$$

$$g(-x) = \frac{5(-x)}{2-(-x)^2}$$

$$= \frac{-5x}{2-x^2}$$

$$= -\left(\frac{5x}{2-x^2}\right)$$

$$= -g(x)$$

$$g(x) = \frac{5x}{2-x^2} \text{ is odd}$$

## Example 3

$$(c) \quad h(x) = \frac{5x}{2 - x^3}$$

## Example 3

$$(c) \quad h(x) = \frac{5x}{2 - x^3}$$

$$h(-x) = \frac{5(-x)}{2 - (-x)^3}$$



## Example 3

$$(c) \quad h(x) = \frac{5x}{2 - x^3}$$

$$\begin{aligned} h(-x) &= \frac{5(-x)}{2 - (-x)^3} \\ &= \frac{-5x}{2 + x^3} \end{aligned}$$

## Example 3

$$(c) \quad h(x) = \frac{5x}{2 - x^3}$$

$$h(-x) = \frac{5(-x)}{2 - (-x)^3}$$

$$= \frac{-5x}{2 + x^3}$$

$$= - \left( \frac{5x}{2 + x^3} \right)$$

## Example 3

$$(c) \quad h(x) = \frac{5x}{2 - x^3}$$

$$h(-x) = \frac{5(-x)}{2 - (-x)^3}$$

$$= \frac{-5x}{2 + x^3}$$

$$= - \left( \frac{5x}{2 + x^3} \right)$$

$$h(x) = \frac{5x}{2 - x^3} \text{ is neither odd nor even}$$

## Example 3

$$(d) \quad i(x) = \frac{5x}{2x - x^3}$$

## Example 3

$$(d) \quad i(x) = \frac{5x}{2x - x^3}$$

$$i(-x) = \frac{5(-x)}{2(-x) - (-x)^3}$$

## Example 3

$$(d) \quad i(x) = \frac{5x}{2x - x^3}$$

$$i(-x) = \frac{5(-x)}{2(-x) - (-x)^3}$$

$$= \frac{-5x}{-2x + x^3}$$

## Example 3

$$(d) \quad i(x) = \frac{5x}{2x - x^3}$$

$$i(-x) = \frac{5(-x)}{2(-x) - (-x)^3}$$

$$= \frac{-5x}{-2x + x^3}$$

$$= \frac{-5x}{-1(2x - x^3)}$$

## Example 3

$$(d) \quad i(x) = \frac{5x}{2x - x^3}$$

$$i(-x) = \frac{5(-x)}{2(-x) - (-x)^3}$$

$$= \frac{-5x}{-2x + x^3}$$

$$= \frac{-5x}{-1(2x - x^3)}$$

$$= \frac{5x}{2x - x^3}$$



## Example 3

$$(d) \quad i(x) = \frac{5x}{2x - x^3}$$

$$i(-x) = \frac{5(-x)}{2(-x) - (-x)^3}$$

$$= \frac{-5x}{-2x + x^3}$$

$$= \frac{-5x}{-1(2x - x^3)}$$

$$= \frac{5x}{2x - x^3}$$

$$= i(x)$$

## Example 3

$$(e) \quad j(x) = x^2 - \frac{x}{100} - 1$$

## Example 3

$$(e) \quad j(x) = x^2 - \frac{x}{100} - 1$$

$$j(-x) = (-x)^2 - \frac{-x}{100} - 1$$

## Example 3

$$(e) \quad j(x) = x^2 - \frac{x}{100} - 1$$

$$j(-x) = (-x)^2 - \frac{-x}{100} - 1$$

$$= x^2 + \frac{x}{100} - 1$$

## Example 3

$$(e) \quad j(x) = x^2 - \frac{x}{100} - 1$$


$$j(-x) = (-x)^2 - \frac{-x}{100} - 1$$

$$= x^2 + \frac{x}{100} - 1$$

$j(x)$  is neither odd nor even.

## Example 4


Given the graph of  $y = f(x)$ , find each.



(a) Domain of  $f$

## Example 4

Given the graph of  $y = f(x)$ , find each.




(a) Domain of  $f$

$[-4, 4]$

## Example 4

Given the graph of  $y = f(x)$ , find each.



(a) Domain of  $f$


$[-4, 4]$

(b) Range of  $f$



## Example 4

Given the graph of  $y = f(x)$ , find each.



(a) Domain of  $f$


$[-4, 4]$

(b) Range of  $f$

$[-3, 3]$

## Example 4


Given the graph of  $y = f(x)$ , find each.



(c) x-intercepts

## Example 4

Given the graph of  $y = f(x)$ , find each.




(c) x-intercepts

$(-2, 0)$  and  $(2, 0)$

## Example 4

Given the graph of  $y = f(x)$ , find each.




(c) x-intercepts

$(-2, 0)$  and  $(2, 0)$

(d) y-intercept

## Example 4

Given the graph of  $y = f(x)$ , find each.



(c)  $x$ -intercepts


$(-2, 0)$  and  $(2, 0)$

(d)  $y$ -intercept

$(0, 3)$

## Example 4


Given the graph of  $y = f(x)$ , find each.



(e) Zeros of  $f$

## Example 4

Given the graph of  $y = f(x)$ , find each.




(e) Zeros of  $f$

$(-2, 0)$  and  $(2, 0)$

## Example 4

Given the graph of  $y = f(x)$ , find each.



(e) Zeros of  $f$


$(-2, 0)$  and  $(2, 0)$

(f) Solve  $f(x) < 0$



## Example 4

Given the graph of  $y = f(x)$ , find each.



(e) Zeros of  $f$


$(-2, 0)$  and  $(2, 0)$

(f) Solve  $f(x) < 0$

$[-4, -2) \cup (2, 4]$

## Example 4


Given the graph of  $y = f(x)$ , find each.



(g) Determine  $f(2)$

## Example 4

Given the graph of  $y = f(x)$ , find each.




(g) Determine  $f(2)$

0

## Example 4

Given the graph of  $y = f(x)$ , find each.




(g) Determine  $f(2)$

0

(h) Solve  $f(x) = -3$

## Example 4

Given the graph of  $y = f(x)$ , find each.



(g) Determine  $f(2)$


0

(h) Solve  $f(x) = -3$

$x = -4$  and  $x = 4$

## Example 4


Given the graph of  $y = f(x)$ , find each.



(i) Number of solutions to  $f(x) = 1$

## Example 4


Given the graph of  $y = f(x)$ , find each.



(i) Number of solutions to  $f(x) = 1$

## Example 4

Given the graph of  $y = f(x)$ , find each.




(i) Number of solutions to  $f(x) = 1$

2



## Example 4

Given the graph of  $y = f(x)$ , find each.




(i) Number of solutions to  $f(x) = 1$

2

(j) Does  $f$  appear even, odd, or neither?

## Example 4

Given the graph of  $y = f(x)$ , find each.



(i) Number of solutions to  $f(x) = 1$


2

(j) Does  $f$  appear even, odd, or neither?

even

## Example 4


Given the graph of  $y = f(x)$ , find each.



(k) List intervals of increasing and decreasing.

## Example 4

Given the graph of  $y = f(x)$ , find each.




(k) List intervals of increasing and decreasing.

Increasing:

## Example 4

Given the graph of  $y = f(x)$ , find each.




(k) List intervals of increasing and decreasing.

Increasing:

$(-4, 0)$

## Example 4

Given the graph of  $y = f(x)$ , find each.



(k) List intervals of increasing and decreasing.


Increasing:

$(-4, 0)$

Decreasing:

## Example 4

Given the graph of  $y = f(x)$ , find each.



(k) List intervals of increasing and decreasing.

Increasing:

$(-4, 0)$

Decreasing:

$(0, 4)$

# Objectives

- 1 Determine increasing, decreasing, and constant intervals of a function
- 2 Determine relative (local) maximum and minimum coordinates
- 3 Determine if a function is even or odd
- 4 Evaluate piecewise-defined functions




# Piecewise-Defined Functions

Piecewise-defined functions take pieces of other functions and put them together (a la Frankenstein).

$$f(x) = \begin{cases} \sin(x - 1) + 2 & \text{if } -9 \leq x < -2 \\ x^2 - 3 & \text{if } -2 \leq x \leq 3 \\ -8x + 30 & \text{if } 3 < x < 4.5 \end{cases}$$

# Piecewise-Defined Functions



# Piecewise-Defined Functions

When evaluating piecewise-defined functions, pay attention to the **domain** of each piece.

## Example 5

Evaluate each for

$$(a) \quad f(-3)$$

$$f(x) = \begin{cases} 4 - x^2 & \text{if } x < 1 \\ x - 3 & \text{if } x \geq 1 \end{cases}$$

## Example 5

Evaluate each for

$$(a) \quad f(-3) = -5$$

$$f(x) = \begin{cases} 4 - x^2 & \text{if } x < 1 \\ x - 3 & \text{if } x \geq 1 \end{cases}$$

## Example 5

Evaluate each for

$$f(x) = \begin{cases} 4 - x^2 & \text{if } x < 1 \\ x - 3 & \text{if } x \geq 1 \end{cases}$$

(a)  $f(-3) = -5$

(b)  $f(0)$

## Example 5

Evaluate each for

$$f(x) = \begin{cases} 4 - x^2 & \text{if } x < 1 \\ x - 3 & \text{if } x \geq 1 \end{cases}$$

(a)  $f(-3) = -5$

(b)  $f(0) = 4$

## Example 5

Evaluate each for

$$f(x) = \begin{cases} 4 - x^2 & \text{if } x < 1 \\ x - 3 & \text{if } x \geq 1 \end{cases}$$

(a)  $f(-3) = -5$

(b)  $f(0) = 4$

(c)  $f(2)$



## Example 5

Evaluate each for

$$f(x) = \begin{cases} 4 - x^2 & \text{if } x < 1 \\ x - 3 & \text{if } x \geq 1 \end{cases}$$

(a)  $f(-3) = -5$

(b)  $f(0) = 4$

(c)  $f(2) = -1$

## Example 5

Evaluate each for

$$f(x) = \begin{cases} 4 - x^2 & \text{if } x < 1 \\ x - 3 & \text{if } x \geq 1 \end{cases}$$

(a)  $f(-3) = -5$

(b)  $f(0) = 4$

(c)  $f(2) = -1$

(d)  $f\left(\frac{3}{2}\right)$

## Example 5

Evaluate each for

$$f(x) = \begin{cases} 4 - x^2 & \text{if } x < 1 \\ x - 3 & \text{if } x \geq 1 \end{cases}$$

(a)  $f(-3) = -5$

(b)  $f(0) = 4$

(c)  $f(2) = -1$

(d)  $f\left(\frac{3}{2}\right) = -\frac{3}{2}$