

# Logarithmic Functions

# Objectives

- 1 Calculate logarithmic values
- 2 Find the domain of logarithmic functions

# Logarithms as Inverse Exponents

The **inverse** of the exponential function  $f(x) = b^x$  is the **logarithmic function**.

# Logarithms as Inverse Exponents

The **inverse** of the exponential function  $f(x) = b^x$  is the **logarithmic function**.

It is denoted

$$f^{-1}(x) = \log_b x$$

# Special Bases

The **common logarithm** of a real number  $x$  is  $\log_{10} x$  and is usually written

$$\log x$$

.

# Special Bases

The **common logarithm** of a real number  $x$  is  $\log_{10} x$  and is usually written

$$\log x$$


.

The **natural logarithm** of a real number  $x$  is  $\log_e x$  and is usually written


$$\ln x$$

.

# Graphs



# Graphs





# Properties

- The domain of  $\log_b x$  is  $(0, \infty)$  and the range is  $(-\infty, \infty)$

# Properties

- The domain of  $\log_b x$  is  $(0, \infty)$  and the range is  $(-\infty, \infty)$
- $(1, 0)$  is on the graph of  $\log_b x$  and  $x = 0$  is a vertical asymptote.

# Properties

- The domain of  $\log_b x$  is  $(0, \infty)$  and the range is  $(-\infty, \infty)$
- $(1, 0)$  is on the graph of  $\log_b x$  and  $x = 0$  is a vertical asymptote.
- $\log_b x$  is one-to-one (has an inverse, passes HLT), continuous, and smooth.

# Properties

- The domain of  $\log_b x$  is  $(0, \infty)$  and the range is  $(-\infty, \infty)$
- $(1, 0)$  is on the graph of  $\log_b x$  and  $x = 0$  is a vertical asymptote.
- $\log_b x$  is one-to-one (has an inverse, passes HLT), continuous, and smooth.
- $b^a = c$  if and only if  $\log_b c = a$ .

# Properties

- The domain of  $\log_b x$  is  $(0, \infty)$  and the range is  $(-\infty, \infty)$
- $(1, 0)$  is on the graph of  $\log_b x$  and  $x = 0$  is a vertical asymptote.
- $\log_b x$  is one-to-one (has an inverse, passes HLT), continuous, and smooth.
- $b^a = c$  if and only if  $\log_b c = a$ .
  - In other words,  $\log_b c$  is the **exponent** you put on  $b$  to get  $c$ .

# Properties

- The domain of  $\log_b x$  is  $(0, \infty)$  and the range is  $(-\infty, \infty)$
- $(1, 0)$  is on the graph of  $\log_b x$  and  $x = 0$  is a vertical asymptote.
- $\log_b x$  is one-to-one (has an inverse, passes HLT), continuous, and smooth.
- $b^a = c$  if and only if  $\log_b c = a$ .
  - In other words,  $\log_b c$  is the **exponent** you put on  $b$  to get  $c$ .
- $\log_b b^x = x$  for all  $x$  and  $b^{\log_b x} = x$  for all  $x > 0$

# Properties

For  $f(x) = \log_b x$  when  $b > 1$ :

# Properties

For  $f(x) = \log_b x$  when  $b > 1$ :

- $f$  is always increasing.



# Properties

For  $f(x) = \log_b x$  when  $b > 1$ :

- $f$  is always increasing.
- $\lim_{x \rightarrow 0^+} f(x) = -\infty$  and  $\lim_{x \rightarrow \infty} f(x) = \infty$

# Properties

For  $f(x) = \log_b x$  when  $b > 1$ :

- $f$  is always increasing.
- $\lim_{x \rightarrow 0^+} f(x) = -\infty$  and  $\lim_{x \rightarrow \infty} f(x) = \infty$

For  $f(x) = \log_b x$  when  $0 < b < 1$ :

# Properties

For  $f(x) = \log_b x$  when  $b > 1$ :

- $f$  is always increasing.
- $\lim_{x \rightarrow 0^+} f(x) = -\infty$  and  $\lim_{x \rightarrow \infty} f(x) = \infty$

For  $f(x) = \log_b x$  when  $0 < b < 1$ :

- $f$  is always decreasing.

# Properties

For  $f(x) = \log_b x$  when  $b > 1$ :

- $f$  is always increasing.
- $\lim_{x \rightarrow 0^+} f(x) = -\infty$  and  $\lim_{x \rightarrow \infty} f(x) = \infty$

For  $f(x) = \log_b x$  when  $0 < b < 1$ :

- $f$  is always decreasing.
- $\lim_{x \rightarrow 0^+} f(x) = \infty$  and  $\lim_{x \rightarrow \infty} f(x) = -\infty$

## Example 1

Simplify each of the following.

(a)  $\log_3 81$

## Example 1

Simplify each of the following.

(a)  $\log_3 81$

$$3^{???} = 81$$

## Example 1

Simplify each of the following.

(a)  $\log_3 81$

$$3^{???} = 81$$

$$3^4 = 81$$

## Example 1

Simplify each of the following.

(a)  $\log_3 81$

$$3^{???} = 81$$

$$3^4 = 81$$

$$\log_3 81 = 4$$



## Example 1

$$(b) \quad \log_2 \left( \frac{1}{8} \right)$$

## Example 1

$$(b) \quad \log_2 \left( \frac{1}{8} \right)$$

$$2^{???} = \frac{1}{8}$$

## Example 1

$$(b) \quad \log_2 \left( \frac{1}{8} \right)$$

$$2^{???} = \frac{1}{8}$$

$$2^{-3} = \frac{1}{8}$$

## Example 1

$$(b) \quad \log_2 \left( \frac{1}{8} \right)$$

$$2^{???} = \frac{1}{8}$$

$$2^{-3} = \frac{1}{8}$$

$$\log_2 \left( \frac{1}{8} \right) = -3$$

## Example 1

(c)  $\log_{\sqrt{5}} 25$

## Example 1

$$(c) \quad \log_{\sqrt{5}} 25$$

$$\left(\sqrt{5}\right)^{???} = 25$$

## Example 1

$$(c) \quad \log_{\sqrt{5}} 25$$

$$\left(\sqrt{5}\right)^{???} = 25$$

$$\left(\sqrt{5}\right)^4 = 25$$

## Example 1

$$(c) \quad \log_{\sqrt{5}} 25$$

$$\left(\sqrt{5}\right)^{???} = 25$$

$$\left(\sqrt{5}\right)^4 = 25$$

$$\log_{\sqrt{5}} 25 = 4$$



## Example 1

$$(d) \quad \ln \left( \sqrt[3]{e^2} \right)$$

## Example 1

$$(d) \quad \ln \left( \sqrt[3]{e^2} \right) = \ln e^{2/3}$$

## Example 1

$$(d) \quad \ln \left( \sqrt[3]{e^2} \right) = \ln e^{2/3} = \log_e e^{2/3}$$

## Example 1

$$(d) \quad \ln \left( \sqrt[3]{e^2} \right) = \ln e^{2/3} = \log_e e^{2/3}$$

$$e^{???} = e^{2/3}$$

## Example 1

$$(d) \quad \ln \left( \sqrt[3]{e^2} \right) = \ln e^{2/3} = \log_e e^{2/3}$$

$$e^{???} = e^{2/3}$$

$$e^{2/3} = e^{2/3}$$

## Example 1

$$(d) \quad \ln \left( \sqrt[3]{e^2} \right) = \ln e^{2/3} = \log_e e^{2/3}$$

$$e^{???} = e^{2/3}$$

$$e^{2/3} = e^{2/3}$$

$$\ln \left( \sqrt[3]{e^2} \right) = \frac{2}{3}$$

# Example 1

(e)  $\log 0.001$

## Example 1

$$(e) \quad \log 0.001 = \log_{10} 0.001$$



## Example 1

$$(e) \quad \log 0.001 = \log_{10} 0.001 = \log_{10} 10^{-3}$$

## Example 1

$$(e) \quad \log 0.001 = \log_{10} 0.001 = \log_{10} 10^{-3}$$

$$10^{???} = 10^{-3}$$

## Example 1

$$(e) \quad \log 0.001 = \log_{10} 0.001 = \log_{10} 10^{-3}$$

$$10^{???} = 10^{-3}$$

$$10^{-3} = 10^{-3}$$

## Example 1

$$(e) \quad \log 0.001 = \log_{10} 0.001 = \log_{10} 10^{-3}$$

$$10^{???} = 10^{-3}$$

$$10^{-3} = 10^{-3}$$

$$\log 0.001 = -3$$

# Example 1

$$(f) \quad 2^{\log_2 8}$$

# Example 1

$$(f) \quad 2^{\log_2 8}$$

$$\log_2 8 = ???$$

# Example 1

$$(f) \quad 2^{\log_2 8}$$

$$\log_2 8 = ???$$

$$2^{???} = 8$$

# Example 1

$$(f) \quad 2^{\log_2 8}$$

$$\log_2 8 = ???$$

$$2^{???} = 8$$

$$2^3 = 8$$



# Example 1

$$(f) \quad 2^{\log_2 8}$$

$$\log_2 8 = ???$$

$$2^{???} = 8$$

$$2^3 = 8$$

$$\log_2 8 = 3$$

# Example 1

$$(f) \quad 2^{\log_2 8}$$

$$\log_2 8 = ???$$

$$2^{???} = 8$$

$$2^3 = 8$$

$$\log_2 8 = 3$$

$$2^{\log_2 8} = 2^3$$

# Example 1

$$(f) \quad 2^{\log_2 8}$$

$$\log_2 8 = ???$$

$$2^{???} = 8$$

$$2^3 = 8$$

$$\log_2 8 = 3$$

$$2^{\log_2 8} = 2^3$$

$$= 8$$

## Example 1

$$(g) \quad 117^{-\log_{117} 6}$$

## Example 1

(g)  $117^{-\log_{117} 6}$

$$117^{-\log_{117} 6} = \frac{1}{117^{\log_{117} 6}}$$

## Example 1

(g)  $117^{-\log_{117} 6}$

$$117^{-\log_{117} 6} = \frac{1}{117^{\log_{117} 6}}$$

$$= \frac{1}{6}$$

# Objectives

- 1 Calculate logarithmic values
- 2 Find the domain of logarithmic functions

# Domains of Logarithmic Functions

Up until now, the only domain restrictions we have had have been

- Denominator can't  $= 0$
- Can't take even root of a negative number



# Domains of Logarithmic Functions

Up until now, the only domain restrictions we have had have been

- Denominator can't  $= 0$
- Can't take even root of a negative number

For logarithms:

$$\log_b(\text{positive value})$$

# Domains of Logarithmic Functions

Up until now, the only domain restrictions we have had have been

- Denominator can't  $= 0$
- Can't take even root of a negative number

For logarithms:

$$\log_b (\text{positive value})$$

$$\log_b ( > 0 )$$

## Example 2

Find the domain of each. Write your answer in interval notation.

(a)  $f(x) = 2 \log(3 - x) - 1$

## Example 2

Find the domain of each. Write your answer in interval notation.

(a)  $f(x) = 2 \log(3 - x) - 1$

$$3 - x > 0$$

## Example 2

Find the domain of each. Write your answer in interval notation.

(a)  $f(x) = 2 \log(3 - x) - 1$

$$3 - x > 0$$

$$x < 3$$

## Example 2

Find the domain of each. Write your answer in interval notation.

(a)  $f(x) = 2 \log(3 - x) - 1$

$$3 - x > 0$$

$$x < 3$$

$$(-\infty, 3)$$

## Example 2

$$(b) \quad g(x) = -\frac{2}{3} \log_8 (x^2 + 6x - 7)$$

## Example 2

$$(b) \quad g(x) = -\frac{2}{3} \log_8 (x^2 + 6x - 7)$$

$$x^2 + 6x - 7 > 0$$



## Example 2

$$(b) \quad g(x) = -\frac{2}{3} \log_8 (x^2 + 6x - 7)$$

$$x^2 + 6x - 7 > 0$$

$$x^2 + 6x - 7 = 0$$

## Example 2

$$(b) \quad g(x) = -\frac{2}{3} \log_8 (x^2 + 6x - 7)$$

$$x^2 + 6x - 7 > 0$$

$$x^2 + 6x - 7 = 0$$

$$x = -7, 1$$

## Example 2

$$(b) \quad g(x) = -\frac{2}{3} \log_8 (x^2 + 6x - 7)$$

$$x^2 + 6x - 7 > 0$$

$$x^2 + 6x - 7 = 0$$

$$x = -7, 1$$



## Example 2

$$(b) \quad g(x) = -\frac{2}{3} \log_8 (x^2 + 6x - 7)$$

$$x^2 + 6x - 7 > 0$$

$$x^2 + 6x - 7 = 0$$

$$x = -7, 1$$




## Example 2

$$(b) \quad g(x) = -\frac{2}{3} \log_8 (x^2 + 6x - 7)$$

$$x^2 + 6x - 7 > 0$$

$$x^2 + 6x - 7 = 0$$

$$x = -7, 1$$




## Example 2

$$(b) \quad g(x) = -\frac{2}{3} \log_8 (x^2 + 6x - 7)$$

$$x^2 + 6x - 7 > 0$$

$$x^2 + 6x - 7 = 0$$

$$x = -7, 1$$



$$(\infty, -7) \cup (1, \infty)$$

## Example 2

$$(c) \quad h(x) = \ln \left( \frac{x}{x-1} \right)$$

## Example 2

$$(c) \quad h(x) = \ln \left( \frac{x}{x-1} \right)$$

$$\frac{x}{x-1} > 0$$



## Example 2

$$(c) \quad h(x) = \ln \left( \frac{x}{x-1} \right)$$

$$\frac{x}{x-1} > 0$$


Critical values:  $x = 0$  and  $x = 1$

## Example 2

$$(c) \quad h(x) = \ln \left( \frac{x}{x-1} \right)$$

$$\frac{x}{x-1} > 0$$

Critical values:  $x = 0$  and  $x = 1$




## Example 2

$$(c) \quad h(x) = \ln \left( \frac{x}{x-1} \right)$$

$$\frac{x}{x-1} > 0$$

Critical values:  $x = 0$  and  $x = 1$



## Example 2

$$(c) \quad h(x) = \ln \left( \frac{x}{x-1} \right)$$

$$\frac{x}{x-1} > 0$$

Critical values:  $x = 0$  and  $x = 1$



## Example 2

$$(c) \quad h(x) = \ln \left( \frac{x}{x-1} \right)$$

$$\frac{x}{x-1} > 0$$

Critical values:  $x = 0$  and  $x = 1$



$$(-\infty, 0) \cup (1, \infty)$$