

Quadratic Functions

Objectives

- 1 Determine the vertex, range, and intercepts of a quadratic function in standard form
- 2 Convert between standard and general form of quadratic functions

Quadratic Functions

A **quadratic function** is a function in the form


$$f(x) = ax^2 + bx + c$$

where a , b , and c are real numbers with $a \neq 0$.

The domain of a quadratic function is $(-\infty, \infty)$.

Graph of a Quadratic Function

For $f(x) = x^2$, the graph below is a **parabola**.



Graph of a Quadratic Function

The point $(0, 0)$ is called the **vertex** of the parabola and can be either a relative minimum or relative maximum point.

Graph of a Quadratic Function

The point $(0, 0)$ is called the **vertex** of the parabola and can be either a relative minimum or relative maximum point.

We can graph parabolas using transformations to the parent function $f(x) = x^2$.

Example 1


Graph the following functions starting with the graph of $f(x) = x^2$ and using transformations. Find the vertex, state the range, and find the x - and y -intercepts, if any exist.

(a) $g(x) = (x + 2)^2 - 3$

Example 1

Graph the following functions starting with the graph of $f(x) = x^2$ and using transformations. Find the vertex, state the range, and find the x - and y -intercepts, if any exist.


(a) $g(x) = (x + 2)^2 - 3$



Example 1

Graph the following functions starting with the graph of $f(x) = x^2$ and using transformations. Find the vertex, state the range, and find the x - and y -intercepts, if any exist.

(a) $g(x) = (x + 2)^2 - 3$




Shift left 2 units

Shift down 3 units

Example 1


(a) $g(x) = (x + 2)^2 - 3$



Example 1

(a) $g(x) = (x + 2)^2 - 3$

Vertex at $(-2, -3)$




Example 1

(a) $g(x) = (x + 2)^2 - 3$


Vertex at $(-2, -3)$

Range $[-3, \infty)$



Example 1

(a) $g(x) = (x + 2)^2 - 3$



Vertex at $(-2, -3)$


Range $[-3, \infty)$

x-intercepts:

$$(x + 2)^2 - 3 = 0$$

Example 1

(a) $g(x) = (x + 2)^2 - 3$



Vertex at $(-2, -3)$

Range $[-3, \infty)$


x-intercepts:

$$(x + 2)^2 - 3 = 0$$

$$(x + 2)^2 = 3$$

Example 1

(a) $g(x) = (x + 2)^2 - 3$



Vertex at $(-2, -3)$

Range $[-3, \infty)$

x-intercepts:


$$(x + 2)^2 - 3 = 0$$

$$(x + 2)^2 = 3$$

$$x + 2 = \pm\sqrt{3}$$

Example 1

(a) $g(x) = (x + 2)^2 - 3$



Vertex at $(-2, -3)$

Range $[-3, \infty)$

x-intercepts:

$$(x + 2)^2 - 3 = 0$$


$$(x + 2)^2 = 3$$

$$x + 2 = \pm\sqrt{3}$$

$$x = -2 \pm \sqrt{3}$$

Example 1

(a) $g(x) = (x + 2)^2 - 3$



Vertex at $(-2, -3)$

Range $[-3, \infty)$

x-intercepts:

$$(x + 2)^2 - 3 = 0$$

$$(x + 2)^2 = 3$$


$$x + 2 = \pm\sqrt{3}$$

$$x = -2 \pm \sqrt{3}$$

$$(-2 \pm \sqrt{3}, 0)$$


Example 1

(a) $g(x) = (x + 2)^2 - 3$



Example 1

(a) $g(x) = (x + 2)^2 - 3$




y -intercept at $(0, 1)$

Example 1

$$(b) \quad h(x) = -2(x - 3)^2 + 1$$


Example 1

(b) $h(x) = -2(x - 3)^2 + 1$



Example 1

(b) $h(x) = -2(x - 3)^2 + 1$



Reflect across x -axis


Vertical stretch by factor
of 2

Shift right 3 units

Shift up 1 unit

Example 1


(b) $h(x) = -2(x - 3)^2 + 1$



Example 1

(b) $h(x) = -2(x - 3)^2 + 1$

Vertex at $(3, 1)$




Example 1

(b) $h(x) = -2(x - 3)^2 + 1$


Vertex at $(3, 1)$

Range $(-\infty, 1]$



Example 1

(b) $h(x) = -2(x - 3)^2 + 1$



Vertex at (3, 1)


Range $(-\infty, 1]$

x-intercepts:

$$-2(x - 3)^2 + 1 = 0$$

Example 1

(b) $h(x) = -2(x - 3)^2 + 1$



Vertex at (3, 1)

Range $(-\infty, 1]$


x-intercepts:

$$-2(x - 3)^2 + 1 = 0$$

$$-2(x - 3)^2 = -1$$

Example 1

(b) $h(x) = -2(x - 3)^2 + 1$



Vertex at (3, 1)

Range $(-\infty, 1]$

x-intercepts:


$$-2(x - 3)^2 + 1 = 0$$

$$-2(x - 3)^2 = -1$$

$$(x - 3)^2 = \frac{1}{2}$$

Example 1

(b) $h(x) = -2(x - 3)^2 + 1$



Vertex at (3, 1)

Range $(-\infty, 1]$

x-intercepts:

$$-2(x - 3)^2 + 1 = 0$$


$$-2(x - 3)^2 = -1$$

$$(x - 3)^2 = \frac{1}{2}$$

$$x - 3 = \pm\sqrt{\frac{1}{2}}$$

Example 1


(b) $h(x) = -2(x - 3)^2 + 1$



$$x - 3 = \pm \sqrt{\frac{1}{2}}$$

Example 1

(b) $h(x) = -2(x - 3)^2 + 1$




$$x - 3 = \pm \sqrt{\frac{1}{2}}$$

$$x - 3 = \pm \frac{\sqrt{2}}{2}$$

Example 1

(b) $h(x) = -2(x - 3)^2 + 1$




$$x - 3 = \pm \sqrt{\frac{1}{2}}$$

$$x - 3 = \pm \frac{\sqrt{2}}{2}$$

$$x = 3 \pm \frac{\sqrt{2}}{2}$$

Example 1

(b) $h(x) = -2(x - 3)^2 + 1$



$$x - 3 = \pm \sqrt{\frac{1}{2}}$$


$$x - 3 = \pm \frac{\sqrt{2}}{2}$$

$$x = 3 \pm \frac{\sqrt{2}}{2}$$

$$\left(3 \pm \frac{\sqrt{2}}{2}, 0 \right)$$

Example 1

(b) $h(x) = -2(x - 3)^2 + 1$



$$x - 3 = \pm \sqrt{\frac{1}{2}}$$

$$x - 3 = \pm \frac{\sqrt{2}}{2}$$

$$x = 3 \pm \frac{\sqrt{2}}{2}$$

$$\left(3 \pm \frac{\sqrt{2}}{2}, 0 \right)$$

y-intercept: $(0, -17)$

Objectives

- 1 Determine the vertex, range, and intercepts of a quadratic function in standard form
- 2 Convert between standard and general form of quadratic functions

Standard and General Form of Quadratic Functions

If f is a quadratic function:

- The **general form** is $f(x) = ax^2 + bx + c$

Standard and General Form of Quadratic Functions

If f is a quadratic function:

- The **general form** is $f(x) = ax^2 + bx + c$
 - a , b , and c are real numbers

Standard and General Form of Quadratic Functions

If f is a quadratic function:

- The **general form** is $f(x) = ax^2 + bx + c$
 - a , b , and c are real numbers
 - $a \neq 0$

Standard and General Form of Quadratic Functions

If f is a quadratic function:

- The **general form** is $f(x) = ax^2 + bx + c$
 - a , b , and c are real numbers
 - $a \neq 0$
- The **standard form** is $f(x) = a(x - h)^2 + k$

Standard and General Form of Quadratic Functions

If f is a quadratic function:

- The **general form** is $f(x) = ax^2 + bx + c$
 - a , b , and c are real numbers
 - $a \neq 0$
- The **standard form** is $f(x) = a(x - h)^2 + k$
 - Vertex is (h, k)

Standard and General Form of Quadratic Functions

If f is a quadratic function:

- The **general form** is $f(x) = ax^2 + bx + c$
 - a , b , and c are real numbers
 - $a \neq 0$
- The **standard form** is $f(x) = a(x - h)^2 + k$
 - Vertex is (h, k)
 - $a \neq 0$

Standard and General Form of Quadratic Functions

If f is a quadratic function:

- The **general form** is $f(x) = ax^2 + bx + c$
 - a , b , and c are real numbers
 - $a \neq 0$
- The **standard form** is $f(x) = a(x - h)^2 + k$
 - Vertex is (h, k)
 - $a \neq 0$
 - a , h , and k are real numbers

Converting From General to Standard Form

To convert from general form $f(x) = ax^2 + bx + c$ to standard form $f(x) = a(x - h)^2 + k$

Converting From General to Standard Form

To convert from general form $f(x) = ax^2 + bx + c$ to standard form $f(x) = a(x - h)^2 + k$

① Find the vertex:

- x-coordinate: $\frac{-b}{2a}$
- y-coordinate: Evaluate function at x-coordinate
- Or use graphing technology

② Use the same value of a

Example 2

Convert each to standard form.

(a) $f(x) = x^2 - 4x + 3$

Example 2

Convert each to standard form.

(a) $f(x) = x^2 - 4x + 3$

Vertex:

Example 2

Convert each to standard form.

(a) $f(x) = x^2 - 4x + 3$

Vertex:

$$x = \frac{-(-4)}{2(1)}$$

Example 2

Convert each to standard form.

(a) $f(x) = x^2 - 4x + 3$

Vertex:

$$x = \frac{-(-4)}{2(1)}$$

$$x = 2$$

Example 2

Convert each to standard form.

(a) $f(x) = x^2 - 4x + 3$

Vertex:

$$x = \frac{-(-4)}{2(1)}$$

$$x = 2$$

$$y = 2^2 - 4(2) + 3$$

Example 2

Convert each to standard form.

$$(a) \quad f(x) = x^2 - 4x + 3$$

Vertex:

$$x = \frac{-(-4)}{2(1)}$$

$$x = 2$$

$$y = 2^2 - 4(2) + 3$$

$$y = -1$$

Example 2

(a) $f(x) = x^2 - 4x + 3$

Vertex: $(2, -1)$

Example 2

$$(a) \quad f(x) = x^2 - 4x + 3$$

Vertex: $(2, -1)$

$$a = 1$$

Example 2

$$(a) \quad f(x) = x^2 - 4x + 3$$

Vertex: $(2, -1)$

$$a = 1$$

$$f(x) = (x - 2)^2 - 1$$

Example 2

Convert each to standard form.

(b) $g(x) = 6 - x - x^2$

Example 2

Convert each to standard form.

(b) $g(x) = 6 - x - x^2$


$$g(x) = -x^2 - x + 6$$

Example 2

Convert each to standard form.

(b) $g(x) = 6 - x - x^2$

$$g(x) = -x^2 - x + 6$$




Example 2

Convert each to standard form.

(b) $g(x) = 6 - x - x^2$

$$g(x) = -x^2 - x + 6$$




Vertex: $\left(-\frac{1}{2}, \frac{25}{4}\right)$

Example 2

Convert each to standard form.

(b) $g(x) = 6 - x - x^2$

$$g(x) = -x^2 - x + 6$$



Vertex: $\left(-\frac{1}{2}, \frac{25}{4}\right)$


$$a = -1$$

Example 2

Convert each to standard form.

(b) $g(x) = 6 - x - x^2$

$$g(x) = -x^2 - x + 6$$




$$\text{Vertex: } \left(-\frac{1}{2}, \frac{25}{4}\right)$$

$$a = -1$$

$$g(x) = -\left(x + \frac{1}{2}\right)^2 + \frac{25}{4}$$


Axis of Symmetry

The graphs of the parabolas have a line of symmetry called the **axis of symmetry** that is a vertical line through the x-coordinate of the vertex:



Axis of Symmetry

The graphs of the parabolas have a line of symmetry called the **axis of symmetry** that is a vertical line through the x-coordinate of the vertex:



$$x = -\frac{1}{2}$$

Converting From Standard to General Form

To convert from

$$f(x) = a(x - h)^2 + k$$

form to

$$f(x) = ax^2 + bx + c$$

form, just **do the math**.

Example 3

Convert $g(x) = (x + 2)^2 - 3$ to general form.

Example 3

Convert $g(x) = (x + 2)^2 - 3$ to general form.

$$(x + 2)^2 - 3 = x^2 + 4x + 4 - 3$$

Example 3

Convert $g(x) = (x + 2)^2 - 3$ to general form.

$$(x + 2)^2 - 3 = x^2 + 4x + 4 - 3$$

$$= x^2 + 4x + 1$$