

# Complex Numbers

# Objectives

- 1 Perform arithmetic operations with complex numbers
- 2 Solve quadratic equations with complex solutions

Complex numbers arose from the need to solve equations such as

$$x^2 + 1 = 0$$

.

Complex numbers arose from the need to solve equations such as

$$x^2 + 1 = 0$$

The solution to the above problem is

$$x = \pm\sqrt{-1}$$

Complex numbers arose from the need to solve equations such as

$$x^2 + 1 = 0$$

The solution to the above problem is

$$x = \pm\sqrt{-1}$$

$\sqrt{-1}$  is the **imaginary unit**  $i$ .

# Properties of $i$

- $i^2 = -1$

# Properties of $i$

- $i^2 = -1$
- If  $c$  is a real number with  $c \geq 0$ , then  $\sqrt{-c} = i\sqrt{c}$

# Complex Numbers

A **complex number** is a number in the form  $a + bi$ , where  $a$  and  $b$  are real numbers and  $i$  is the imaginary unit.



# Complex Numbers

A **complex number** is a number in the form  $a + bi$ , where  $a$  and  $b$  are real numbers and  $i$  is the imaginary unit.

A pure real number can be written as  $a + 0i$  and a pure imaginary number can be written as  $0 + bi$ .

# Complex Numbers

A **complex number** is a number in the form  $a + bi$ , where  $a$  and  $b$  are real numbers and  $i$  is the imaginary unit.

A pure real number can be written as  $a + 0i$  and a pure imaginary number can be written as  $0 + bi$ .

Adding, subtracting, and multiplying complex numbers is a lot like that of real numbers.

# Complex Numbers

A **complex number** is a number in the form  $a + bi$ , where  $a$  and  $b$  are real numbers and  $i$  is the imaginary unit.

A pure real number can be written as  $a + 0i$  and a pure imaginary number can be written as  $0 + bi$ .

Adding, subtracting, and multiplying complex numbers is a lot like that of real numbers.

However, keep in mind that  $i^2 = -1$ .

## Example 1

Perform each indicated operation. Write your answers in  $a + bi$  form.

(a)  $(1 - 2i) - (3 + 4i)$

## Example 1

Perform each indicated operation. Write your answers in  $a + bi$  form.

$$(a) \quad (1 - 2i) - (3 + 4i)$$

$$= 1 - 2i - 3 - 4i$$

## Example 1

Perform each indicated operation. Write your answers in  $a + bi$  form.

$$(a) \quad (1 - 2i) - (3 + 4i)$$

$$= 1 - 2i - 3 - 4i$$

$$= -2 - 6i$$

## Example 1

$$(b) \quad (1 - 2i)(3 + 4i)$$

## Example 1

$$(b) \quad (1 - 2i)(3 + 4i)$$

$$= 3 + 4i - 6i - 8i^2$$



## Example 1

$$(b) \quad (1 - 2i)(3 + 4i)$$

$$= 3 + 4i - 6i - 8i^2$$

$$= 3 - 2i - 8(-1)$$

## Example 1

$$(b) \quad (1 - 2i)(3 + 4i)$$

$$= 3 + 4i - 6i - 8i^2$$

$$= 3 - 2i - 8(-1)$$

$$= 3 - 2i + 8$$

## Example 1

$$(b) \quad (1 - 2i)(3 + 4i)$$

$$= 3 + 4i - 6i - 8i^2$$

$$= 3 - 2i - 8(-1)$$

$$= 3 - 2i + 8$$

$$= 11 - 2i$$

## Example 1

$$(c) \quad \sqrt{-3}\sqrt{-12}$$

## Example 1

$$(c) \quad \sqrt{-3}\sqrt{-12}$$

$$= i\sqrt{3} \cdot 2i\sqrt{3}$$

## Example 1

$$(c) \quad \sqrt{-3}\sqrt{-12}$$

$$= i\sqrt{3} \cdot 2i\sqrt{3}$$

$$= 2i^2\sqrt{9}$$

## Example 1

$$(c) \quad \sqrt{-3}\sqrt{-12}$$

$$= i\sqrt{3} \cdot 2i\sqrt{3}$$

$$= 2i^2\sqrt{9}$$

$$= -2(3)$$

## Example 1

$$(c) \quad \sqrt{-3}\sqrt{-12}$$

$$= i\sqrt{3} \cdot 2i\sqrt{3}$$

$$= 2i^2\sqrt{9}$$

$$= -2(3)$$

$$= -6$$



## Example 1

$$(d) \quad \sqrt{(-3)(-12)}$$

## Example 1

$$(d) \quad \sqrt{(-3)(-12)}$$

$$= \sqrt{36}$$

## Example 1

$$(d) \quad \sqrt{(-3)(-12)}$$

$$= \sqrt{36}$$

$$= 6$$

## Example 1

$$(e) \quad (x - (1 + 2i))(x - (1 - 2i))$$

## Example 1

$$\begin{aligned} \text{(e)} \quad & (x - (1 + 2i))(x - (1 - 2i)) \\ &= x^2 - x(1 - 2i) - x(1 + 2i) + (1 + 2i)(1 - 2i) \end{aligned}$$

## Example 1

$$\begin{aligned} \text{(e)} \quad & (x - (1 + 2i))(x - (1 - 2i)) \\ &= x^2 - x(1 - 2i) - x(1 + 2i) + (1 + 2i)(1 - 2i) \\ &= x^2 - x + 2ix - x - 2ix + 1 - 2i + 2i - 4i^2 \end{aligned}$$

## Example 1

$$\begin{aligned} \text{(e)} \quad & (x - (1 + 2i))(x - (1 - 2i)) \\ &= x^2 - x(1 - 2i) - x(1 + 2i) + (1 + 2i)(1 - 2i) \\ &= x^2 - x + 2ix - x - 2ix + 1 - 2i + 2i - 4i^2 \\ &= x^2 - 2x + 1 - 4(-1) \end{aligned}$$

## Example 1

$$\begin{aligned} \text{(e)} \quad & (x - (1 + 2i))(x - (1 - 2i)) \\ &= x^2 - x(1 - 2i) - x(1 + 2i) + (1 + 2i)(1 - 2i) \\ &= x^2 - x + 2ix - x - 2ix + 1 - 2i + 2i - 4i^2 \\ &= x^2 - 2x + 1 - 4(-1) \\ &= x^2 - 2x + 5 \end{aligned}$$



# Complex Conjugates

If  $z = a + bi$  is a complex number, then the **conjugate** of  $z$  (denoted  $\bar{z}$ ) is

$$\bar{z} = a - bi$$

# Complex Conjugates

If  $z = a + bi$  is a complex number, then the **conjugate** of  $z$  (denoted  $\bar{z}$ ) is

$$\bar{z} = a - bi$$

Likewise, if  $z = a - bi$ ,

$$\bar{z} = a + bi$$

# Complex Conjugates

If  $z = a + bi$  is a complex number, then the **conjugate** of  $z$  (denoted  $\bar{z}$ ) is

$$\bar{z} = a - bi$$

Likewise, if  $z = a - bi$ ,

$$\bar{z} = a + bi$$

Complex conjugates are used to divide complex numbers and find complex solutions to equations.

## Example 2

Write your answer in  $a + bi$  form for

$$\frac{1 - 2i}{3 - 4i}$$

## Example 2

Write your answer in  $a + bi$  form for

$$\frac{1 - 2i}{3 - 4i}$$

$$= \frac{1 - 2i}{3 - 4i} \left( \frac{3 + 4i}{3 + 4i} \right)$$

## Example 2

Write your answer in  $a + bi$  form for

$$\frac{1 - 2i}{3 - 4i}$$

$$= \frac{1 - 2i}{3 - 4i} \left( \frac{3 + 4i}{3 + 4i} \right)$$

$$= \frac{3 + 4i - 6i - 8i^2}{9 - 12i + 12i - 16i^2}$$

## Example 2

Write your answer in  $a + bi$  form for

$$\frac{1 - 2i}{3 - 4i}$$

$$= \frac{1 - 2i}{3 - 4i} \left( \frac{3 + 4i}{3 + 4i} \right)$$

$$= \frac{3 + 4i - 6i - 8i^2}{9 - 12i + 12i - 16i^2}$$

$$= \frac{11 - 2i}{25}$$

## Example 2

Write your answer in  $a + bi$  form for

$$\frac{1 - 2i}{3 - 4i}$$

$$= \frac{1 - 2i}{3 - 4i} \left( \frac{3 + 4i}{3 + 4i} \right)$$

$$= \frac{3 + 4i - 6i - 8i^2}{9 - 12i + 12i - 16i^2}$$

$$= \frac{11 - 2i}{25}$$

$$= \frac{11}{25} - \frac{2}{25}i$$



# Properties of Complex Conjugates

Let  $z$  and  $w$  be complex numbers.

- $\overline{\overline{z}} = z$
- $\overline{z + w} = \overline{z} + \overline{w}$
- $\overline{z \cdot w} = \overline{z} \cdot \overline{w}$
- $(\overline{z})^n = \overline{z^n}$  for any natural number  $n$
- $z$  is a real number if and only if  $\overline{z} = z$

# Objectives

- 1 Perform arithmetic operations with complex numbers
- 2 Solve quadratic equations with complex solutions

# Quadratic Equations with Complex Solutions

In the quadratic formula

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$


the discriminant

$$b^2 - 4ac$$

tells us what type of solutions we will have.


# The Discriminant

$b^2 - 4ac > 0$  gives us 2 unique real solutions.




# The Discriminant

$b^2 - 4ac = 0$  gives us 1 unique real solution (a double root).



# The Discriminant

$b^2 - 4ac < 0$  gives us 2 complex solutions that are *conjugates*.



## Example 3

Solve  $x^2 - 2x + 5 = 0$ . Exact answers only.

## Example 3

Solve  $x^2 - 2x + 5 = 0$ . Exact answers only.

$$x = \frac{-(-2) \pm \sqrt{(-2)^2 - 4(1)(5)}}{2(1)}$$



## Example 3

Solve  $x^2 - 2x + 5 = 0$ . Exact answers only.

$$x = \frac{-(-2) \pm \sqrt{(-2)^2 - 4(1)(5)}}{2(1)}$$

$$x = \frac{2 \pm \sqrt{-16}}{2}$$

## Example 3

Solve  $x^2 - 2x + 5 = 0$ . Exact answers only.

$$x = \frac{-(-2) \pm \sqrt{(-2)^2 - 4(1)(5)}}{2(1)}$$

$$x = \frac{2 \pm \sqrt{-16}}{2}$$

$$x = \frac{2 \pm 4i}{2}$$

## Example 3

Solve  $x^2 - 2x + 5 = 0$ . Exact answers only.

$$x = \frac{-(-2) \pm \sqrt{(-2)^2 - 4(1)(5)}}{2(1)}$$

$$x = \frac{2 \pm \sqrt{-16}}{2}$$

$$x = \frac{2 \pm 4i}{2}$$

$$x = 1 \pm 2i$$