

# Exponential Functions

# Objectives

- 1 Use exponential functions to solve problems

# Intro


An **exponential function** is a function in which each successive output value is obtained by **multiplying** the previous one by a constant value.

An **exponential function** is a function in which each successive output value is obtained by **multiplying** the previous one by a constant value.

One example of an exponential function is the **doubling function**

$$f(x) = 2^x$$

# Doubling Function



$x$	$f(x)$
-3	$\frac{1}{8}$
-2	$\frac{1}{4}$
-1	$\frac{1}{2}$
0	1
1	2
2	4
3	8

# End Behavior of the Doubling Function

As the values of  $x \rightarrow -\infty$ ,

$$2^{\text{very big negative number}} \rightarrow 0$$

# End Behavior of the Doubling Function

As the values of  $x \rightarrow -\infty$ ,

$$2^{\text{very big negative number}} \rightarrow 0$$

For instance,

$$2^{-50} \approx 0.00000000000000008882$$

# End Behavior of the Doubling Function

As the values of  $x \rightarrow -\infty$ ,

$$2^{\text{very big negative number}} \rightarrow 0$$

For instance,

$$2^{-50} \approx 0.00000000000000008882$$

As the values of  $x \rightarrow \infty$ ,

$$2^{\text{very big positive number}} \rightarrow \infty$$



# End Behavior of the Doubling Function

As the values of  $x \rightarrow -\infty$ ,

$$2^{\text{very big negative number}} \rightarrow 0$$

For instance,

$$2^{-50} \approx 0.00000000000000008882$$

As the values of  $x \rightarrow \infty$ ,

$$2^{\text{very big positive number}} \rightarrow \infty$$

For instance,



$$2^{50} = 1,125,899,906,842,624$$

# Exponential Functions

A function in the form  $f(x) = b^x$  where  $b$  is a fixed real number,  $b > 0$ ,  $b \neq 1$  is called an **exponential function of base  $b$**

# Exponential Functions

A function in the form  $f(x) = b^x$  where  $b$  is a fixed real number,  $b > 0$ ,  $b \neq 1$  is called an **exponential function of base  $b$**



# Properties of Exponential Functions

For  $f(x) = b^x$ :

- Domain is  $(-\infty, \infty)$  and the range is  $[0, \infty)$

# Properties of Exponential Functions

For  $f(x) = b^x$ :

- Domain is  $(-\infty, \infty)$  and the range is  $[0, \infty)$
- $(0, 1)$  is on the graph of  $f$  and  $y = 0$  is a horizontal asymptote.

# Properties of Exponential Functions

For  $f(x) = b^x$ :

- Domain is  $(-\infty, \infty)$  and the range is  $[0, \infty)$
- $(0, 1)$  is on the graph of  $f$  and  $y = 0$  is a horizontal asymptote.
- $f$  is one-to-one (has an inverse), continuous, and smooth.

# Properties of Exponential Functions

For  $f(x) = b^x$  when  $b > 1$ :

- $f$  is always increasing

# Properties of Exponential Functions

For  $f(x) = b^x$  when  $b > 1$ :


- $f$  is always increasing
- $\lim_{x \rightarrow -\infty} = 0$  and  $\lim_{x \rightarrow \infty} = \infty$



# Properties of Exponential Functions

For  $f(x) = b^x$  when  $b > 1$ :

- $f$  is always increasing
- $\lim_{x \rightarrow -\infty} = 0$  and  $\lim_{x \rightarrow \infty} = \infty$
- The graph of  $f$  resembles



# Properties of Exponential Functions

For  $f(x) = b^x$  when  $0 < b < 1$ :

- $f$  is always decreasing

# Properties of Exponential Functions


For  $f(x) = b^x$  when  $0 < b < 1$ :

- $f$  is always decreasing
- $\lim_{x \rightarrow -\infty} = \infty$  and  $\lim_{x \rightarrow \infty} = 0$

# Properties of Exponential Functions

For  $f(x) = b^x$  when  $0 < b < 1$ :

- $f$  is always decreasing
- $\lim_{x \rightarrow -\infty} = \infty$  and  $\lim_{x \rightarrow \infty} = 0$
- The graph of  $f$  resembles



# Special Bases

Of all possible bases for exponential functions, the 2 that occur most are base 10 (**common base**) and irrational base  $e \approx 2.718$  (**natural base**).

$$e = \lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^x$$

## Example 1

The value of a car can be modeled by  $V(x) = 25 \left(\frac{4}{5}\right)^x$ , where  $x \geq 0$  is the age of the car in years and  $V(x)$  is the value in thousands of dollars.

- (a) Find and interpret  $V(0)$

## Example 1

The value of a car can be modeled by  $V(x) = 25 \left(\frac{4}{5}\right)^x$ , where  $x \geq 0$  is the age of the car in years and  $V(x)$  is the value in thousands of dollars.

(a) Find and interpret  $V(0)$

$V(0)$  is the value of the car (in thousands of dollars) when it's brand new.

## Example 1

The value of a car can be modeled by  $V(x) = 25 \left(\frac{4}{5}\right)^x$ , where  $x \geq 0$  is the age of the car in years and  $V(x)$  is the value in thousands of dollars.

(a) Find and interpret  $V(0)$

$V(0)$  is the value of the car (in thousands of dollars) when it's brand new.

$$V(0) = 25 \left(\frac{4}{5}\right)^0$$



## Example 1

The value of a car can be modeled by  $V(x) = 25 \left(\frac{4}{5}\right)^x$ , where  $x \geq 0$  is the age of the car in years and  $V(x)$  is the value in thousands of dollars.

(a) Find and interpret  $V(0)$

$V(0)$  is the value of the car (in thousands of dollars) when it's brand new.

$$\begin{aligned} V(0) &= 25 \left(\frac{4}{5}\right)^0 \\ &= 25 \end{aligned}$$

## Example 1

The value of a car can be modeled by  $V(x) = 25 \left(\frac{4}{5}\right)^x$ , where  $x \geq 0$  is the age of the car in years and  $V(x)$  is the value in thousands of dollars.

(a) Find and interpret  $V(0)$

$V(0)$  is the value of the car (in thousands of dollars) when it's brand new.

$$\begin{aligned} V(0) &= 25 \left(\frac{4}{5}\right)^0 \\ &= 25 \end{aligned}$$

Brand new, the car is valued at \$25,000.

## Example 1

(b) Find the parent function and describe the function

$$V(x) = 25 \left( \frac{4}{5} \right)^x \text{ using transformations.}$$

## Example 1

(b) Find the parent function and describe the function

$$V(x) = 25 \left(\frac{4}{5}\right)^x \text{ using transformations.}$$

$$\text{Parent function is } f(x) = \left(\frac{4}{5}\right)^x$$

## Example 1

(b) Find the parent function and describe the function

$$V(x) = 25 \left(\frac{4}{5}\right)^x \text{ using transformations.}$$

$$\text{Parent function is } f(x) = \left(\frac{4}{5}\right)^x$$

$$V(x) = 25 \left(\frac{4}{5}\right)^x \text{ is a vertical stretch by a factor of 25.}$$

## Example 1

- (c) Find and interpret the horizontal asymptote of the graph of  $V(x)$ .

## Example 1

(c) Find and interpret the horizontal asymptote of the graph of  $V(x)$ .

Horizontal asymptote is  $y = 0$ .

## Example 1

(c) Find and interpret the horizontal asymptote of the graph of  $V(x)$ .

Horizontal asymptote is  $y = 0$ .

Over time, the value of the car will approach 0.



## Example 2

According to Newton's Law of Cooling, the temperature of coffee  $T$  (in degrees Fahrenheit)  $t$  minutes after it is served can be modeled by

$$T(t) = 70 + 90e^{-0.1t}$$

- (a) Find and interpret  $T(0)$ .

## Example 2

According to Newton's Law of Cooling, the temperature of coffee  $T$  (in degrees Fahrenheit)  $t$  minutes after it is served can be modeled by

$$T(t) = 70 + 90e^{-0.1t}$$

(a) Find and interpret  $T(0)$ .

$$T(0) = 70 + 90e^{-0.1(0)}$$

## Example 2

According to Newton's Law of Cooling, the temperature of coffee  $T$  (in degrees Fahrenheit)  $t$  minutes after it is served can be modeled by

$$T(t) = 70 + 90e^{-0.1t}$$

(a) Find and interpret  $T(0)$ .

$$\begin{aligned} T(0) &= 70 + 90e^{-0.1(0)} \\ &= 160 \end{aligned}$$

## Example 2

According to Newton's Law of Cooling, the temperature of coffee  $T$  (in degrees Fahrenheit)  $t$  minutes after it is served can be modeled by

$$T(t) = 70 + 90e^{-0.1t}$$

(a) Find and interpret  $T(0)$ .

$$\begin{aligned} T(0) &= 70 + 90e^{-0.1(0)} \\ &= 160 \end{aligned}$$

The coffee was served at  $160^{\circ}\text{F}$ .

## Example 2

$$T(t) = 70 + 90e^{-0.1t}$$

(b) Find the parent function and describe the function  $T(t)$  using transformations.

## Example 2

$$T(t) = 70 + 90e^{-0.1t}$$

(b) Find the parent function and describe the function  $T(t)$  using transformations.

Parent function is  $f(t) = e^t$

## Example 2

$$T(t) = 70 + 90e^{-0.1t}$$

(b) Find the parent function and describe the function  $T(t)$  using transformations.

Parent function is  $f(t) = e^t$

$T(t)$  is the parent function with the following transformations:

- Reflection across  $y$ -axis (multiplying  $t$  by  $-1$ )

## Example 2

$$T(t) = 70 + 90e^{-0.1t}$$

(b) Find the parent function and describe the function  $T(t)$  using transformations.

Parent function is  $f(t) = e^t$

$T(t)$  is the parent function with the following transformations:

- Reflection across  $y$ -axis (multiplying  $t$  by  $-1$ )
- Horizontal stretch by factor of 10 (multiplying  $t$  by 0.1)



## Example 2

$$T(t) = 70 + 90e^{-0.1t}$$

(b) Find the parent function and describe the function  $T(t)$  using transformations.

Parent function is  $f(t) = e^t$

$T(t)$  is the parent function with the following transformations:

- Reflection across  $y$ -axis (multiplying  $t$  by  $-1$ )
- Horizontal stretch by factor of 10 (multiplying  $t$  by 0.1)
- Vertical stretch by factor of 90

## Example 2

$$T(t) = 70 + 90e^{-0.1t}$$

(b) Find the parent function and describe the function  $T(t)$  using transformations.

Parent function is  $f(t) = e^t$

$T(t)$  is the parent function with the following transformations:

- Reflection across  $y$ -axis (multiplying  $t$  by  $-1$ )
- Horizontal stretch by factor of 10 (multiplying  $t$  by 0.1)
- Vertical stretch by factor of 90
- Shift up 70 degrees Fahrenheit

## Example 2

- (c) Find and interpret the horizontal asymptote of the graph.

## Example 2

(c) Find and interpret the horizontal asymptote of the graph.

Horizontal asymptote = 70

## Example 2

(c) Find and interpret the horizontal asymptote of the graph.

Horizontal asymptote = 70

Over time, the coffee will cool to a temperature of  $70^{\circ}\text{F}$ .