

Factoring Techniques

Objectives

- 1 Find the greatest common factor of a quadratic expression.
- 2 Factor trinomials in the form $x^2 + bx + c$
- 3 Factor trinomials in the form $ax^2 + bx + c$
- 4 Use “reverse engineering” to factor quadratic expressions

Greatest Common Factor

The **greatest common factor**, or GCF, of an expression is the largest term that divides into each term of that expression.

Greatest Common Factor

The **greatest common factor**, or GCF, of an expression is the largest term that divides into each term of that expression.

To find the greatest common factor of an expression:

Greatest Common Factor

The **greatest common factor**, or GCF, of an expression is the largest term that divides into each term of that expression.

To find the greatest common factor of an expression:

- Find the greatest common factor of the coefficients.

Greatest Common Factor

The **greatest common factor**, or GCF, of an expression is the largest term that divides into each term of that expression.

To find the greatest common factor of an expression:

- Find the greatest common factor of the coefficients.
- Select the lowest power of each unique variable.

Greatest Common Factor

The **greatest common factor**, or GCF, of an expression is the largest term that divides into each term of that expression.

To find the greatest common factor of an expression:

- Find the greatest common factor of the coefficients.
- Select the lowest power of each unique variable.

Then divide each term by that GCF.

Example 1

Factor the GCF from each.

(a) $16x^5 + 72x^3 - 36x^2$

Example 1

Factor the GCF from each.

(a) $16x^5 + 72x^3 - 36x^2$

GCF of 16, 72, and 36 is 4

Example 1

Factor the GCF from each.

(a) $16x^5 + 72x^3 - 36x^2$

GCF of 16, 72, and 36 is 4

Lowest power of x is 2, so the GCF of the variables is x^2

Example 1

Factor the GCF from each.

(a) $16x^5 + 72x^3 - 36x^2$

GCF of 16, 72, and 36 is 4

Lowest power of x is 2, so the GCF of the variables is x^2

GCF is $4x^2$

Example 1

Factor the GCF from each.

$$(a) \quad 16x^5 + 72x^3 - 36x^2$$

GCF of 16, 72, and 36 is 4

Lowest power of x is 2, so the GCF of the variables is x^2

GCF is $4x^2$

$$4x^2 (4x^3 + 18x - 9)$$

Example 1

$$(b) \quad 5a^4 + 70a^2 - 25a$$

Example 1

$$(b) \quad 5a^4 + 70a^2 - 25a$$

GCF of 5, 70, and 25 is 5

Example 1

$$(b) \quad 5a^4 + 70a^2 - 25a$$

GCF of 5, 70, and 25 is 5

Lowest power of a is 1, so the GCF of the variables is a

Example 1

$$(b) \quad 5a^4 + 70a^2 - 25a$$

GCF of 5, 70, and 25 is 5

Lowest power of a is 1, so the GCF of the variables is a

GCF is $5a$

Example 1

$$(b) \quad 5a^4 + 70a^2 - 25a$$

GCF of 5, 70, and 25 is 5

Lowest power of a is 1, so the GCF of the variables is a

GCF is $5a$

$$5a(a^3 + 14a - 5)$$

Objectives

- 1 Find the greatest common factor of a quadratic expression.
- 2 Factor trinomials in the form $x^2 + bx + c$
- 3 Factor trinomials in the form $ax^2 + bx + c$
- 4 Use “reverse engineering” to factor quadratic expressions

$$x^2 + bx + c$$

When factoring trinomials in the form $x^2 + bx + c$, we are looking to find 2 numbers that **multiply** to make c and **add** to make b .

$$x^2 + bx + c$$

When factoring trinomials in the form $x^2 + bx + c$, we are looking to find 2 numbers that **multiply** to make c and **add** to make b .

We can then write the factorization in the form

$$(x + m)(x + n)$$

where $mn = c$ and $m + n = b$

Example 2

Factor each completely.

(a) $x^2 + 15x + 36$

Example 2

Factor each completely.

(a) $x^2 + 15x + 36$

Product of the two numbers is 36.

Example 2

Factor each completely.

(a) $x^2 + 15x + 36$

Product of the two numbers is 36.

Sum of the two numbers is 15.

Example 2

Factor each completely.

(a) $x^2 + 15x + 36$

Product of the two numbers is 36.

Sum of the two numbers is 15.

Numbers are 3 and 12

Example 2

Factor each completely.

(a) $x^2 + 15x + 36$

Product of the two numbers is 36.

Sum of the two numbers is 15.

Numbers are 3 and 12


$$(x + 3)(x + 12)$$

Visual Approach to Factoring

$$x^2 + 15x + 36$$


Visual Approach to Factoring

$$x^2 + 15x + 36$$




Visual Approach to Factoring

$$x^2 + 15x + 36$$




Visual Approach to Factoring

$$x^2 + 15x + 36$$



Visual Approach to Factoring

$$x^2 + 15x + 36$$



Visual Approach to Factoring

$$x^2 + 15x + 36$$

	x	3
x	x^2	
12		36

Visual Approach to Factoring

$$x^2 + 15x + 36$$

	x	3
x	x^2	$3x$
12	$12x$	36

Visual Approach to Factoring

$$x^2 + 15x + 36$$

	x	3
x	x^2	$3x$
12	$12x$	36

$$(x + 3)(x + 12)$$

Example 2

(b) $x^2 - 9x + 14$

Example 2

(b) $x^2 - 9x + 14$

Product of the two numbers is 14.

Example 2

(b) $x^2 - 9x + 14$

Product of the two numbers is 14.

Sum of the two numbers is -9 .

Example 2

(b) $x^2 - 9x + 14$

Product of the two numbers is 14.

Sum of the two numbers is -9 .

Numbers are -7 and -2

Example 2

(b) $x^2 - 9x + 14$

Product of the two numbers is 14.

Sum of the two numbers is -9 .

Numbers are -7 and -2

$$(x - 7)(x - 2)$$

Example 2

(c) $x^2 + 5x - 6$

Example 2

(c) $x^2 + 5x - 6$

Product of the two numbers is -6 .

Example 2

(c) $x^2 + 5x - 6$

Product of the two numbers is -6 .

Sum of the two numbers is 5.

Example 2

(c) $x^2 + 5x - 6$

Product of the two numbers is -6 .

Sum of the two numbers is 5.

Numbers are -1 and 6

Example 2

(c) $x^2 + 5x - 6$

Product of the two numbers is -6 .

Sum of the two numbers is 5 .

Numbers are -1 and 6

$$(x - 1)(x + 6)$$

Example 2

(d) $x^2 - 7x - 8$

Example 2

(d) $x^2 - 7x - 8$

Product of the two numbers is -8 .

Example 2

(d) $x^2 - 7x - 8$

Product of the two numbers is -8 .

Sum of the two numbers is -7 .

Example 2

(d) $x^2 - 7x - 8$

Product of the two numbers is -8 .

Sum of the two numbers is -7 .

Numbers are -8 and 1

Example 2

(d) $x^2 - 7x - 8$

Product of the two numbers is -8 .

Sum of the two numbers is -7 .

Numbers are -8 and 1

$$(x - 8)(x + 1)$$

Objectives

- 1 Find the greatest common factor of a quadratic expression.
- 2 Factor trinomials in the form $x^2 + bx + c$
- 3 Factor trinomials in the form $ax^2 + bx + c$
- 4 Use “reverse engineering” to factor quadratic expressions

$$ax^2 + bx + c$$

To factor quadratic expressions in the form $ax^2 + bx + c$, we will first **transform** them into the form $x^2 + bx + c$ by **multiplying** the values of a and c to get:

$$ax^2 + bx + c$$

To factor quadratic expressions in the form $ax^2 + bx + c$, we will first **transform** them into the form $x^2 + bx + c$ by **multiplying** the values of a and c to get:

$$x^2 + bx + ac$$

$$ax^2 + bx + c$$

To factor quadratic expressions in the form $ax^2 + bx + c$, we will first **transform** them into the form $x^2 + bx + c$ by **multiplying** the values of a and c to get:

$$x^2 + bx + ac$$

We can then factor like in the last example, but we will have to remember to **transform the expression back**; as we will see.

Example 3

Factor each completely.

(a) $3x^2 - 11x - 4$

Example 3

Factor each completely.

(a) $3x^2 - 11x - 4$

Multiply $a = 3$ and $c = -4$ to get $ac = -12$.

Example 3

Factor each completely.

(a) $3x^2 - 11x - 4$

Multiply $a = 3$ and $c = -4$ to get $ac = -12$.

Factor $x^2 - 11x - 12$

Example 3

Factor each completely.

(a) $3x^2 - 11x - 4$

Multiply $a = 3$ and $c = -4$ to get $ac = -12$.

Factor $x^2 - 11x - 12$

$$(x - 12)(x + 1)$$

Example 3

Factor each completely.

(a) $3x^2 - 11x - 4$

Multiply $a = 3$ and $c = -4$ to get $ac = -12$.

Factor $x^2 - 11x - 12$

$$(x - 12)(x + 1)$$

We are not done yet!!

Example 3 $3x^2 - 11x - 4$

$$(x - 12)(x + 1)$$

Example 3 $3x^2 - 11x - 4$

$$(x - 12)(x + 1)$$

Divide the -12 and 1 by **3** and simplify.

Example 3 $3x^2 - 11x - 4$

$$(x - 12)(x + 1)$$

Divide the -12 and 1 by **3** and simplify.

$$\left(x - \frac{12}{3}\right) \left(x + \frac{1}{3}\right)$$

Example 3 $3x^2 - 11x - 4$

$$(x - 12)(x + 1)$$

Divide the -12 and 1 by **3** and simplify.

$$\left(x - \frac{12}{3}\right) \left(x + \frac{1}{3}\right)$$

$$(x - 4) \left(x + \frac{1}{3}\right)$$

Example 3 $3x^2 - 11x - 4$

$$(x - 12)(x + 1)$$

Divide the -12 and 1 by **3** and simplify.

$$\left(x - \frac{12}{3}\right) \left(x + \frac{1}{3}\right)$$

$$(x - 4) \left(x + \frac{1}{3}\right)$$

Slide any remaining denominators in front of their variable.

Example 3 $3x^2 - 11x - 4$

$$(x - 12)(x + 1)$$

Divide the -12 and 1 by **3** and simplify.

$$\left(x - \frac{12}{3}\right) \left(x + \frac{1}{3}\right)$$

$$(x - 4) \left(x + \frac{1}{3}\right)$$

Slide any remaining denominators in front of their variable.

$$(x - 4)(3x + 1)$$

Example 3

(b) $8x^2 + 2x - 3$

Example 3

(b) $8x^2 + 2x - 3$

Multiply $a = 8$ and $c = -3$ to get $ac = -24$

Example 3

(b) $8x^2 + 2x - 3$

Multiply $a = 8$ and $c = -3$ to get $ac = -24$

Factor $x^2 + 2x - 24$

Example 3

(b) $8x^2 + 2x - 3$

Multiply $a = 8$ and $c = -3$ to get $ac = -24$

Factor $x^2 + 2x - 24$

$$(x + 6)(x - 4)$$

Example 3

(b) $8x^2 + 2x - 3$

Multiply $a = 8$ and $c = -3$ to get $ac = -24$

Factor $x^2 + 2x - 24$

$$(x + 6)(x - 4)$$

$$\left(x + \frac{6}{8}\right) \left(x - \frac{4}{8}\right)$$

Example 3

(b) $8x^2 + 2x - 3$

Multiply $a = 8$ and $c = -3$ to get $ac = -24$

Factor $x^2 + 2x - 24$

$$(x + 6)(x - 4)$$

$$\left(x + \frac{6}{8}\right) \left(x - \frac{4}{8}\right)$$

$$\left(x + \frac{3}{4}\right) \left(x - \frac{1}{2}\right)$$

Example 3

(b) $8x^2 + 2x - 3$

Multiply $a = 8$ and $c = -3$ to get $ac = -24$

Factor $x^2 + 2x - 24$

$$(x + 6)(x - 4)$$

$$\left(x + \frac{6}{8}\right) \left(x - \frac{4}{8}\right)$$

$$\left(x + \frac{3}{4}\right) \left(x - \frac{1}{2}\right)$$

$$(4x + 3)(2x - 1)$$

Example 3

(c) $16x^2 + 46x + 15$

Example 3

(c) $16x^2 + 46x + 15$

Multiply $a = 16$ and $c = 15$ to get $ac = 240$

Example 3

(c) $16x^2 + 46x + 15$

Multiply $a = 16$ and $c = 15$ to get $ac = 240$

Factor $x^2 + 46x + 240$

Example 3

(c) $16x^2 + 46x + 15$

Multiply $a = 16$ and $c = 15$ to get $ac = 240$

Factor $x^2 + 46x + 240$

$$(x + 40)(x + 6)$$

Example 3

(c) $16x^2 + 46x + 15$

Multiply $a = 16$ and $c = 15$ to get $ac = 240$

Factor $x^2 + 46x + 240$

$$(x + 40)(x + 6)$$

$$\left(x + \frac{40}{16}\right) \left(x + \frac{6}{16}\right)$$

$$\left(x + \frac{5}{2}\right) \left(x + \frac{3}{8}\right)$$

Example 3

$$(c) \quad 16x^2 + 46x + 15$$

Multiply $a = 16$ and $c = 15$ to get $ac = 240$

Factor $x^2 + 46x + 240$

$$(x + 40)(x + 6)$$

$$\left(x + \frac{40}{16}\right) \left(x + \frac{6}{16}\right)$$

$$\left(x + \frac{5}{2}\right) \left(x + \frac{3}{8}\right)$$

$$(2x + 5)(8x + 3)$$

Example 3

(d) $7x^2 - 55x + 42$

Example 3

(d) $7x^2 - 55x + 42$

Multiply $a = 7$ and $c = 42$ to get $ac = 294$

Example 3

(d) $7x^2 - 55x + 42$

Multiply $a = 7$ and $c = 42$ to get $ac = 294$

Factor $x^2 - 55x + 294$

Example 3

(d) $7x^2 - 55x + 42$

Multiply $a = 7$ and $c = 42$ to get $ac = 294$

Factor $x^2 - 55x + 294$

$$(x - 49)(x - 6)$$

Example 3

$$(d) \quad 7x^2 - 55x + 42$$

Multiply $a = 7$ and $c = 42$ to get $ac = 294$

Factor $x^2 - 55x + 294$

$$(x - 49)(x - 6)$$

$$\left(x - \frac{49}{7}\right) \left(x - \frac{6}{7}\right)$$

$$(x - 7) \left(x - \frac{6}{7}\right)$$

Example 3

$$(d) \quad 7x^2 - 55x + 42$$

Multiply $a = 7$ and $c = 42$ to get $ac = 294$

Factor $x^2 - 55x + 294$

$$(x - 49)(x - 6)$$

$$\left(x - \frac{49}{7}\right) \left(x - \frac{6}{7}\right)$$

$$(x - 7) \left(x - \frac{6}{7}\right)$$

$$(x - 7)(7x - 6)$$

Objectives

- 1 Find the greatest common factor of a quadratic expression.
- 2 Factor trinomials in the form $x^2 + bx + c$
- 3 Factor trinomials in the form $ax^2 + bx + c$
- 4 Use “reverse engineering” to factor quadratic expressions

Using the Graph

You can also use the x -intercepts of the graph of the quadratic expression to help you factor.

Using the Graph

You can also use the x -intercepts of the graph of the quadratic expression to help you factor.

For instance, the graph for $x^2 + 5x - 6$ has x -intercepts at $x = -6$ and $x = 1$.

Using the Graph

You can also use the x -intercepts of the graph of the quadratic expression to help you factor.

For instance, the graph for $x^2 + 5x - 6$ has x -intercepts at $x = -6$ and $x = 1$.

$$x = -6$$

$$x = 1$$

Using the Graph

You can also use the x -intercepts of the graph of the quadratic expression to help you factor.

For instance, the graph for $x^2 + 5x - 6$ has x -intercepts at $x = -6$ and $x = 1$.

$$x = -6$$

$$x + 6 = 0$$

$$x = 1$$

$$x - 1 = 0$$

Using the Graph

You can also use the x -intercepts of the graph of the quadratic expression to help you factor.

For instance, the graph for $x^2 + 5x - 6$ has x -intercepts at $x = -6$ and $x = 1$.

$$x = -6$$

$$x + 6 = 0$$

$$x = 1$$

$$x - 1 = 0$$

$$x^2 + 5x - 6 \text{ factors as } (x + 6)(x - 1)$$

Using the Graph for $ax^2 + bx + c$

For expressions in the form $ax^2 + bx + c$:

Using the Graph for $ax^2 + bx + c$

For expressions in the form $ax^2 + bx + c$:

- 1 Get the transformation expression by multiplying a and c .

Using the Graph for $ax^2 + bx + c$

For expressions in the form $ax^2 + bx + c$:

- 1 Get the transformation expression by multiplying a and c .
- 2 Find those x -intercepts

Using the Graph for $ax^2 + bx + c$

For expressions in the form $ax^2 + bx + c$:

- 1 Get the transformation expression by multiplying a and c .
- 2 Find those x -intercepts
- 3 Transform back by dividing by the value of a and simplifying like in Example 3.

Using the Graph for $ax^2 + bx + c$

For instance, we transformed $3x^2 - 11x - 4$ to $x^2 - 11x - 12$

Using the Graph for $ax^2 + bx + c$

For instance, we transformed $3x^2 - 11x - 4$ to $x^2 - 11x - 12$

The x -intercepts of $x^2 - 11x - 12$ are $x = -1$ and $x = 12$

Using the Graph for $ax^2 + bx + c$

For instance, we transformed $3x^2 - 11x - 4$ to $x^2 - 11x - 12$

The x -intercepts of $x^2 - 11x - 12$ are $x = -1$ and $x = 12$

Divide each of those by 3 to get $x = -\frac{1}{3}$ and $x = 4$

Using the Graph for $ax^2 + bx + c$

For instance, we transformed $3x^2 - 11x - 4$ to $x^2 - 11x - 12$

The x -intercepts of $x^2 - 11x - 12$ are $x = -1$ and $x = 12$

Divide each of those by 3 to get $x = -\frac{1}{3}$ and $x = 4$

Move any remaining denominators in front of their variable:

Using the Graph for $ax^2 + bx + c$

For instance, we transformed $3x^2 - 11x - 4$ to $x^2 - 11x - 12$

The x -intercepts of $x^2 - 11x - 12$ are $x = -1$ and $x = 12$

Divide each of those by 3 to get $x = -\frac{1}{3}$ and $x = 4$

Move any remaining denominators in front of their variable:

$$3x = -1 \quad \text{and} \quad x = 4$$

Using the Graph for $ax^2 + bx + c$

For instance, we transformed $3x^2 - 11x - 4$ to $x^2 - 11x - 12$

The x -intercepts of $x^2 - 11x - 12$ are $x = -1$ and $x = 12$

Divide each of those by 3 to get $x = -\frac{1}{3}$ and $x = 4$

Move any remaining denominators in front of their variable:

$$3x = -1 \quad \text{and} \quad x = 4$$

Get each equal to 0:

Using the Graph for $ax^2 + bx + c$

For instance, we transformed $3x^2 - 11x - 4$ to $x^2 - 11x - 12$

The x -intercepts of $x^2 - 11x - 12$ are $x = -1$ and $x = 12$

Divide each of those by 3 to get $x = -\frac{1}{3}$ and $x = 4$

Move any remaining denominators in front of their variable:

$$3x = -1 \quad \text{and} \quad x = 4$$

Get each equal to 0:

$$3x + 1 \quad \text{and} \quad x - 4$$

Using the Graph for $ax^2 + bx + c$

For instance, we transformed $3x^2 - 11x - 4$ to $x^2 - 11x - 12$

The x -intercepts of $x^2 - 11x - 12$ are $x = -1$ and $x = 12$

Divide each of those by 3 to get $x = -\frac{1}{3}$ and $x = 4$

Move any remaining denominators in front of their variable:

$$3x = -1 \quad \text{and} \quad x = 4$$

Get each equal to 0:

$$3x + 1 \quad \text{and} \quad x - 4$$

$$(3x + 1)(x - 4)$$