

Quadratic Formula

Objectives

The Quadratic Formula

The **quadratic formula** can be used to solve any quadratic equation that is equal to 0.

The Quadratic Formula

The **quadratic formula** can be used to solve any quadratic equation that is equal to 0.

$$\text{For } ax^2 + bx + c = 0$$

Quadratic Formula

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Example 1

Solve each of the following using the quadratic formula. Exact answers only.

(a) $3x^2 + 8x - 28 = 0$

Example 1

Solve each of the following using the quadratic formula. Exact answers only.

(a) $3x^2 + 8x - 28 = 0$

$$a = 3 \quad b = 8 \quad c = -28$$

Example 1

Solve each of the following using the quadratic formula. Exact answers only.

$$(a) \quad 3x^2 + 8x - 28 = 0$$

$$a = 3 \quad b = 8 \quad c = -28$$

$$x = \frac{-8 \pm \sqrt{8^2 - 4(3)(-28)}}{2(3)}$$

Example 1

Solve each of the following using the quadratic formula. Exact answers only.

$$(a) \quad 3x^2 + 8x - 28 = 0$$

$$a = 3 \quad b = 8 \quad c = -28$$

$$x = \frac{-8 \pm \sqrt{8^2 - 4(3)(-28)}}{2(3)}$$

$$x = \frac{-8 \pm \sqrt{400}}{6}$$

Example 1

Solve each of the following using the quadratic formula. Exact answers only.

$$(a) \quad 3x^2 + 8x - 28 = 0$$

$$a = 3 \quad b = 8 \quad c = -28$$

$$x = \frac{-8 \pm \sqrt{8^2 - 4(3)(-28)}}{2(3)}$$

$$x = \frac{-8 \pm \sqrt{400}}{6}$$

$$x = \frac{-8 \pm 20}{6}$$

Example 1

$$x = \frac{-8 \pm 20}{6}$$

Example 1

$$x = \frac{-8 \pm 20}{6}$$

$$x = \frac{-8 + 20}{6}$$

$$x = \frac{-8 - 20}{6}$$

Example 1

$$x = \frac{-8 \pm 20}{6}$$

$$x = \frac{-8 + 20}{6}$$

$$x = 2$$

$$x = \frac{-8 - 20}{6}$$

$$x = -\frac{14}{3}$$

Example 1

$$(b) \quad 5x^2 + 9x - 5 = 0$$

Example 1

$$(b) \quad 5x^2 + 9x - 5 = 0$$

$$a = 5 \quad b = 9 \quad c = -5$$

Example 1

$$(b) \quad 5x^2 + 9x - 5 = 0$$

$$a = 5 \quad b = 9 \quad c = -5$$

$$x = \frac{-9 \pm \sqrt{9^2 - 4(5)(-5)}}{2(5)}$$

Example 1

$$(b) \quad 5x^2 + 9x - 5 = 0$$

$$a = 5 \quad b = 9 \quad c = -5$$

$$x = \frac{-9 \pm \sqrt{9^2 - 4(5)(-5)}}{2(5)}$$

$$x = \frac{-9 \pm \sqrt{181}}{10}$$

Example 1

$$(b) \quad 5x^2 + 9x - 5 = 0$$

$$a = 5 \quad b = 9 \quad c = -5$$

$$x = \frac{-9 \pm \sqrt{9^2 - 4(5)(-5)}}{2(5)}$$

$$x = \frac{-9 \pm \sqrt{181}}{10}$$

$$x = \frac{-9 + \sqrt{181}}{10} \quad x = \frac{-9 - \sqrt{181}}{10}$$

Example 1

$$(c) \quad 6x^2 - 6x - 15 = -10$$

Example 1

$$(c) \quad 6x^2 - 6x - 15 = -10$$

$$6x^2 - 6x - 5 = 0$$

Example 1

$$(c) \quad 6x^2 - 6x - 15 = -10$$

$$6x^2 - 6x - 5 = 0$$

$$a = 6 \quad b = -6 \quad c = -5$$

Example 1

$$(c) \quad 6x^2 - 6x - 15 = -10$$

$$6x^2 - 6x - 5 = 0$$

$$a = 6 \quad b = -6 \quad c = -5$$

$$x = \frac{6 \pm \sqrt{6^2 - 4(6)(-5)}}{2(6)}$$

Example 1

$$(c) \quad 6x^2 - 6x - 15 = -10$$

$$6x^2 - 6x - 5 = 0$$

$$a = 6 \quad b = -6 \quad c = -5$$

$$x = \frac{6 \pm \sqrt{6^2 - 4(6)(-5)}}{2(6)}$$

$$x = \frac{6 \pm \sqrt{156}}{12}$$

Example 1

$$x = \frac{6 \pm \sqrt{156}}{12}$$

Example 1

$$x = \frac{6 \pm \sqrt{156}}{12}$$

$$x = \frac{6 \pm 2\sqrt{39}}{12}$$

Example 1

$$x = \frac{6 \pm \sqrt{156}}{12}$$

$$x = \frac{6 \pm 2\sqrt{39}}{12}$$

$$x = \frac{2(3 \pm \sqrt{39})}{12}$$

Example 1

$$x = \frac{6 \pm \sqrt{156}}{12}$$

$$x = \frac{6 \pm 2\sqrt{39}}{12}$$

$$x = \frac{2(3 \pm \sqrt{39})}{12}$$

$$x = \frac{3 \pm \sqrt{39}}{6}$$

Example 1

$$(d) \quad 7x^2 + 20x - 8 = -3x^2 - 1 + 10x$$

Example 1

$$(d) \quad 7x^2 + 20x - 8 = -3x^2 - 1 + 10x$$

$$10x^2 + 10x - 7$$

Example 1

$$(d) \quad 7x^2 + 20x - 8 = -3x^2 - 1 + 10x$$

$$10x^2 + 10x - 7$$

$$a = 10 \quad b = 10 \quad c = -7$$

Example 1

$$(d) \quad 7x^2 + 20x - 8 = -3x^2 - 1 + 10x$$

$$10x^2 + 10x - 7$$

$$a = 10 \quad b = 10 \quad c = -7$$

$$x = \frac{-10 \pm \sqrt{10^2 - 4(10)(-7)}}{2(10)}$$

Example 1

$$(d) \quad 7x^2 + 20x - 8 = -3x^2 - 1 + 10x$$

$$10x^2 + 10x - 7$$

$$a = 10 \quad b = 10 \quad c = -7$$

$$x = \frac{-10 \pm \sqrt{10^2 - 4(10)(-7)}}{2(10)}$$

$$x = \frac{-10 \pm \sqrt{380}}{20}$$

Example 1

$$x = \frac{-10 \pm \sqrt{380}}{20}$$

Example 1

$$x = \frac{-10 \pm \sqrt{380}}{20}$$

$$x = \frac{-10 \pm 2\sqrt{95}}{20}$$

Example 1

$$x = \frac{-10 \pm \sqrt{380}}{20}$$

$$x = \frac{-10 \pm 2\sqrt{95}}{20}$$

$$x = \frac{2(-5 \pm \sqrt{95})}{20}$$

Example 1

$$x = \frac{-10 \pm \sqrt{380}}{20}$$

$$x = \frac{-10 \pm 2\sqrt{95}}{20}$$

$$x = \frac{2(-5 \pm \sqrt{95})}{20}$$

$$x = \frac{-5 \pm \sqrt{95}}{10}$$

Objectives

The Discriminant

The expression $b^2 - 4ac$ in the square root is called the **discriminant**. It can tell us about the solutions to a quadratic equation:

The Discriminant

The expression $b^2 - 4ac$ in the square root is called the **discriminant**. It can tell us about the solutions to a quadratic equation:

- **Discriminant is negative:** No real values of x make the original equation true.

The Discriminant

The expression $b^2 - 4ac$ in the square root is called the **discriminant**. It can tell us about the solutions to a quadratic equation:

- **Discriminant is negative:** No real values of x make the original equation true.
- **Discriminant is 0:** There is one value of x (called a *double root*).

The Discriminant

The expression $b^2 - 4ac$ in the square root is called the **discriminant**. It can tell us about the solutions to a quadratic equation:

- **Discriminant is negative:** No real values of x make the original equation true.
- **Discriminant is 0:** There is one value of x (called a *double root*).
- **Discriminant is positive:** There are 2 unique answers for x .

The Discriminant

In addition, if $\sqrt{b^2 - 4ac}$ equals a **rational number**, then the quadratic equation is factorable over the integers (only use integers in your factoring).