

# Factoring Techniques

# Objectives

- 1 Find the greatest common factor of a quadratic expression.
- 2 Factor trinomials in the form  $x^2 + bx + c$
- 3 Factor trinomials in the form  $ax^2 + bx + c$
- 4 Use “reverse engineering” to factor quadratic expressions

# Greatest Common Factor

The **greatest common factor**, or GCF, of an expression is the largest term that divides into each term of that expression.

# Greatest Common Factor

The **greatest common factor**, or GCF, of an expression is the largest term that divides into each term of that expression.

To find the greatest common factor of an expression:

# Greatest Common Factor

The **greatest common factor**, or GCF, of an expression is the largest term that divides into each term of that expression.

To find the greatest common factor of an expression:

- Find the greatest common factor of the coefficients.

# Greatest Common Factor

The **greatest common factor**, or GCF, of an expression is the largest term that divides into each term of that expression.

To find the greatest common factor of an expression:

- Find the greatest common factor of the coefficients.
- Select the lowest power of each unique variable.

# Greatest Common Factor

The **greatest common factor**, or GCF, of an expression is the largest term that divides into each term of that expression.

To find the greatest common factor of an expression:

- Find the greatest common factor of the coefficients.
- Select the lowest power of each unique variable.

Then divide each term by that GCF.

## Example 1

Factor the GCF from each.

(a)  $16x^5 + 72x^3 - 36x^2$



## Example 1

Factor the GCF from each.

(a)  $16x^5 + 72x^3 - 36x^2$

GCF of 16, 72, and 36 is 4

## Example 1

Factor the GCF from each.

(a)  $16x^5 + 72x^3 - 36x^2$

GCF of 16, 72, and 36 is 4

Lowest power of  $x$  is 2, so the GCF of the variables is  $x^2$

## Example 1

Factor the GCF from each.

(a)  $16x^5 + 72x^3 - 36x^2$

GCF of 16, 72, and 36 is 4

Lowest power of  $x$  is 2, so the GCF of the variables is  $x^2$

GCF is  $4x^2$

## Example 1

Factor the GCF from each.

$$(a) \quad 16x^5 + 72x^3 - 36x^2$$

GCF of 16, 72, and 36 is 4

Lowest power of  $x$  is 2, so the GCF of the variables is  $x^2$

GCF is  $4x^2$

$$4x^2 (4x^3 + 18x - 9)$$

## Example 1

$$(b) \quad 5a^4 + 70a^2 - 25a$$

## Example 1

$$(b) \quad 5a^4 + 70a^2 - 25a$$

GCF of 5, 70, and 25 is 5

## Example 1

$$(b) \quad 5a^4 + 70a^2 - 25a$$

GCF of 5, 70, and 25 is 5

Lowest power of  $a$  is 1, so the GCF of the variables is  $a$

## Example 1

$$(b) \quad 5a^4 + 70a^2 - 25a$$

GCF of 5, 70, and 25 is 5

Lowest power of  $a$  is 1, so the GCF of the variables is  $a$

GCF is  $5a$



## Example 1

$$(b) \quad 5a^4 + 70a^2 - 25a$$

GCF of 5, 70, and 25 is 5

Lowest power of  $a$  is 1, so the GCF of the variables is  $a$

GCF is  $5a$

$$5a(a^3 + 14a - 5)$$

# Objectives

- 1 Find the greatest common factor of a quadratic expression.
- 2 Factor trinomials in the form  $x^2 + bx + c$
- 3 Factor trinomials in the form  $ax^2 + bx + c$
- 4 Use “reverse engineering” to factor quadratic expressions

$$x^2 + bx + c$$

When factoring trinomials in the form  $x^2 + bx + c$ , we are looking to find 2 numbers that **multiply** to make  $c$  and **add** to make  $b$ .

$$x^2 + bx + c$$

When factoring trinomials in the form  $x^2 + bx + c$ , we are looking to find 2 numbers that **multiply** to make  $c$  and **add** to make  $b$ .

We can then write the factorization in the form

$$(x + m)(x + n)$$

where  $mn = c$  and  $m + n = b$

## Example 2

Factor each completely.

(a)  $x^2 + 15x + 36$

## Example 2

Factor each completely.

(a)  $x^2 + 15x + 36$

Product of the two numbers is 36.

## Example 2

Factor each completely.

(a)  $x^2 + 15x + 36$

Product of the two numbers is 36.

Sum of the two numbers is 15.

## Example 2

Factor each completely.

(a)  $x^2 + 15x + 36$

Product of the two numbers is 36.

Sum of the two numbers is 15.

Numbers are 3 and 12



## Example 2

Factor each completely.

(a)  $x^2 + 15x + 36$

Product of the two numbers is 36.

Sum of the two numbers is 15.

Numbers are 3 and 12


$$(x + 3)(x + 12)$$

# Visual Approach to Factoring

$$x^2 + 15x + 36$$


# Visual Approach to Factoring

$$x^2 + 15x + 36$$




# Visual Approach to Factoring

$$x^2 + 15x + 36$$




# Visual Approach to Factoring

$$x^2 + 15x + 36$$



# Visual Approach to Factoring

$$x^2 + 15x + 36$$



# Visual Approach to Factoring

$$x^2 + 15x + 36$$

	$x$	$3$
$x$	$x^2$	
$12$		$36$

# Visual Approach to Factoring

$$x^2 + 15x + 36$$

	$x$	$3$
$x$	$x^2$	$3x$
$12$	$12x$	$36$



# Visual Approach to Factoring

$$x^2 + 15x + 36$$

	$x$	$3$
$x$	$x^2$	$3x$
$12$	$12x$	$36$

$$(x + 3)(x + 12)$$

## Example 2

(b)  $x^2 - 9x + 14$

## Example 2

(b)  $x^2 - 9x + 14$

Product of the two numbers is 14.

## Example 2

(b)  $x^2 - 9x + 14$

Product of the two numbers is 14.

Sum of the two numbers is  $-9$ .

## Example 2

(b)  $x^2 - 9x + 14$

Product of the two numbers is 14.

Sum of the two numbers is  $-9$ .

Numbers are  $-7$  and  $-2$

## Example 2

(b)  $x^2 - 9x + 14$

Product of the two numbers is 14.

Sum of the two numbers is  $-9$ .

Numbers are  $-7$  and  $-2$

$$(x - 7)(x - 2)$$

## Example 2

(c)  $x^2 + 5x - 6$

## Example 2

(c)  $x^2 + 5x - 6$

Product of the two numbers is  $-6$ .



## Example 2

(c)  $x^2 + 5x - 6$

Product of the two numbers is  $-6$ .

Sum of the two numbers is 5.

## Example 2

(c)  $x^2 + 5x - 6$

Product of the two numbers is  $-6$ .

Sum of the two numbers is 5.

Numbers are  $-1$  and 6

## Example 2

(c)  $x^2 + 5x - 6$

Product of the two numbers is  $-6$ .

Sum of the two numbers is  $5$ .

Numbers are  $-1$  and  $6$

$$(x - 1)(x + 6)$$

## Example 2

(d)  $x^2 - 7x - 8$

## Example 2

(d)  $x^2 - 7x - 8$

Product of the two numbers is  $-8$ .

## Example 2

(d)  $x^2 - 7x - 8$

Product of the two numbers is  $-8$ .

Sum of the two numbers is  $-7$ .

## Example 2

(d)  $x^2 - 7x - 8$

Product of the two numbers is  $-8$ .

Sum of the two numbers is  $-7$ .

Numbers are  $-8$  and  $1$

## Example 2

(d)  $x^2 - 7x - 8$

Product of the two numbers is  $-8$ .

Sum of the two numbers is  $-7$ .

Numbers are  $-8$  and  $1$

$$(x - 8)(x + 1)$$



# Objectives

- 1 Find the greatest common factor of a quadratic expression.
- 2 Factor trinomials in the form  $x^2 + bx + c$
- 3 Factor trinomials in the form  $ax^2 + bx + c$
- 4 Use “reverse engineering” to factor quadratic expressions

$$ax^2 + bx + c$$

To factor quadratic expressions in the form  $ax^2 + bx + c$ , we will first **transform** them into the form  $x^2 + bx + c$  by **multiplying** the values of  $a$  and  $c$  to get:

$$ax^2 + bx + c$$

To factor quadratic expressions in the form  $ax^2 + bx + c$ , we will first **transform** them into the form  $x^2 + bx + c$  by **multiplying** the values of  $a$  and  $c$  to get:

$$x^2 + bx + ac$$

$$ax^2 + bx + c$$

To factor quadratic expressions in the form  $ax^2 + bx + c$ , we will first **transform** them into the form  $x^2 + bx + c$  by **multiplying** the values of  $a$  and  $c$  to get:

$$x^2 + bx + ac$$

We can then factor like in the last example, but we will have to remember to **transform the expression back**; as we will see.

## Example 3

Factor each completely.

(a)  $3x^2 - 11x - 4$

## Example 3

Factor each completely.

(a)  $3x^2 - 11x - 4$

Multiply  $a = 3$  and  $c = -4$  to get  $ac = -12$ .

## Example 3

Factor each completely.

(a)  $3x^2 - 11x - 4$

Multiply  $a = 3$  and  $c = -4$  to get  $ac = -12$ .

Factor  $x^2 - 11x - 12$

## Example 3

Factor each completely.

(a)  $3x^2 - 11x - 4$

Multiply  $a = 3$  and  $c = -4$  to get  $ac = -12$ .

Factor  $x^2 - 11x - 12$

$$(x - 12)(x + 1)$$



## Example 3

Factor each completely.

(a)  $3x^2 - 11x - 4$

Multiply  $a = 3$  and  $c = -4$  to get  $ac = -12$ .

Factor  $x^2 - 11x - 12$

$$(x - 12)(x + 1)$$

**We are not done yet!!**

### Example 3 $3x^2 - 11x - 4$

$$(x - 12)(x + 1)$$

### Example 3 $3x^2 - 11x - 4$

$$(x - 12)(x + 1)$$

Divide the  $-12$  and  $1$  by **3** and simplify.

### Example 3 $3x^2 - 11x - 4$

$$(x - 12)(x + 1)$$

Divide the  $-12$  and  $1$  by **3** and simplify.

$$\left(x - \frac{12}{3}\right) \left(x + \frac{1}{3}\right)$$

### Example 3 $3x^2 - 11x - 4$

$$(x - 12)(x + 1)$$

Divide the  $-12$  and  $1$  by **3** and simplify.

$$\left(x - \frac{12}{3}\right) \left(x + \frac{1}{3}\right)$$

$$(x - 4) \left(x + \frac{1}{3}\right)$$

### Example 3 $3x^2 - 11x - 4$

$$(x - 12)(x + 1)$$

Divide the  $-12$  and  $1$  by **3** and simplify.

$$\left(x - \frac{12}{3}\right) \left(x + \frac{1}{3}\right)$$

$$(x - 4) \left(x + \frac{1}{3}\right)$$

Slide any remaining denominators in front of their variable.

### Example 3 $3x^2 - 11x - 4$

$$(x - 12)(x + 1)$$

Divide the  $-12$  and  $1$  by **3** and simplify.

$$\left(x - \frac{12}{3}\right) \left(x + \frac{1}{3}\right)$$

$$(x - 4) \left(x + \frac{1}{3}\right)$$

Slide any remaining denominators in front of their variable.

$$(x - 4)(3x + 1)$$

## Example 3

(b)  $8x^2 + 2x - 3$



## Example 3

(b)  $8x^2 + 2x - 3$

Multiply  $a = 8$  and  $c = -3$  to get  $ac = -24$

## Example 3

(b)  $8x^2 + 2x - 3$

Multiply  $a = 8$  and  $c = -3$  to get  $ac = -24$

Factor  $x^2 + 2x - 24$

## Example 3

(b)  $8x^2 + 2x - 3$

Multiply  $a = 8$  and  $c = -3$  to get  $ac = -24$

Factor  $x^2 + 2x - 24$

$$(x + 6)(x - 4)$$

## Example 3

(b)  $8x^2 + 2x - 3$

Multiply  $a = 8$  and  $c = -3$  to get  $ac = -24$

Factor  $x^2 + 2x - 24$

$$(x + 6)(x - 4)$$

$$\left(x + \frac{6}{8}\right) \left(x - \frac{4}{8}\right)$$

## Example 3

(b)  $8x^2 + 2x - 3$

Multiply  $a = 8$  and  $c = -3$  to get  $ac = -24$

Factor  $x^2 + 2x - 24$

$$(x + 6)(x - 4)$$

$$\left(x + \frac{6}{8}\right) \left(x - \frac{4}{8}\right)$$

$$\left(x + \frac{3}{4}\right) \left(x - \frac{1}{2}\right)$$

## Example 3

(b)  $8x^2 + 2x - 3$

Multiply  $a = 8$  and  $c = -3$  to get  $ac = -24$

Factor  $x^2 + 2x - 24$

$$(x + 6)(x - 4)$$

$$\left(x + \frac{6}{8}\right) \left(x - \frac{4}{8}\right)$$

$$\left(x + \frac{3}{4}\right) \left(x - \frac{1}{2}\right)$$

$$(4x + 3)(2x - 1)$$

## Example 3

(c)  $16x^2 + 46x + 15$

## Example 3

(c)  $16x^2 + 46x + 15$

Multiply  $a = 16$  and  $c = 15$  to get  $ac = 240$



## Example 3

(c)  $16x^2 + 46x + 15$

Multiply  $a = 16$  and  $c = 15$  to get  $ac = 240$

Factor  $x^2 + 46x + 240$

## Example 3

(c)  $16x^2 + 46x + 15$

Multiply  $a = 16$  and  $c = 15$  to get  $ac = 240$

Factor  $x^2 + 46x + 240$

$$(x + 40)(x + 6)$$

## Example 3

(c)  $16x^2 + 46x + 15$

Multiply  $a = 16$  and  $c = 15$  to get  $ac = 240$

Factor  $x^2 + 46x + 240$

$$(x + 40)(x + 6)$$

$$\left(x + \frac{40}{16}\right) \left(x + \frac{6}{16}\right)$$

$$\left(x + \frac{5}{2}\right) \left(x + \frac{3}{8}\right)$$

## Example 3

$$(c) \quad 16x^2 + 46x + 15$$

Multiply  $a = 16$  and  $c = 15$  to get  $ac = 240$

Factor  $x^2 + 46x + 240$

$$(x + 40)(x + 6)$$

$$\left(x + \frac{40}{16}\right) \left(x + \frac{6}{16}\right)$$

$$\left(x + \frac{5}{2}\right) \left(x + \frac{3}{8}\right)$$

$$(2x + 5)(8x + 3)$$

## Example 3

(d)  $7x^2 - 55x + 42$

## Example 3

(d)  $7x^2 - 55x + 42$

Multiply  $a = 7$  and  $c = 42$  to get  $ac = 294$

## Example 3

(d)  $7x^2 - 55x + 42$

Multiply  $a = 7$  and  $c = 42$  to get  $ac = 294$

Factor  $x^2 - 55x + 294$

## Example 3

(d)  $7x^2 - 55x + 42$

Multiply  $a = 7$  and  $c = 42$  to get  $ac = 294$

Factor  $x^2 - 55x + 294$

$$(x - 49)(x - 6)$$



## Example 3

$$(d) \quad 7x^2 - 55x + 42$$

Multiply  $a = 7$  and  $c = 42$  to get  $ac = 294$

Factor  $x^2 - 55x + 294$

$$(x - 49)(x - 6)$$

$$\left(x - \frac{49}{1}\right) \left(x - \frac{6}{1}\right)$$

$$(x - 7) \left(x - \frac{6}{7}\right)$$

## Example 3

$$(d) \quad 7x^2 - 55x + 42$$

Multiply  $a = 7$  and  $c = 42$  to get  $ac = 294$

Factor  $x^2 - 55x + 294$

$$(x - 49)(x - 6)$$

$$\left(x - \frac{49}{7}\right) \left(x - \frac{6}{7}\right)$$

$$(x - 7) \left(x - \frac{6}{7}\right)$$

$$(x - 7)(7x - 6)$$

# Objectives

- 1 Find the greatest common factor of a quadratic expression.
- 2 Factor trinomials in the form  $x^2 + bx + c$
- 3 Factor trinomials in the form  $ax^2 + bx + c$
- 4 Use “reverse engineering” to factor quadratic expressions

# Using the Graph

You can also use the  $x$ -intercepts of the graph of the quadratic expression to help you factor.

# Using the Graph

You can also use the  $x$ -intercepts of the graph of the quadratic expression to help you factor.

For instance, the graph for  $x^2 + 5x - 6$  has  $x$ -intercepts at  $x = -6$  and  $x = 1$ .

# Using the Graph

You can also use the  $x$ -intercepts of the graph of the quadratic expression to help you factor.

For instance, the graph for  $x^2 + 5x - 6$  has  $x$ -intercepts at  $x = -6$  and  $x = 1$ .

$$x = -6$$

$$x = 1$$

# Using the Graph

You can also use the  $x$ -intercepts of the graph of the quadratic expression to help you factor.

For instance, the graph for  $x^2 + 5x - 6$  has  $x$ -intercepts at  $x = -6$  and  $x = 1$ .

$$x = -6$$

$$x + 6 = 0$$

$$x = 1$$

$$x - 1 = 0$$

# Using the Graph

You can also use the  $x$ -intercepts of the graph of the quadratic expression to help you factor.

For instance, the graph for  $x^2 + 5x - 6$  has  $x$ -intercepts at  $x = -6$  and  $x = 1$ .

$$x = -6$$

$$x + 6 = 0$$

$$x = 1$$

$$x - 1 = 0$$

$$x^2 + 5x - 6 \text{ factors as } (x + 6)(x - 1)$$



# Using the Graph for $ax^2 + bx + c$

For expressions in the form  $ax^2 + bx + c$ :

# Using the Graph for $ax^2 + bx + c$

For expressions in the form  $ax^2 + bx + c$ :

- 1 Get the transformation expression by multiplying  $a$  and  $c$ .

# Using the Graph for $ax^2 + bx + c$

For expressions in the form  $ax^2 + bx + c$ :

- 1 Get the transformation expression by multiplying  $a$  and  $c$ .
- 2 Find those  $x$ -intercepts

# Using the Graph for $ax^2 + bx + c$

For expressions in the form  $ax^2 + bx + c$ :

- 1 Get the transformation expression by multiplying  $a$  and  $c$ .
- 2 Find those  $x$ -intercepts
- 3 Transform back by dividing by the value of  $a$  and simplifying like in Example 3.

## Using the Graph for $ax^2 + bx + c$

For instance, we transformed  $3x^2 - 11x - 4$  to  $x^2 - 11x - 12$

## Using the Graph for $ax^2 + bx + c$

For instance, we transformed  $3x^2 - 11x - 4$  to  $x^2 - 11x - 12$

The  $x$ -intercepts of  $x^2 - 11x - 12$  are  $x = -1$  and  $x = 12$

## Using the Graph for $ax^2 + bx + c$

For instance, we transformed  $3x^2 - 11x - 4$  to  $x^2 - 11x - 12$

The  $x$ -intercepts of  $x^2 - 11x - 12$  are  $x = -1$  and  $x = 12$

Divide each of those by 3 to get  $x = -\frac{1}{3}$  and  $x = 4$

## Using the Graph for $ax^2 + bx + c$

For instance, we transformed  $3x^2 - 11x - 4$  to  $x^2 - 11x - 12$

The  $x$ -intercepts of  $x^2 - 11x - 12$  are  $x = -1$  and  $x = 12$

Divide each of those by 3 to get  $x = -\frac{1}{3}$  and  $x = 4$

Move any remaining denominators in front of their variable:



## Using the Graph for $ax^2 + bx + c$

For instance, we transformed  $3x^2 - 11x - 4$  to  $x^2 - 11x - 12$

The  $x$ -intercepts of  $x^2 - 11x - 12$  are  $x = -1$  and  $x = 12$

Divide each of those by 3 to get  $x = -\frac{1}{3}$  and  $x = 4$

Move any remaining denominators in front of their variable:

$$3x = -1 \quad \text{and} \quad x = 4$$

## Using the Graph for $ax^2 + bx + c$

For instance, we transformed  $3x^2 - 11x - 4$  to  $x^2 - 11x - 12$

The  $x$ -intercepts of  $x^2 - 11x - 12$  are  $x = -1$  and  $x = 12$

Divide each of those by 3 to get  $x = -\frac{1}{3}$  and  $x = 4$

Move any remaining denominators in front of their variable:

$$3x = -1 \quad \text{and} \quad x = 4$$

Get each equal to 0:

## Using the Graph for $ax^2 + bx + c$

For instance, we transformed  $3x^2 - 11x - 4$  to  $x^2 - 11x - 12$

The  $x$ -intercepts of  $x^2 - 11x - 12$  are  $x = -1$  and  $x = 12$

Divide each of those by 3 to get  $x = -\frac{1}{3}$  and  $x = 4$

Move any remaining denominators in front of their variable:

$$3x = -1 \quad \text{and} \quad x = 4$$

Get each equal to 0:

$$3x + 1 \quad \text{and} \quad x - 4$$

## Using the Graph for $ax^2 + bx + c$

For instance, we transformed  $3x^2 - 11x - 4$  to  $x^2 - 11x - 12$

The  $x$ -intercepts of  $x^2 - 11x - 12$  are  $x = -1$  and  $x = 12$

Divide each of those by 3 to get  $x = -\frac{1}{3}$  and  $x = 4$

Move any remaining denominators in front of their variable:

$$3x = -1 \quad \text{and} \quad x = 4$$

Get each equal to 0:

$$3x + 1 \quad \text{and} \quad x - 4$$

$$(3x + 1)(x - 4)$$