

Objectives

1 Find the domain of a function

2 Find the range of a function

Intro

Often times when evaluating functions, we are interested in which values we are allowed to put into the function, as well as which values the functions will give us back.

Often times when evaluating functions, we are interested in which values we are allowed to put into the function, as well as which values the functions will give us back.

Domain

The **domain** of a function is the set of all possible legal input values (x) of the function.

Intro

Often times when evaluating functions, we are interested in which values we are allowed to put into the function, as well as which values the functions will give us back.

Domain

The **domain** of a function is the set of all possible legal input values (x) of the function.

Range

The **range** of a function is the set of all possible output values (y) from the domain.

Domain Restrictions

Right now, there are only 2 things to worry about when finding the domain of a function:

Domain Restrictions

Right now, there are only 2 things to worry about when finding the domain of a function:

- You are not allowed to divide by 0

Domain Restrictions

Right now, there are only 2 things to worry about when finding the domain of a function:

- You are not allowed to divide by 0
- You can't take the square root (or any even root such as $\sqrt[4]{}$, $\sqrt[6]{}$, ...) of a negative number.

Domain Restrictions

Right now, there are only 2 things to worry about when finding the domain of a function:

- You are not allowed to divide by 0
- You can't take the square root (or any even root such as $\sqrt[4]{}$, $\sqrt[6]{}$, ...) of a negative number.

Both of the above issues will result in an error message from your calculator, and we would like to avoid those.

Domain

Aside from the 2 scenarios previously listed, most of the functions will have domains in which you can evaluate any value of x you like.

Domain

Aside from the 2 scenarios previously listed, most of the functions will have domains in which you can evaluate any value of x you like.

Those domains are **all real numbers**, or \mathbb{R} .

Example 1

State the domain of each.

(a) $f(x) = -2x + 7$

Example 1

State the domain of each.

(a) $f(x) = -2x + 7$

All real numbers, \mathbb{R}

Example 1

State the domain of each.

(a) $f(x) = -2x + 7$

All real numbers, \mathbb{R}

(b) $f(x) = (x + 4)^2$

Example 1

State the domain of each.

(a) $f(x) = -2x + 7$

All real numbers, \mathbb{R}

(b) $f(x) = (x + 4)^2$

\mathbb{R}

Example 1

$$(c) \quad f(x) = \sqrt{x - 3}$$

Example 1

$$(c) \quad f(x) = \sqrt{x-3}$$

$$x - 3 \geq 0$$

Example 1

$$(c) \quad f(x) = \sqrt{x-3}$$

$$x - 3 \geq 0$$

$$x \geq 3$$

Example 1

$$(c) \quad f(x) = \sqrt{x-3}$$

$$x - 3 \geq 0$$

$$x \geq 3$$

$$(d) \quad f(x) = \sqrt[3]{x-3}$$

Example 1

$$(c) \quad f(x) = \sqrt{x-3}$$

$$x - 3 \geq 0$$

$$x \geq 3$$

$$(d) \quad f(x) = \sqrt[3]{x-3}$$

\mathbb{R}

Example 1

$$(e) \quad f(x) = \frac{3}{2x + 5}$$

Example 1

$$(e) \quad f(x) = \frac{3}{2x + 5}$$

$$2x + 5 \neq 0$$

Example 1

$$(e) \quad f(x) = \frac{3}{2x + 5}$$

$$2x + 5 \neq 0$$

$$2x \neq -5$$

Example 1

$$(e) \quad f(x) = \frac{3}{2x + 5}$$

$$2x + 5 \neq 0$$

$$2x \neq -5$$

$$x \neq -\frac{5}{2}$$

Objectives

- 1 Find the domain of a function
- 2 Find the range of a function

Range

Finding the range can be a bit more challenging with some functions.

Range

Finding the range can be a bit more challenging with some functions.

When we look at finding the inverse of a function, we'll learn a way to find the range of a given function.

Range

Finding the range can be a bit more challenging with some functions.

When we look at finding the inverse of a function, we'll learn a way to find the range of a given function.

For now, we can look at the graphs of functions to assist us in finding the range.

Example 2

State the range of each.

(a) $f(x) = -2x + 7$

Example 2

State the range of each.

(a) $f(x) = -2x + 7$

All real numbers, \mathbb{R}

Example 2

$$(b) \quad f(x) = (x + 4)^2$$

Example 2

$$(b) \quad f(x) = (x + 4)^2$$

$$y \geq 0$$

Example 2

$$(c) \quad f(x) = \sqrt{x - 3}$$

Example 2

$$(c) \quad f(x) = \sqrt{x - 3}$$

$$y \geq 0$$

Example 2

$$(d) \quad f(x) = -\sqrt{x-3}$$

Example 2

$$(d) \quad f(x) = -\sqrt{x-3}$$

$$y \leq 0$$

Example 2

$$(e) \quad f(x) = \sqrt{x-3} + 6$$

Example 2

$$(e) \quad f(x) = \sqrt{x-3} + 6$$

$$y \geq 6$$

Example 2

$$(f) \quad f(x) = \frac{3}{2x + 5}$$

Example 2

$$(f) \quad f(x) = \frac{3}{2x + 5}$$

$$y \neq 0$$