Systems of Equations

Objectives

Write a system of equations using matrices

2 Solve a system of equations using inverse matrices

3 Solve applications of systems of equations

Matrix Multiplication

Previously, we looked at the method for multiplying matrices:

$$\begin{bmatrix} 7 & 2 & -1 \\ 0 & 5 & 4 \\ -3 & 6 & 2 \end{bmatrix} \cdot \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}$$

Matrix Multiplication

Previously, we looked at the method for multiplying matrices:

$$\begin{bmatrix} 7 & 2 & -1 \\ 0 & 5 & 4 \\ -3 & 6 & 2 \end{bmatrix} \cdot \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix} = \begin{bmatrix} 13 \\ 17 \\ 6 \end{bmatrix}$$

Matrix Multiplication

Previously, we looked at the method for multiplying matrices:

$$\begin{bmatrix} 7 & 2 & -1 \\ 0 & 5 & 4 \\ -3 & 6 & 2 \end{bmatrix} \cdot \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix} = \begin{bmatrix} 13 \\ 17 \\ 6 \end{bmatrix}$$

In this section, we will use matrix multiplication to solve a system of equations.

System of Equations

A **system of equations** consists of 2 or more equations with 2 or more variables.

System of Equations

A **system of equations** consists of 2 or more equations with 2 or more variables.

EXAMPLES

System of Equations

A **system of equations** consists of 2 or more equations with 2 or more variables.

EXAMPLES

$$2x + 5y = 8$$
$$-7x - 3y = 9$$

$$4x - 9y + 2z = 10$$
$$-x + z = 15$$
$$3x + 10y - 12z = 0$$

We can use matrix multiplication with the coefficients and variables to write a system of equations using matrices.

We can use matrix multiplication with the coefficients and variables to write a system of equations using matrices.

So

$$2x + 5y = 8$$
$$-7x - 3y = 9$$

We can use matrix multiplication with the coefficients and variables to write a system of equations using matrices.

So

$$2x + 5y = 8$$
$$-7x - 3y = 9$$

becomes

$$\begin{bmatrix} 2 & 5 \\ -7 & -3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 8 \\ 9 \end{bmatrix}$$

And

$$4x - 9y + 2z = 10$$
$$-x + z = 15$$
$$3x + 10y - 12z = 0$$

And

$$4x - 9y + 2z = 10$$
$$-x + z = 15$$
$$3x + 10y - 12z = 0$$

becomes

$$\begin{bmatrix} 4 & -9 & 2 \\ -1 & 0 & 1 \\ 3 & 10 & -12 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 10 \\ 15 \\ 0 \end{bmatrix}$$

Write each of the following systems of equations using matrices.

$$4y + z = -1$$
$$-4x - 5y + 5z = 0$$
$$6x + 5y = 25$$

Write each of the following systems of equations using matrices.

$$4y + z = -1$$
$$-4x - 5y + 5z = 0$$
$$6x + 5y = 25$$

$$\begin{bmatrix} 0 & 4 & 1 \\ -4 & -5 & 5 \\ 6 & 5 & 0 \end{bmatrix}$$

Write each of the following systems of equations using matrices.

$$4y + z = -1$$
$$-4x - 5y + 5z = 0$$
$$6x + 5y = 25$$

$$\begin{bmatrix} 0 & 4 & 1 \\ -4 & -5 & 5 \\ 6 & 5 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

Write each of the following systems of equations using matrices.

$$4y + z = -1$$
$$-4x - 5y + 5z = 0$$
$$6x + 5y = 25$$

$$\begin{bmatrix} 0 & 4 & 1 \\ -4 & -5 & 5 \\ 6 & 5 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -1 \\ 0 \\ 25 \end{bmatrix}$$

$$x - y = -9$$
$$-3x - 4y = -8$$

(b)

$$x - y = -9$$
$$-3x - 4y = -8$$

$$\begin{bmatrix} 1 & -1 \\ -3 & -4 \end{bmatrix}$$

(b)

$$x - y = -9$$
$$-3x - 4y = -8$$

$$\begin{bmatrix} 1 & -1 \\ -3 & -4 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

(b)

$$x - y = -9$$
$$-3x - 4y = -8$$

$$\begin{bmatrix} 1 & -1 \\ -3 & -4 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -9 \\ -8 \end{bmatrix}$$

Objectives

1 Write a system of equations using matrices

2 Solve a system of equations using inverse matrices

3 Solve applications of systems of equations

Solving Equations Using Inverse Operations

Long ago, you learned that to solve something like

$$5x = 10$$

you divide (the inverse operation of multiplication) both sides by 5.

Solving Equations Using Inverse Operations

Long ago, you learned that to solve something like

$$5x = 10$$

you divide (the inverse operation of multiplication) both sides by 5.

We don't "divide" matrices in this sense, but we do need to use an inverse operation to solve the previous examples.

Identity Matrices

Identity Matrix

An **identity matrix** is a square matrix with 1s along the diagonal from top left to bottom right, and 0s elsewhere.

Identity Matrices

Identity Matrix

An **identity matrix** is a square matrix with 1s along the diagonal from top left to bottom right, and 0s elsewhere.

$$I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$
 $I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ etc.

Inverse Matrix

The **inverse matrix** for square matrix A is A^{-1} .

Inverse Matrix

The **inverse matrix** for square matrix A is A^{-1} .

Inverse matrices are such that

$$A\cdot A^{-1}=I$$

Inverse Matrix

The **inverse matrix** for square matrix A is A^{-1} .

Inverse matrices are such that

$$A \cdot A^{-1} = I$$

and that if AX = B where A is the matrix of coefficients and B are the constants on the right side, then

$$X = A^{-1} \cdot B$$

Inverse Matrix

The **inverse matrix** for square matrix A is A^{-1} .

Inverse matrices are such that

$$A \cdot A^{-1} = I$$

and that if AX = B where A is the matrix of coefficients and B are the constants on the right side, then

$$X = A^{-1} \cdot B$$

Note: We will not discuss the techniques of how to actually find the inverse of a matrix without a calculator.

Solve each of the following using matrices.

$$4y + z = -1$$
$$-4x - 5y + 5z = 0$$
$$6x + 5y = 25$$

Solve each of the following using matrices.

$$4y + z = -1$$
$$-4x - 5y + 5z = 0$$
$$6x + 5y = 25$$

$$\begin{bmatrix} 0 & 4 & 1 \\ -4 & -5 & 5 \\ 6 & 5 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -1 \\ 0 \\ 25 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = A^{-1}B$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = A^{-1}B$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 5 \\ -1 \\ 3 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = A^{-1}B$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 5 \\ -1 \\ 3 \end{bmatrix}$$

$$x = 5$$
, $y = -1$, $z = 3$

$$x - y = -9$$
$$-3x - 4y = -8$$

$$x - y = -9$$
$$-3x - 4y = -8$$

$$\begin{bmatrix} 1 & -1 \\ -3 & -4 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -9 \\ -8 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = A^{-1}B$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = A^{-1}B$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -4 \\ 5 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = A^{-1}B$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -4 \\ 5 \end{bmatrix}$$

$$x = -4, y = 5$$

(c)

$$x - 3y - 5z = -18$$
$$-2x + 3y + 5z = 23$$
$$-x + 3y + 6z = 17$$

$$x - 3y - 5z = -18$$
$$-2x + 3y + 5z = 23$$
$$-x + 3y + 6z = 17$$

$$\begin{bmatrix} 1 & -3 & -5 \\ -2 & 3 & 5 \\ -1 & 3 & 6 \end{bmatrix} \cdot \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -18 \\ 23 \\ 17 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = A^{-1}B$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = A^{-1}B$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -5 \\ 6 \\ -1 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = A^{-1}B$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -5 \\ 6 \\ -1 \end{bmatrix}$$

$$x = -5$$
, $y = 6$, $z = -1$

Objectives

1 Write a system of equations using matrices

2 Solve a system of equations using inverse matrices

Solve applications of systems of equations

Applied Systems of Equations

Setting up many applied systems of equations problems boils down to

unit rate \times amount = total amount

For instance, \$2.00 per gallon times 5 gallons of gas costs a total amount of \$10.

(a) How many mL of a solution containing 10% pure hydrochloric acid must be mixed with a solution containing 15% hydrochloric acid to produce 30 mL of a solution that is 11% hydrochloric acid?

(a) How many mL of a solution containing 10% pure hydrochloric acid must be mixed with a solution containing 15% hydrochloric acid to produce 30 mL of a solution that is 11% hydrochloric acid?

x = mL of the 10% acid solution y = mL of the 15% acid solution

(a) How many mL of a solution containing 10% pure hydrochloric acid must be mixed with a solution containing 15% hydrochloric acid to produce 30 mL of a solution that is 11% hydrochloric acid?

x = mL of the 10% acid solution y = mL of the 15% acid solution

$$x+y=30$$
 volume of total liquid $0.10x+0.15y=30(0.11)$ volume of total hydrocholic acid

$$\begin{bmatrix} 1 & 1 \\ 0.1 & 0.15 \end{bmatrix} \cdot \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 30 \\ 3.3 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 \\ 0.1 & 0.15 \end{bmatrix} \cdot \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 30 \\ 3.3 \end{bmatrix}$$
$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 24 \\ 6 \end{bmatrix}$$

We need to mix 24 mL of a 10% hydrochloric acid solution with 6 mL of a 15% hydrochloric acid solution.

(b) A coffee shop wants to sell 8 pound bags of a coffee blend for \$28.64. They will do this by blending coffee that costs \$2.25 per pound with coffee that costs \$4.00 per pound. How much of each type of coffee should they use in the blend?

(b) A coffee shop wants to sell 8 pound bags of a coffee blend for \$28.64. They will do this by blending coffee that costs \$2.25 per pound with coffee that costs \$4.00 per pound. How much of each type of coffee should they use in the blend?

x = number of pounds of cheaper coffee (\$2.25 per pound) y = number of pounds of more expensive coffee (\$4.00 per pound)

(b) A coffee shop wants to sell 8 pound bags of a coffee blend for \$28.64. They will do this by blending coffee that costs \$2.25 per pound with coffee that costs \$4.00 per pound. How much of each type of coffee should they use in the blend?

x = number of pounds of cheaper coffee (\$2.25 per pound) y = number of pounds of more expensive coffee (\$4.00 per pound)

$$x + y = 8$$
 weight of each bag $2.25x + 4.00y = 28.64$ total cost

$$\begin{bmatrix} 1 & 1 \\ 2.25 & 4 \end{bmatrix} \cdot \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 8 \\ 28.64 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 \\ 2.25 & 4 \end{bmatrix} \cdot \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 8 \\ 28.64 \end{bmatrix}$$
$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1.92 \\ 6.08 \end{bmatrix}$$

They will need 1.92 pounds of the cheaper coffee and 6.08 pounds of the more expensive coffee.