

Trig Identities

Objectives

- 1 Derive the Pythagorean identities
- 2 Derive the quotient identities
- 3 Use the sum and difference identities to verify other identities
- 4 Derive double-angle identities
- 5 Derive power-reducing identities


Pythagorean Identities

These are some of the most important identities in this course.

They are based on the Pythagorean Theorem.

Example 1


Verify $\cos^2 \theta + \sin^2 \theta = 1$



Example 1


Verify $\cos^2 \theta + \sin^2 \theta = 1$

$$x^2 + y^2 = r^2$$



Example 1

Verify $\cos^2 \theta + \sin^2 \theta = 1$




$$x^2 + y^2 = r^2$$

$$\frac{x^2}{r^2} + \frac{y^2}{r^2} = \frac{r^2}{r^2}$$

Example 1

Verify $\cos^2 \theta + \sin^2 \theta = 1$




$$x^2 + y^2 = r^2$$

$$\frac{x^2}{r^2} + \frac{y^2}{r^2} = \frac{r^2}{r^2}$$

$$\left(\frac{x}{r}\right)^2 + \left(\frac{y}{r}\right)^2 = 1$$

Example 1

Verify $\cos^2 \theta + \sin^2 \theta = 1$



$$x^2 + y^2 = r^2$$


$$\frac{x^2}{r^2} + \frac{y^2}{r^2} = \frac{r^2}{r^2}$$

$$\left(\frac{x}{r}\right)^2 + \left(\frac{y}{r}\right)^2 = 1$$

$$(\cos \theta)^2 + (\sin \theta)^2 = 1$$

Example 1

Verify $\cos^2 \theta + \sin^2 \theta = 1$



$$x^2 + y^2 = r^2$$

$$\frac{x^2}{r^2} + \frac{y^2}{r^2} = \frac{r^2}{r^2}$$


$$\left(\frac{x}{r}\right)^2 + \left(\frac{y}{r}\right)^2 = 1$$

$$(\cos \theta)^2 + (\sin \theta)^2 = 1$$

$$\cos^2 \theta + \sin^2 \theta = 1$$

Example 2


Verify $1 + \tan^2 \theta = \sec^2 \theta$



Example 2


Verify $1 + \tan^2 \theta = \sec^2 \theta$

$$x^2 + y^2 = r^2$$



Example 2

Verify $1 + \tan^2 \theta = \sec^2 \theta$




$$x^2 + y^2 = r^2$$

$$\frac{x^2}{x^2} + \frac{y^2}{x^2} = \frac{r^2}{x^2}$$

Example 2

Verify $1 + \tan^2 \theta = \sec^2 \theta$




$$x^2 + y^2 = r^2$$

$$\frac{x^2}{x^2} + \frac{y^2}{x^2} = \frac{r^2}{x^2}$$

$$1 + \left(\frac{y}{x}\right)^2 = \left(\frac{r}{x}\right)^2$$

Example 2

Verify $1 + \tan^2 \theta = \sec^2 \theta$



$$x^2 + y^2 = r^2$$


$$\frac{x^2}{x^2} + \frac{y^2}{x^2} = \frac{r^2}{x^2}$$

$$1 + \left(\frac{y}{x}\right)^2 = \left(\frac{r}{x}\right)^2$$

$$1 + (\tan \theta)^2 = (\sec \theta)^2$$

Example 2

Verify $1 + \tan^2 \theta = \sec^2 \theta$



$$x^2 + y^2 = r^2$$

$$\frac{x^2}{x^2} + \frac{y^2}{x^2} = \frac{r^2}{x^2}$$


$$1 + \left(\frac{y}{x}\right)^2 = \left(\frac{r}{x}\right)^2$$

$$1 + (\tan \theta)^2 = (\sec \theta)^2$$

$$1 + \tan^2 \theta = \sec^2 \theta$$

Example 3


Verify $\cot^2 \theta + 1 = \csc^2 \theta$



Example 3

Verify $\cot^2 \theta + 1 = \csc^2 \theta$

$$x^2 + y^2 = r^2$$




Example 3

Verify $\cot^2 \theta + 1 = \csc^2 \theta$


$$x^2 + y^2 = r^2$$

$$\frac{x^2}{y^2} + \frac{y^2}{y^2} = \frac{r^2}{y^2}$$



Example 3

Verify $\cot^2 \theta + 1 = \csc^2 \theta$




$$x^2 + y^2 = r^2$$

$$\frac{x^2}{y^2} + \frac{y^2}{y^2} = \frac{r^2}{y^2}$$

$$\left(\frac{x}{y}\right)^2 + 1 = \left(\frac{r}{y}\right)^2$$

Example 3

Verify $\cot^2 \theta + 1 = \csc^2 \theta$



$$x^2 + y^2 = r^2$$


$$\frac{x^2}{y^2} + \frac{y^2}{y^2} = \frac{r^2}{y^2}$$

$$\left(\frac{x}{y}\right)^2 + 1 = \left(\frac{r}{y}\right)^2$$

$$(\cot \theta)^2 + 1 = (\csc \theta)^2$$

Example 3

Verify $\cot^2 \theta + 1 = \csc^2 \theta$



$$x^2 + y^2 = r^2$$

$$\frac{x^2}{y^2} + \frac{y^2}{y^2} = \frac{r^2}{y^2}$$

$$\left(\frac{x}{y}\right)^2 + 1 = \left(\frac{r}{y}\right)^2$$

$$(\cot \theta)^2 + 1 = (\csc \theta)^2$$

$$\cot^2 \theta + 1 = \csc^2 \theta$$

Alternate Forms of Pythagorean Identities

$$\cos^2 \theta + \sin^2 \theta = 1$$

Alternate Forms of Pythagorean Identities

$$\cos^2 \theta + \sin^2 \theta = 1$$

- $1 - \sin^2 \theta = \cos^2 \theta$

Alternate Forms of Pythagorean Identities

$$\cos^2 \theta + \sin^2 \theta = 1$$

- $1 - \sin^2 \theta = \cos^2 \theta$
 - $(1 - \sin \theta)(1 + \sin \theta) = \cos^2 \theta$

Alternate Forms of Pythagorean Identities

$$\cos^2 \theta + \sin^2 \theta = 1$$

- $1 - \sin^2 \theta = \cos^2 \theta$
 - $(1 - \sin \theta)(1 + \sin \theta) = \cos^2 \theta$
- $1 - \cos^2 \theta = \sin^2 \theta$

Alternate Forms of Pythagorean Identities

$$\cos^2 \theta + \sin^2 \theta = 1$$

- $1 - \sin^2 \theta = \cos^2 \theta$
 - $(1 - \sin \theta)(1 + \sin \theta) = \cos^2 \theta$
- $1 - \cos^2 \theta = \sin^2 \theta$
 - $(1 + \cos \theta)(1 - \cos \theta) = \sin^2 \theta$

Alternate Forms of Pythagorean Identities

$$1 + \tan^2 \theta = \sec^2 \theta$$

Alternate Forms of Pythagorean Identities

$$1 + \tan^2 \theta = \sec^2 \theta$$

- $\sec^2 \theta - 1 = \tan^2 \theta$

Alternate Forms of Pythagorean Identities

$$1 + \tan^2 \theta = \sec^2 \theta$$

- $\sec^2 \theta - 1 = \tan^2 \theta$
 - $(\sec \theta + 1)(\sec \theta - 1)$

Alternate Forms of Pythagorean Identities

$$1 + \tan^2 \theta = \sec^2 \theta$$

- $\sec^2 \theta - 1 = \tan^2 \theta$
 - $(\sec \theta + 1)(\sec \theta - 1)$
- $1 = \sec^2 \theta - \tan^2 \theta$

Alternate Forms of Pythagorean Identities

$$1 + \tan^2 \theta = \sec^2 \theta$$

- $\sec^2 \theta - 1 = \tan^2 \theta$
 - $(\sec \theta + 1)(\sec \theta - 1)$
- $1 = \sec^2 \theta - \tan^2 \theta$
 - $1 = (\sec \theta + \tan \theta)(\sec \theta - \tan \theta)$

Alternate Forms of Pythagorean Identities

$$1 + \cot^2 \theta = \csc^2 \theta$$

Alternate Forms of Pythagorean Identities

$$1 + \cot^2 \theta = \csc^2 \theta$$

- $\csc^2 \theta - \cot^2 \theta = 1$

Alternate Forms of Pythagorean Identities

$$1 + \cot^2 \theta = \csc^2 \theta$$

- $\csc^2 \theta - \cot^2 \theta = 1$

- $(\csc \theta + \cot \theta)(\csc \theta - \cot \theta) = 1$

Alternate Forms of Pythagorean Identities

$$1 + \cot^2 \theta = \csc^2 \theta$$

- $\csc^2 \theta - \cot^2 \theta = 1$
 - $(\csc \theta + \cot \theta)(\csc \theta - \cot \theta) = 1$
- $\csc^2 \theta - 1 = \cot^2 \theta$

Alternate Forms of Pythagorean Identities

$$1 + \cot^2 \theta = \csc^2 \theta$$

- $\csc^2 \theta - \cot^2 \theta = 1$
 - $(\csc \theta + \cot \theta)(\csc \theta - \cot \theta) = 1$
- $\csc^2 \theta - 1 = \cot^2 \theta$
 - $(\csc \theta + 1)(\csc \theta - 1) = \cot^2 \theta$

Objectives

- 1 Derive the Pythagorean identities
- 2 Derive the quotient identities
- 3 Use the sum and difference identities to verify other identities
- 4 Derive double-angle identities
- 5 Derive power-reducing identities


Quotient Identities

The Quotient Identities are

$$\tan \theta = \frac{\sin \theta}{\cos \theta} \quad \text{and} \quad \cot \theta = \frac{\cos \theta}{\sin \theta}$$


Example 4

Use the triangle to verify $\tan \theta = \frac{\sin \theta}{\cos \theta}$



Example 4


Use the triangle to verify $\tan \theta = \frac{\sin \theta}{\cos \theta}$



$$\frac{\sin \theta}{\cos \theta} = \frac{\left(\frac{y}{r}\right)}{\left(\frac{x}{r}\right)}$$

Example 4


Use the triangle to verify $\tan \theta = \frac{\sin \theta}{\cos \theta}$



$$\frac{\sin \theta}{\cos \theta} = \frac{\left(\frac{y}{r}\right)}{\left(\frac{x}{r}\right)} \left(\frac{r}{r}\right)$$

Example 4


Use the triangle to verify $\tan \theta = \frac{\sin \theta}{\cos \theta}$



$$\begin{aligned}\frac{\sin \theta}{\cos \theta} &= \frac{\left(\frac{y}{r}\right)}{\left(\frac{x}{r}\right)} \left(\frac{r}{r}\right) \\ &= \frac{y}{x}\end{aligned}$$

Example 4

Use the triangle to verify $\tan \theta = \frac{\sin \theta}{\cos \theta}$



$$\begin{aligned}\frac{\sin \theta}{\cos \theta} &= \frac{\left(\frac{y}{r}\right)}{\left(\frac{x}{r}\right)} \left(\frac{r}{r}\right) \\ &= \frac{y}{x} \\ &= \tan \theta\end{aligned}$$

Objectives

- 1 Derive the Pythagorean identities
- 2 Derive the quotient identities
- 3 Use the sum and difference identities to verify other identities
- 4 Derive double-angle identities
- 5 Derive power-reducing identities

Additional Angle Sum and Difference Identities

$$\cos(A - B) = \cos A \cos B + \sin A \sin B$$

$$\cos(A + B) = \cos A \cos B - \sin A \sin B$$

$$\sin(A - B) = \sin A \cos B - \sin B \cos A$$

$$\sin(A + B) = \sin A \cos B + \sin B \cos A$$

$$\tan(A - B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$$

$$\tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$$

$$** \text{ Note: } \tan(A \pm B) = \frac{\sin(A \pm B)}{\cos(A \pm B)}$$

Example 5

Verify $\cos\left(\frac{\pi}{2} - \theta\right) = \sin \theta$

Example 5

Verify $\cos\left(\frac{\pi}{2} - \theta\right) = \sin \theta$

$$\cos\left(\frac{\pi}{2} - \theta\right) = \cos \frac{\pi}{2} \cdot \cos \theta + \sin \frac{\pi}{2} \cdot \sin \theta$$

Example 5

Verify $\cos\left(\frac{\pi}{2} - \theta\right) = \sin \theta$

$$\begin{aligned}\cos\left(\frac{\pi}{2} - \theta\right) &= \cos \frac{\pi}{2} \cdot \cos \theta + \sin \frac{\pi}{2} \cdot \sin \theta \\ &= 0 \cos \theta + 1 \sin \theta\end{aligned}$$

Example 5

Verify $\cos\left(\frac{\pi}{2} - \theta\right) = \sin \theta$

$$\cos\left(\frac{\pi}{2} - \theta\right) = \cos \frac{\pi}{2} \cdot \cos \theta + \sin \frac{\pi}{2} \cdot \sin \theta$$

$$= 0 \cos \theta + 1 \sin \theta$$

$$= \sin \theta$$

Example 6

Verify $\cos(-\theta) = \cos \theta$

Example 6

Verify $\cos(-\theta) = \cos \theta$

$$\cos(-\theta) = \cos(0 - \theta)$$

Example 6

Verify $\cos(-\theta) = \cos \theta$

$$\cos(-\theta) = \cos(0 - \theta)$$

$$= \cos 0 \cdot \cos \theta + \sin 0 \cdot \sin \theta$$

Example 6

Verify $\cos(-\theta) = \cos \theta$

$$\cos(-\theta) = \cos(0 - \theta)$$

$$= \cos 0 \cdot \cos \theta + \sin 0 \cdot \sin \theta$$

$$= 1 \cos \theta + 0 \sin \theta$$

Example 6

Verify $\cos(-\theta) = \cos \theta$

$$\begin{aligned}\cos(-\theta) &= \cos(0 - \theta) \\ &= \cos 0 \cdot \cos \theta + \sin 0 \cdot \sin \theta \\ &= 1 \cos \theta + 0 \sin \theta \\ &= \cos \theta\end{aligned}$$

Objectives

- 1 Derive the Pythagorean identities
- 2 Derive the quotient identities
- 3 Use the sum and difference identities to verify other identities
- 4 Derive double-angle identities**
- 5 Derive power-reducing identities

Double-Angle Identities

The double-angle identities are an extension of the **angle sum identities**.

In this case, the angles would be equal.

Double-Angle Identities

The double-angle identities are an extension of the **angle sum identities**.

In this case, the angles would be equal.

$$\sin(2A) = \sin(A + A)$$

Double-Angle Identities

The double-angle identities are an extension of the **angle sum identities**.

In this case, the angles would be equal.

$$\begin{aligned}\sin(2A) &= \sin(A + A) \\ &= \sin A \cdot \cos A + \sin A \cdot \cos A\end{aligned}$$

Double-Angle Identities

The double-angle identities are an extension of the **angle sum identities**.

In this case, the angles would be equal.

$$\begin{aligned}\sin(2A) &= \sin(A + A) \\ &= \sin A \cdot \cos A + \sin A \cdot \cos A \\ &= 2 \sin A \cos A\end{aligned}$$

Double-Angle Identities

$$\sin(2A) = 2 \sin A \cos A$$

$$\cos(2A) = \cos^2 A - \sin^2 A$$

$$\tan(2A) = \frac{2 \tan A}{1 - \tan^2 A}$$

Alternate Forms of $\cos(2A)$

$\cos(2A)$ has two alternate forms.

Alternate form # 1:

Alternate Forms of $\cos(2A)$

$\cos(2A)$ has two alternate forms.

Alternate form # 1:

$$\cos(2A) = \cos^2 A - \sin^2 A$$

Alternate Forms of $\cos(2A)$

$\cos(2A)$ has two alternate forms.

Alternate form # 1:

$$\begin{aligned}\cos(2A) &= \cos^2 A - \sin^2 A \\ &= 1 - \sin^2 A - \sin^2 A\end{aligned}$$

Alternate Forms of $\cos(2A)$

$\cos(2A)$ has two alternate forms.

Alternate form # 1:

$$\begin{aligned}\cos(2A) &= \cos^2 A - \sin^2 A \\ &= 1 - \sin^2 A - \sin^2 A \\ &= 1 - 2\sin^2 A\end{aligned}$$

Alternate Forms of $\cos(2A)$

Alternate form # 2:

Alternate Forms of $\cos(2A)$

Alternate form # 2:

$$\cos(2A) = \cos^2 A - \sin^2 A$$

Alternate Forms of $\cos(2A)$

Alternate form # 2:

$$\begin{aligned}\cos(2A) &= \cos^2 A - \sin^2 A \\ &= \cos^2 A - (1 - \cos^2 A)\end{aligned}$$

Alternate Forms of $\cos(2A)$

Alternate form # 2:

$$\cos(2A) = \cos^2 A - \sin^2 A$$

$$= \cos^2 A - (1 - \cos^2 A)$$

$$= \cos^2 A - 1 + \cos^2 A$$

Alternate Forms of $\cos(2A)$

Alternate form # 2:

$$\begin{aligned}\cos(2A) &= \cos^2 A - \sin^2 A \\ &= \cos^2 A - (1 - \cos^2 A) \\ &= \cos^2 A - 1 + \cos^2 A \\ &= 2\cos^2 A - 1\end{aligned}$$

Objectives

- 1 Derive the Pythagorean identities
- 2 Derive the quotient identities
- 3 Use the sum and difference identities to verify other identities
- 4 Derive double-angle identities
- 5 Derive power-reducing identities

Power-Reduction Identities

The power-reduction identities can be derived from the alternate equations for $\cos(2A)$.

Power-Reduction Identities

The power-reduction identities can be derived from the alternate equations for $\cos(2A)$.

For $\cos(2A) = 2 \cos^2 A - 1$, if we solve for $\cos^2 A$, we get

$$\cos^2 A = \frac{1 + \cos(2A)}{2}$$

Power-Reduction Identities

The power-reduction identities can be derived from the alternate equations for $\cos(2A)$.

For $\cos(2A) = 2 \cos^2 A - 1$, if we solve for $\cos^2 A$, we get

$$\cos^2 A = \frac{1 + \cos(2A)}{2}$$

And for $\cos(2A) = 1 - 2 \sin^2 A$, solving for $\sin^2 A$ gives us

$$\sin^2 A = \frac{1 - \cos(2A)}{2}$$