

# Inverse Trig Functions

# Objectives

- 1 Find the values of inverse trig functions
- 2 Find the values of the compositions of inverse trig functions
- 3 Write an algebraic expression for the compositions of inverse trig functions

# Trig Functions and the Horizontal Line Test


Recall that for a function to have an **inverse**, any horizontal line can only hit the function at most once.

# Trig Functions and the Horizontal Line Test

Recall that for a function to have an **inverse**, any horizontal line can only hit the function at most once.

This creates a problem with periodic functions such as  $y = \sin x$ .

Graph of  $y = \sin x$  and  $y = \frac{1}{2}$




# Passing the Horizontal Line Test

If we limit the graph to a smaller domain, then the Horizontal Line Test will demonstrate the function  $y = \sin x$  will have an inverse:

# Passing the Horizontal Line Test

If we limit the graph to a smaller domain, then the Horizontal Line Test will demonstrate the function  $y = \sin x$  will have an inverse:



# Graph of $y = \sin^{-1} x$

The inverse of  $y = \sin x$  is


$$y = \sin^{-1} x \quad \text{or} \quad y = \arcsin x$$



# Graph of $y = \sin^{-1} x$

The inverse of  $y = \sin x$  is

$$y = \sin^{-1} x \quad \text{or} \quad y = \arcsin x$$



# Domains and Ranges for Inverse Trig Functions

<b>Inverse Function</b>	<b>Domain (Input)</b>	<b>Range (Output)</b>
-------------------------	-----------------------	-----------------------

---

# Domains and Ranges for Inverse Trig Functions

Inverse Function	Domain (Input)	Range (Output)
$y = \sin^{-1} x$	$[-1, 1]$	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$

# Domains and Ranges for Inverse Trig Functions

Inverse Function	Domain (Input)	Range (Output)
$y = \sin^{-1} x$	$[-1, 1]$	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$
$y = \cos^{-1} x$	$[-1, 1]$	$[0, \pi]$

# Domains and Ranges for Inverse Trig Functions

Inverse Function	Domain (Input)	Range (Output)
$y = \sin^{-1} x$	$[-1, 1]$	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$
$y = \cos^{-1} x$	$[-1, 1]$	$[0, \pi]$
$y = \tan^{-1} x$	$(-\infty, \infty)$	$\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$

# Domains and Ranges for Inverse Trig Functions

Inverse Function	Domain (Input)	Range (Output)
$y = \sin^{-1} x$	$[-1, 1]$	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$
$y = \cos^{-1} x$	$[-1, 1]$	$[0, \pi]$
$y = \tan^{-1} x$	$(-\infty, \infty)$	$\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$
$y = \csc^{-1} x$	$(-\infty, -1] \cup [1, \infty)$	$\left[-\frac{\pi}{2}, 0\right) \cup \left(0, \frac{\pi}{2}\right]$

# Domains and Ranges for Inverse Trig Functions

Inverse Function	Domain (Input)	Range (Output)
$y = \sin^{-1} x$	$[-1, 1]$	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$
$y = \cos^{-1} x$	$[-1, 1]$	$[0, \pi]$
$y = \tan^{-1} x$	$(-\infty, \infty)$	$\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$
$y = \csc^{-1} x$	$(-\infty, -1] \cup [1, \infty)$	$\left[-\frac{\pi}{2}, 0\right) \cup \left(0, \frac{\pi}{2}\right]$
$y = \sec^{-1} x$	$(-\infty, -1] \cup [1, \infty)$	$\left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$

# Domains and Ranges for Inverse Trig Functions

Inverse Function	Domain (Input)	Range (Output)
$y = \sin^{-1} x$	$[-1, 1]$	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$
$y = \cos^{-1} x$	$[-1, 1]$	$[0, \pi]$
$y = \tan^{-1} x$	$(-\infty, \infty)$	$\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$
$y = \csc^{-1} x$	$(-\infty, -1] \cup [1, \infty)$	$\left[-\frac{\pi}{2}, 0\right) \cup \left(0, \frac{\pi}{2}\right]$
$y = \sec^{-1} x$	$(-\infty, -1] \cup [1, \infty)$	$\left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$
$y = \cot^{-1} x$	$(-\infty, \infty)$	$(0, \pi)$



## Example 1


Find the exact value of each.

(a)  $\sin^{-1}\left(\frac{\sqrt{2}}{2}\right)$

## Example 1

Find the exact value of each.


(a)  $\sin^{-1}\left(\frac{\sqrt{2}}{2}\right)$



## Example 1

Find the exact value of each.

(a)  $\sin^{-1}\left(\frac{\sqrt{2}}{2}\right)$




$$\theta = \sin^{-1}\left(\frac{\sqrt{2}}{2}\right)$$

## Example 1

Find the exact value of each.

(a)  $\sin^{-1}\left(\frac{\sqrt{2}}{2}\right)$




$$\theta = \sin^{-1}\left(\frac{\sqrt{2}}{2}\right)$$

$$\theta = 45^\circ$$

## Example 1

Find the exact value of each.

(a)  $\sin^{-1}\left(\frac{\sqrt{2}}{2}\right)$



$$\theta = \sin^{-1}\left(\frac{\sqrt{2}}{2}\right)$$

$$\theta = 45^\circ$$

$$\theta = \frac{\pi}{4}$$

## Example 1


Find the exact value of each.

$$(b) \quad \cos^{-1} \left( -\frac{\sqrt{3}}{2} \right)$$

## Example 1

Find the exact value of each.


(b)  $\cos^{-1}\left(-\frac{\sqrt{3}}{2}\right)$



## Example 1

Find the exact value of each.

(b)  $\cos^{-1}\left(-\frac{\sqrt{3}}{2}\right)$




$$\theta = \cos^{-1}\left(\frac{-\sqrt{3}}{2}\right)$$



## Example 1

Find the exact value of each.

(b)  $\cos^{-1}\left(-\frac{\sqrt{3}}{2}\right)$




$$\theta = \cos^{-1}\left(-\frac{\sqrt{3}}{2}\right)$$

$$\theta = -150^\circ$$

## Example 1

Find the exact value of each.

$$(b) \quad \cos^{-1} \left( -\frac{\sqrt{3}}{2} \right)$$



$$\theta = \cos^{-1} \left( -\frac{\sqrt{3}}{2} \right)$$

$$\theta = -150^\circ$$

$$\theta = -\frac{5\pi}{6}$$

## Example 1


Find the exact value of each.

(c)  $\arctan(\sqrt{3})$

## Example 1

Find the exact value of each.


(c)  $\arctan(\sqrt{3})$



## Example 1

Find the exact value of each.

(c)  $\arctan(\sqrt{3})$




$$\theta = \arctan(\sqrt{3})$$

## Example 1

Find the exact value of each.

(c)  $\arctan(\sqrt{3})$




$$\theta = \arctan(\sqrt{3})$$

$$\theta = 60^\circ$$

## Example 1

Find the exact value of each.

(c)  $\arctan(\sqrt{3})$



$$\theta = \arctan(\sqrt{3})$$

$$\theta = 60^\circ$$

$$\theta = \frac{\pi}{3}$$

## Example 1

Find the exact value of each.


(d)  $\sec^{-1}(2)$



## Example 1

Find the exact value of each.


(d)  $\sec^{-1}(2)$



## Example 1

Find the exact value of each.

(d)  $\sec^{-1}(2)$




$$\theta = \sec^{-1}(2)$$

## Example 1

Find the exact value of each.

(d)  $\sec^{-1}(2)$




$$\theta = \sec^{-1}(2)$$

$$\theta = 60^\circ$$

## Example 1

Find the exact value of each.

(d)  $\sec^{-1}(2)$



$$\theta = \sec^{-1}(2)$$

$$\theta = 60^\circ$$

$$\theta = \frac{\pi}{3}$$

# Objectives

- 1 Find the values of inverse trig functions
- 2 Find the values of the compositions of inverse trig functions
- 3 Write an algebraic expression for the compositions of inverse trig functions

# Compositions of Inverse Trig Functions

When dealing with problems like these, it helps to sketch a right triangle, much like in the previous example.

You may need to use the Pythagorean Theorem.

## Example 2


Find the exact values of each (if possible).

(a)  $\cos\left(\cos^{-1}\left(\frac{1}{3}\right)\right)$

## Example 2

Find the exact values of each (if possible).

(a)  $\cos\left(\cos^{-1}\left(\frac{1}{3}\right)\right)$






## Example 2

Find the exact values of each (if possible).

(a)  $\cos\left(\cos^{-1}\left(\frac{1}{3}\right)\right)$




$$\theta = \cos^{-1}\left(\frac{1}{3}\right)$$

## Example 2

Find the exact values of each (if possible).

(a)  $\cos\left(\cos^{-1}\left(\frac{1}{3}\right)\right)$




$$\theta = \cos^{-1}\left(\frac{1}{3}\right)$$

$$\cos \theta = \frac{x}{r}$$

## Example 2

Find the exact values of each (if possible).

(a)  $\cos\left(\cos^{-1}\left(\frac{1}{3}\right)\right)$



$$\theta = \cos^{-1}\left(\frac{1}{3}\right)$$

$$\cos \theta = \frac{x}{r}$$


$$\cos \theta = \frac{1}{3}$$

## Example 2

$$(b) \quad \sin \left( \arcsin \left( -\frac{3}{7} \right) \right)$$


## Example 2

$$(b) \quad \sin \left( \arcsin \left( -\frac{3}{7} \right) \right)$$



## Example 2


$$(b) \quad \sin \left( \arcsin \left( -\frac{3}{7} \right) \right)$$



$$\theta = \arcsin \left( -\frac{3}{7} \right)$$

## Example 2

$$(b) \quad \sin \left( \arcsin \left( -\frac{3}{7} \right) \right)$$




$$\theta = \arcsin \left( -\frac{3}{7} \right)$$

$$\sin \theta = \frac{y}{r}$$

## Example 2

$$(b) \quad \sin \left( \arcsin \left( -\frac{3}{7} \right) \right)$$



$$\theta = \arcsin \left( -\frac{3}{7} \right)$$

$$\sin \theta = \frac{y}{r}$$

$$\sin \theta = -\frac{3}{7}$$




## Example 2

$$(c) \quad \cos \left( \arccos \left( \frac{5}{4} \right) \right)$$


## Example 2

$$(c) \quad \cos \left( \arccos \left( \frac{5}{4} \right) \right)$$



## Example 2


$$(c) \quad \cos \left( \arccos \left( \frac{5}{4} \right) \right)$$



$$\theta = \arccos \left( \frac{5}{4} \right)$$

## Example 2

(c)  $\cos\left(\arccos\left(\frac{5}{4}\right)\right)$




$$\theta = \arccos\left(\frac{5}{4}\right)$$

Domain error.

## Example 2

(c)  $\cos\left(\arccos\left(\frac{5}{4}\right)\right)$



$$\theta = \arccos\left(\frac{5}{4}\right)$$

Domain error.


$\emptyset$

## Example 2

$$(d) \quad \cos \left( \arctan \left( \frac{5}{12} \right) \right)$$

## Example 2


$$(d) \quad \cos \left( \arctan \left( \frac{5}{12} \right) \right)$$



## Example 2

$$(d) \quad \cos \left( \arctan \left( \frac{5}{12} \right) \right)$$


$$\theta = \arctan \left( \frac{5}{12} \right)$$





## Example 2

$$(d) \quad \cos \left( \arctan \left( \frac{5}{12} \right) \right)$$




$$\theta = \arctan \left( \frac{5}{12} \right)$$

$$12^2 + 5^2 = r^2$$

## Example 2

$$(d) \quad \cos \left( \arctan \left( \frac{5}{12} \right) \right)$$




$$\theta = \arctan \left( \frac{5}{12} \right)$$

$$12^2 + 5^2 = r^2$$

$$r = \sqrt{169} = 13$$

## Example 2

$$(d) \quad \cos \left( \arctan \left( \frac{5}{12} \right) \right)$$



$$\theta = \arctan \left( \frac{5}{12} \right)$$

$$12^2 + 5^2 = r^2$$

$$r = \sqrt{169} = 13$$


$$\cos \theta = \frac{12}{13}$$

## Example 2

$$(e) \quad \sin \left( \arctan \left( \frac{3}{4} \right) \right)$$

## Example 2


$$(e) \quad \sin \left( \arctan \left( \frac{3}{4} \right) \right)$$



## Example 2


(e)  $\sin \left( \arctan \left( \frac{3}{4} \right) \right)$

$$\theta = \arctan \left( \frac{3}{4} \right)$$



## Example 2

$$(e) \quad \sin \left( \arctan \left( \frac{3}{4} \right) \right)$$




$$\theta = \arctan \left( \frac{3}{4} \right)$$

$$3^2 + 4^2 = r^2$$

## Example 2

$$(e) \quad \sin \left( \arctan \left( \frac{3}{4} \right) \right)$$



$$\theta = \arctan \left( \frac{3}{4} \right)$$


$$3^2 + 4^2 = r^2$$

$$r = \sqrt{25} = 5$$



## Example 2

$$(e) \quad \sin \left( \arctan \left( \frac{3}{4} \right) \right)$$



$$\theta = \arctan \left( \frac{3}{4} \right)$$

$$3^2 + 4^2 = r^2$$

$$r = \sqrt{25} = 5$$


$$\sin \theta = \frac{3}{5}$$

## Example 2

$$(f) \quad \cot \left( \arcsin \left( -\frac{1}{3} \right) \right)$$

## Example 2


$$(f) \quad \cot \left( \arcsin \left( -\frac{1}{3} \right) \right)$$



## Example 2


$$(f) \quad \cot \left( \arcsin \left( -\frac{1}{3} \right) \right)$$

$$\theta = \arcsin \left( -\frac{1}{3} \right)$$



## Example 2

$$(f) \quad \cot \left( \arcsin \left( -\frac{1}{3} \right) \right)$$




$$\theta = \arcsin \left( -\frac{1}{3} \right)$$

$$x^2 + 1^2 = 3^2$$

## Example 2

$$(f) \quad \cot \left( \arcsin \left( -\frac{1}{3} \right) \right)$$




$$\theta = \arcsin \left( -\frac{1}{3} \right)$$

$$x^2 + 1^2 = 3^2$$

$$x = \sqrt{8} = 2\sqrt{2}$$

## Example 2

$$(f) \quad \cot \left( \arcsin \left( -\frac{1}{3} \right) \right)$$



$$\theta = \arcsin \left( -\frac{1}{3} \right)$$


$$x^2 + 1^2 = 3^2$$

$$x = \sqrt{8} = 2\sqrt{2}$$

$$\cot \theta = \frac{x}{y}$$

## Example 2

$$(f) \quad \cot \left( \arcsin \left( -\frac{1}{3} \right) \right)$$



$$\theta = \arcsin \left( -\frac{1}{3} \right)$$

$$x^2 + 1^2 = 3^2$$

$$x = \sqrt{8} = 2\sqrt{2}$$

$$\cot \theta = \frac{x}{y}$$

$$\cot \theta = \frac{2\sqrt{2}}{-1} = -2\sqrt{2}$$




## Example 2

$$(g) \quad \sec \left( \arccos \left( \frac{3}{4} \right) \right)$$


## Example 2

$$(g) \quad \sec \left( \arccos \left( \frac{3}{4} \right) \right)$$



## Example 2


$$(g) \quad \sec \left( \arccos \left( \frac{3}{4} \right) \right)$$



$$\theta = \arccos \left( \frac{3}{4} \right)$$

## Example 2

$$(g) \quad \sec \left( \arccos \left( \frac{3}{4} \right) \right)$$




$$\theta = \arccos \left( \frac{3}{4} \right)$$

$$\sec \theta = \frac{r}{x}$$

## Example 2

$$(g) \quad \sec \left( \arccos \left( \frac{3}{4} \right) \right)$$



$$\theta = \arccos \left( \frac{3}{4} \right)$$

$$\sec \theta = \frac{r}{x}$$

$$\sec \theta = \frac{4}{3}$$

# Objectives

- 1 Find the values of inverse trig functions
- 2 Find the values of the compositions of inverse trig functions
- 3 Write an algebraic expression for the compositions of inverse trig functions

# Algebraic Inverse Trig

In calculus, inverse trigonometric functions will be expressed algebraically (i.e. in terms of  $x$ ).

Set up a right triangle and use the **Pythagorean Theorem** to write the missing side in terms of  $x$ .

## Example 3

Write each as an algebraic expression of  $x$ .


(a)  $\cos(\sin^{-1} x)$



## Example 3

Write each as an algebraic expression of  $x$ .


(a)  $\cos(\sin^{-1} x)$



## Example 3

Write each as an algebraic expression of  $x$ .

(a)  $\cos(\sin^{-1} x)$




$$\theta = \arcsin(x)$$

## Example 3

Write each as an algebraic expression of  $x$ .

(a)  $\cos(\sin^{-1} x)$




$$\theta = \arcsin(x)$$

$$a^2 + x^2 = 1^2$$

## Example 3

Write each as an algebraic expression of  $x$ .

(a)  $\cos(\sin^{-1} x)$



$$\theta = \arcsin(x)$$


$$a^2 + x^2 = 1^2$$

$$a = \sqrt{1 - x^2}$$

## Example 3

Write each as an algebraic expression of  $x$ .

(a)  $\cos(\sin^{-1} x)$



$$\theta = \arcsin(x)$$

$$a^2 + x^2 = 1^2$$


$$a = \sqrt{1 - x^2}$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}}$$

## Example 3

Write each as an algebraic expression of  $x$ .

(a)  $\cos(\sin^{-1} x)$



$$\theta = \arcsin(x)$$

$$a^2 + x^2 = 1^2$$

$$a = \sqrt{1-x^2}$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}}$$


$$\cos \theta = \frac{\sqrt{1-x^2}}{1} = \sqrt{1-x^2}$$

## Example 3

(b)  $\sec(\tan^{-1} x)$

## Example 3


(b)  $\sec(\tan^{-1} x)$





## Example 3


(b)  $\sec(\tan^{-1} x)$



$$\theta = \arctan(x)$$

## Example 3

(b)  $\sec(\tan^{-1} x)$




$$\theta = \arctan(x)$$

$$1^2 + x^2 = c^2$$

## Example 3

(b)  $\sec(\tan^{-1} x)$




$$\theta = \arctan(x)$$

$$1^2 + x^2 = c^2$$

$$c = \sqrt{1 + x^2}$$

## Example 3

(b)  $\sec(\tan^{-1} x)$



$$\theta = \arctan(x)$$


$$1^2 + x^2 = c^2$$

$$c = \sqrt{1 + x^2}$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}}$$

## Example 3

(b)  $\sec(\tan^{-1} x)$



$$\theta = \arctan(x)$$

$$1^2 + x^2 = c^2$$

$$c = \sqrt{1 + x^2}$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}}$$


$$\sec \theta = \frac{\sqrt{1 + x^2}}{1} = \sqrt{1 + x^2}$$

## Example 3

(c)  $\tan(\sin^{-1}(2x))$


## Example 3

(c)  $\tan(\sin^{-1}(2x))$



## Example 3

(c)  $\tan(\sin^{-1}(2x))$




$$\theta = \arcsin(2x)$$



## Example 3

(c)  $\tan(\sin^{-1}(2x))$




$$\theta = \arcsin(2x)$$

$$a^2 + (2x)^2 = 1^2$$

## Example 3

(c)  $\tan(\sin^{-1}(2x))$




$$\theta = \arcsin(2x)$$

$$a^2 + (2x)^2 = 1^2$$

$$a^2 + 4x^2 = 1$$

## Example 3

(c)  $\tan(\sin^{-1}(2x))$



$$\theta = \arcsin(2x)$$


$$a^2 + (2x)^2 = 1^2$$

$$a^2 + 4x^2 = 1$$

$$a = \sqrt{1 - 4x^2}$$


## Example 3

(c)  $\tan(\sin^{-1}(2x))$



## Example 3


(c)  $\tan(\sin^{-1}(2x))$



$$\tan \theta = \frac{\text{opp}}{\text{adj}}$$

## Example 3

(c)  $\tan(\sin^{-1}(2x))$




$$\tan \theta = \frac{\text{opp}}{\text{adj}}$$

$$\tan \theta = \frac{2x}{\sqrt{1 - 4x^2}}$$

## Example 3

(c)  $\tan(\sin^{-1}(2x))$



$$\tan \theta = \frac{\text{opp}}{\text{adj}}$$

$$\tan \theta = \frac{2x}{\sqrt{1 - 4x^2}}$$

$$\tan \theta = \frac{2x\sqrt{1 - 4x^2}}{1 - 4x^2}$$


## Example 3

$$(d) \quad \sin \left( \cot^{-1} \left( \frac{x}{5} \right) \right)$$




## Example 3

(d)  $\sin \left( \cot^{-1} \left( \frac{x}{5} \right) \right)$



## Example 3


(d)  $\sin \left( \cot^{-1} \left( \frac{x}{5} \right) \right)$



$$\theta = \cot^{-1} \left( \frac{x}{5} \right)$$

## Example 3

(d)  $\sin \left( \cot^{-1} \left( \frac{x}{5} \right) \right)$




$$\theta = \cot^{-1} \left( \frac{x}{5} \right)$$

$$5^2 + x^2 = c^2$$

## Example 3

(d)  $\sin \left( \cot^{-1} \left( \frac{x}{5} \right) \right)$




$$\theta = \cot^{-1} \left( \frac{x}{5} \right)$$

$$5^2 + x^2 = c^2$$

$$c = \sqrt{25 + x^2}$$


## Example 3

(d)  $\sin \left( \cot^{-1} \left( \frac{x}{5} \right) \right)$



## Example 3


(d)  $\sin \left( \cot^{-1} \left( \frac{x}{5} \right) \right)$



$$\sec \theta = \frac{\text{hyp}}{\text{adj}}$$

## Example 3

(d)  $\sin \left( \cot^{-1} \left( \frac{x}{5} \right) \right)$



$$\sec \theta = \frac{\text{hyp}}{\text{adj}}$$

$$\sec \theta = \frac{\sqrt{25 + x^2}}{x}$$