

# Inverse Functions

# Intro

An inverse function works like an “undo” function.

# Intro

An inverse function works like an “undo” function.

If we give a function input it gives us output in return.

# Intro

An inverse function works like an “undo” function.


If we give a function input it gives us output in return.

If we want our original input back, we can give our output to the inverse function and get our original input back.

# Visual Interpretation

Original Function  $f(x) = x^2$

Inverse Function:  $g(x) = \sqrt{x}$



# Mathematical Definition of Inverse Functions

Mathematically, two functions  $f(x)$  and  $g(x)$  are inverses of each other if **both** of the following are true:

# Mathematical Definition of Inverse Functions

Mathematically, two functions  $f(x)$  and  $g(x)$  are inverses of each other if **both** of the following are true:

- $(g \circ f)(x) = x$  for all  $x$  in the domain of  $f$

# Mathematical Definition of Inverse Functions

Mathematically, two functions  $f(x)$  and  $g(x)$  are inverses of each other if **both** of the following are true:

- $(g \circ f)(x) = x$  for all  $x$  in the domain of  $f$
- $(f \circ g)(x) = x$  for all  $x$  in the domain of  $g$



# Objectives

- 1 Determine if a relation or function has an inverse
- 2 Find the inverse of a function
- 3 Find the domain and range of the inverse of a function

# Horizontal Line Test

When we first looked at relations, we knew they were functions if they passed the **Vertical Line Test**.

# Horizontal Line Test

When we first looked at relations, we knew they were functions if they passed the **Vertical Line Test**.

For a function to have an inverse that is also a function, it must pass a similar test known as the **Horizontal Line Test**.

# Horizontal Line Test

When we first looked at relations, we knew they were functions if they passed the **Vertical Line Test**.

For a function to have an inverse that is also a function, it must pass a similar test known as the **Horizontal Line Test**.

## Horizontal Line Test

If each horizontal line intersects the graph **at most once**, then the function has an inverse.

# Horizontal Line Test

When we first looked at relations, we knew they were functions if they passed the **Vertical Line Test**.

For a function to have an inverse that is also a function, it must pass a similar test known as the **Horizontal Line Test**.

## Horizontal Line Test

If each horizontal line intersects the graph at most once, then the function has an inverse.

*Note:* A function that passes the Horizontal Line Test is also said to be **invertible** or **one-to-one**.

## Example 1


Determine if the following relations or functions have an inverse.

(a)  $F = \{(-1, 1), (0, 2), (2, 1)\}$

## Example 1

Determine if the following relations or functions have an inverse.


(a)  $F = \{(-1, 1), (0, 2), (2, 1)\}$



## Example 1

Determine if the following relations or functions have an inverse.

(a)  $F = \{(-1, 1), (0, 2), (2, 1)\}$






## Example 1

Determine if the following relations or functions have an inverse.

(a)  $F = \{(-1, 1), (0, 2), (2, 1)\}$




Does not have an inverse.

## Example 1

$$(b) \quad f(x) = \frac{1 - 2x}{5}$$


## Example 1

$$(b) \quad f(x) = \frac{1 - 2x}{5}$$




## Example 1

$$(b) \quad f(x) = \frac{1 - 2x}{5}$$



## Example 1

(b)  $f(x) = \frac{1 - 2x}{5}$




Has an inverse.

## Example 1

$$(c) \quad g(x) = \frac{2x}{1-x}$$


## Example 1

(c)  $g(x) = \frac{2x}{1-x}$



## Example 1


(c)  $g(x) = \frac{2x}{1-x}$





## Example 1

(c)  $g(x) = \frac{2x}{1-x}$




Has an inverse.

## Example 1

$$(d) \quad h(x) = x^2 - 2x + 4$$


## Example 1

(d)  $h(x) = x^2 - 2x + 4$




## Example 1

(d)  $h(x) = x^2 - 2x + 4$



## Example 1

(d)  $h(x) = x^2 - 2x + 4$



Does not have an inverse.

# Objectives

- 1 Determine if a relation or function has an inverse
- 2 Find the inverse of a function
- 3 Find the domain and range of the inverse of a function

# Inverse Function Notation

For a given function  $f(x)$ , the notation for the inverse of  $f(x)$  is

$$f^{-1}(x)$$

# How to Find the Inverse of a Function

- 1 Write as  $y = f(x)$



# How to Find the Inverse of a Function

- 1 Write as  $y = f(x)$
- 2 Switch the  $x$  and  $y$

# How to Find the Inverse of a Function

- 1 Write as  $y = f(x)$
- 2 Switch the  $x$  and  $y$
- 3 Solve for  $y$  and write using inverse notation  $f^{-1}(x)$ .

## Example 2

Find the inverse of each.

(a)  $f(x) = \frac{1 - 2x}{5}$

## Example 2

Find the inverse of each.

$$(a) \quad f(x) = \frac{1 - 2x}{5}$$

$$y = \frac{1 - 2x}{5}$$

## Example 2

Find the inverse of each.

$$(a) \quad f(x) = \frac{1 - 2x}{5}$$

$$y = \frac{1 - 2x}{5}$$

$$x = \frac{1 - 2y}{5}$$

## Example 2

Find the inverse of each.

$$(a) \quad f(x) = \frac{1 - 2x}{5}$$

$$y = \frac{1 - 2x}{5}$$

$$x = \frac{1 - 2y}{5}$$

$$5x = 1 - 2y$$

## Example 2

Find the inverse of each.

$$(a) \quad f(x) = \frac{1 - 2x}{5}$$

$$y = \frac{1 - 2x}{5}$$

$$x = \frac{1 - 2y}{5}$$

$$5x = 1 - 2y$$

$$-2y = 5x - 1$$

## Example 2

Find the inverse of each.

$$(a) \quad f(x) = \frac{1 - 2x}{5}$$

$$y = \frac{1 - 2x}{5}$$

$$x = \frac{1 - 2y}{5}$$

$$5x = 1 - 2y$$

$$-2y = 5x - 1$$

$$y = \frac{5x - 1}{-2}$$



## Example 2

$$f^{-1}(x) = \frac{-5x + 1}{2}$$

## Example 2

$$(b) \quad g(x) = \frac{2x}{1-x}$$

## Example 2

$$(b) \quad g(x) = \frac{2x}{1-x}$$

$$y = \frac{2x}{1-x}$$

## Example 2

$$(b) \quad g(x) = \frac{2x}{1-x}$$

$$y = \frac{2x}{1-x}$$

$$x = \frac{2y}{1-y}$$

## Example 2

$$(b) \quad g(x) = \frac{2x}{1-x}$$

$$y = \frac{2x}{1-x}$$

$$x = \frac{2y}{1-y}$$

$$x(1-y) = 2y$$

## Example 2

$$(b) \quad g(x) = \frac{2x}{1-x}$$

$$y = \frac{2x}{1-x}$$

$$x = \frac{2y}{1-y}$$

$$x(1-y) = 2y$$

$$x - xy = 2y$$

## Example 2

$$(b) \quad g(x) = \frac{2x}{1-x}$$

$$y = \frac{2x}{1-x}$$

$$x = \frac{2y}{1-y}$$

$$x(1-y) = 2y$$

$$x - xy = 2y$$

$$x = 2y + xy$$

## Example 2

$$(b) \quad g(x) = \frac{2x}{1-x}$$

$$y = \frac{2x}{1-x}$$

$$x = \frac{2y}{1-y}$$

$$x(1-y) = 2y$$

$$x - xy = 2y$$

$$x = 2y + xy$$

$$x = y(2 + x)$$



Example 2  $g(x) = \frac{2x}{1-x}$

$$\frac{x}{2+x} = y$$

Example 2  $g(x) = \frac{2x}{1-x}$

$$\frac{x}{2+x} = y$$

$$g^{-1}(x) = \frac{x}{x+2}$$

## Example 2

$$(c) \quad h(x) = \sqrt{x+2} - 1$$

## Example 2

$$(c) \quad h(x) = \sqrt{x+2} - 1$$

$$y = \sqrt{x+2} - 1$$

## Example 2

$$(c) \quad h(x) = \sqrt{x+2} - 1$$

$$y = \sqrt{x+2} - 1$$

$$x = \sqrt{\textcolor{red}{y}+2} - 1$$

## Example 2

$$(c) \quad h(x) = \sqrt{x+2} - 1$$

$$y = \sqrt{x+2} - 1$$

$$x = \sqrt{\textcolor{red}{y}+2} - 1$$

$$x+1 = \sqrt{\textcolor{red}{y}+2}$$

## Example 2

$$(c) \quad h(x) = \sqrt{x+2} - 1$$

$$y = \sqrt{x+2} - 1$$

$$x = \sqrt{y+2} - 1$$

$$x + 1 = \sqrt{y+2}$$

$$(x + 1)^2 = y + 2$$

## Example 2

$$(c) \quad h(x) = \sqrt{x+2} - 1$$

$$y = \sqrt{x+2} - 1$$

$$x = \sqrt{y+2} - 1$$

$$x + 1 = \sqrt{y+2}$$

$$(x + 1)^2 = y + 2$$

$$(x + 1)^2 - 2 = y$$



## Example 2

$$(c) \quad h(x) = \sqrt{x+2} - 1$$

$$y = \sqrt{x+2} - 1$$

$$x = \sqrt{y+2} - 1$$

$$x + 1 = \sqrt{y+2}$$

$$(x + 1)^2 = y + 2$$

$$(x + 1)^2 - 2 = y$$

$$h^{-1}(x) = (x + 1)^2 - 2$$

# Objectives

- 1 Determine if a relation or function has an inverse
- 2 Find the inverse of a function
- 3 Find the domain and range of the inverse of a function

# Domain and Range of Inverse Functions

In the previous example, we found the inverse of a function by switching the  $x$  and the  $y$  in the equation.

# Domain and Range of Inverse Functions

In the previous example, we found the inverse of a function by switching the  $x$  and the  $y$  in the equation.

By doing this, we also switch the **domain and range** of the original function and its inverse.

# Domain and Range of Inverse Functions

In the previous example, we found the inverse of a function by switching the  $x$  and the  $y$  in the equation.

By doing this, we also switch the **domain and range** of the original function and its inverse.


	Domain	Range
$f(x)$		
$f^{-1}(x)$		

# Domain and Range of Inverse Functions

In the previous example, we found the inverse of a function by switching the  $x$  and the  $y$  in the equation.

By doing this, we also switch the **domain and range** of the original function and its inverse.

	Domain	Range
$f(x)$		
$f^{-1}(x)$		




# Domain and Range of Inverse Functions

In the previous example, we found the inverse of a function by switching the  $x$  and the  $y$  in the equation.

By doing this, we also switch the **domain and range** of the original function and its inverse.

	Domain	Range
$f(x)$		
$f^{-1}(x)$		




# Visual Interpretation of Finding Inverse

Switching the  $x$  and  $y$  variables when finding the inverse of a function involves reflecting the graph of the function along the line  $y = x$ .



# Visual Interpretation of Finding Inverse

Switching the  $x$  and  $y$  variables when finding the inverse of a function involves **reflecting the graph of the function along the line  $y = x$** .



# Domain and Range Restrictions

Sometimes you have to restrict the domain of the function so that it is a reflection of its inverse across the line  $y = x$

## Example 3

Find the domain and range of the inverse of each.

(a)  $f(x) = \frac{1 - 2x}{5}$

## Example 3

Find the domain and range of the inverse of each.

$$(a) \quad f(x) = \frac{1 - 2x}{5} \quad f^{-1}(x) = \frac{-5x + 1}{2}$$

## Example 3

Find the domain and range of the inverse of each.

$$(a) \quad f(x) = \frac{1-2x}{5} \quad f^{-1}(x) = \frac{-5x+1}{2}$$

	Domain	Range
$f(x)$		
$f^{-1}(x)$		

## Example 3

Find the domain and range of the inverse of each.

$$(a) \quad f(x) = \frac{1 - 2x}{5} \quad f^{-1}(x) = \frac{-5x + 1}{2}$$

	Domain	Range
$f(x)$	$(-\infty, \infty)$	
$f^{-1}(x)$		

## Example 3

Find the domain and range of the inverse of each.

$$(a) \quad f(x) = \frac{1-2x}{5} \quad f^{-1}(x) = \frac{-5x+1}{2}$$

	Domain	Range
$f(x)$	$(-\infty, \infty)$	
$f^{-1}(x)$		$(-\infty, \infty)$

## Example 3

Find the domain and range of the inverse of each.

$$(a) \quad f(x) = \frac{1-2x}{5} \quad f^{-1}(x) = \frac{-5x+1}{2}$$

	Domain	Range
$f(x)$	$(-\infty, \infty)$	$(-\infty, \infty)$
$f^{-1}(x)$		$(-\infty, \infty)$



## Example 3

Find the domain and range of the inverse of each.

$$(a) \quad f(x) = \frac{1-2x}{5} \quad f^{-1}(x) = \frac{-5x+1}{2}$$

	Domain	Range
$f(x)$	$(-\infty, \infty)$	$(-\infty, \infty)$
$f^{-1}(x)$	$(-\infty, \infty)$	$(-\infty, \infty)$

## Example 3

Find the domain and range of the inverse of each.

$$(a) \quad f(x) = \frac{1-2x}{5} \quad f^{-1}(x) = \frac{-5x+1}{2}$$

	Domain	Range
$f(x)$	$(-\infty, \infty)$	$(-\infty, \infty)$
$f^{-1}(x)$	$(-\infty, \infty)$	$(-\infty, \infty)$

Domain of  $f^{-1}(x)$  :  $(-\infty, \infty)$

## Example 3

Find the domain and range of the inverse of each.

$$(a) \quad f(x) = \frac{1-2x}{5} \quad f^{-1}(x) = \frac{-5x+1}{2}$$

	Domain	Range
$f(x)$	$(-\infty, \infty)$	$(-\infty, \infty)$
$f^{-1}(x)$	$(-\infty, \infty)$	$(-\infty, \infty)$

$$\text{Domain of } f^{-1}(x) : (-\infty, \infty)$$

$$\text{Range of } f^{-1}(x) : (-\infty, \infty)$$

## Example 3

$$(b) \quad g(x) = \frac{2x}{1-x}$$

## Example 3

$$(b) \quad g(x) = \frac{2x}{1-x} \quad g^{-1}(x) = \frac{x}{x+2}$$

## Example 3

$$(b) \quad g(x) = \frac{2x}{1-x} \quad g^{-1}(x) = \frac{x}{x+2}$$

	Domain	Range
$g(x)$		
$g^{-1}(x)$		

## Example 3

$$(b) \quad g(x) = \frac{2x}{1-x} \quad g^{-1}(x) = \frac{x}{x+2}$$

	Domain	Range
$g(x)$	$x \neq 1$	
$g^{-1}(x)$		

## Example 3

$$(b) \quad g(x) = \frac{2x}{1-x} \quad g^{-1}(x) = \frac{x}{x+2}$$

	Domain	Range
$g(x)$	$x \neq 1$	
$g^{-1}(x)$		$y \neq 1$



## Example 3

$$(b) \quad g(x) = \frac{2x}{1-x} \quad g^{-1}(x) = \frac{x}{x+2}$$

	Domain	Range
$g(x)$	$x \neq 1$	
$g^{-1}(x)$	$x \neq -2$	$y \neq 1$

## Example 3

$$(b) \quad g(x) = \frac{2x}{1-x} \quad g^{-1}(x) = \frac{x}{x+2}$$

	Domain	Range
$g(x)$	$x \neq 1$	$y \neq -2$
$g^{-1}(x)$	$x \neq -2$	$y \neq 1$

## Example 3

$$(b) \quad g(x) = \frac{2x}{1-x} \quad g^{-1}(x) = \frac{x}{x+2}$$

	Domain	Range
$g(x)$	$x \neq 1$	$y \neq -2$
$g^{-1}(x)$	$x \neq -2$	$y \neq 1$

Domain of  $g^{-1}(x)$  :  $(-\infty, -2) \cup (-2, \infty)$

## Example 3

$$(b) \quad g(x) = \frac{2x}{1-x} \quad g^{-1}(x) = \frac{x}{x+2}$$

	Domain	Range
$g(x)$	$x \neq 1$	$y \neq -2$
$g^{-1}(x)$	$x \neq -2$	$y \neq 1$

Domain of  $g^{-1}(x)$  :  $(-\infty, -2) \cup (-2, \infty)$

Range of  $g^{-1}(x)$  :  $(-\infty, 1) \cup (1, \infty)$

## Example 3

$$(c) \quad h(x) = \sqrt{x+2} - 1$$

## Example 3

$$(c) \quad h(x) = \sqrt{x+2} - 1 \quad h^{-1}(x) = (x+1)^2 - 2$$

## Example 3

(c)  $h(x) = \sqrt{x+2} - 1$      $h^{-1}(x) = (x+1)^2 - 2$

	Domain	Range
$h(x)$		
$h^{-1}(x)$		

## Example 3

(c)  $h(x) = \sqrt{x+2} - 1$      $h^{-1}(x) = (x+1)^2 - 2$

	Domain	Range
$h(x)$	$x \geq -2$	
$h^{-1}(x)$		



## Example 3

(c)  $h(x) = \sqrt{x+2} - 1$      $h^{-1}(x) = (x+1)^2 - 2$

	Domain	Range
$h(x)$	$x \geq -2$	
$h^{-1}(x)$		$y \geq -2$

## Example 3

(c)  $h(x) = \sqrt{x+2} - 1$      $h^{-1}(x) = (x+1)^2 - 2$

	Domain	Range
$h(x)$	$x \geq -2$	$y \geq -1$
$h^{-1}(x)$		$y \geq -2$

## Example 3

(c)  $h(x) = \sqrt{x+2} - 1$      $h^{-1}(x) = (x+1)^2 - 2$

	Domain	Range
$h(x)$	$x \geq -2$	$y \geq -1$
$h^{-1}(x)$	$x \geq -1$	$y \geq -2$

## Example 3

$$(c) \quad h(x) = \sqrt{x+2} - 1 \quad h^{-1}(x) = (x+1)^2 - 2$$

	Domain	Range
$h(x)$	$x \geq -2$	$y \geq -1$
$h^{-1}(x)$	$x \geq -1$	$y \geq -2$

Domain of  $h^{-1}(x)$  :  $[-1, \infty)$

## Example 3

(c)  $h(x) = \sqrt{x+2} - 1$      $h^{-1}(x) = (x+1)^2 - 2$

	Domain	Range
$h(x)$	$x \geq -2$	$y \geq -1$
$h^{-1}(x)$	$x \geq -1$	$y \geq -2$

Domain of  $h^{-1}(x)$  :  $[-1, \infty)$


Range of  $h^{-1}(x)$  :  $[-2, \infty)$

## Example 3

$$(d) \quad j(x) = x^2 - 2x + 4, \quad x \leq 1$$


## Example 3

(d)  $j(x) = x^2 - 2x + 4, \quad x \leq 1$



## Example 3

(d)  $j(x) = x^2 - 2x + 4, \quad x \leq 1$





## Example 3

$$(d) \quad j(x) = x^2 - 2x + 4 \quad x \leq 1$$

## Example 3

$$(d) \quad j(x) = x^2 - 2x + 4 \quad x \leq 1$$

$$y = x^2 - 2x + 4$$

## Example 3

$$(d) \quad j(x) = x^2 - 2x + 4 \quad x \leq 1$$

$$y = x^2 - 2x + 4$$

$$x = y^2 - 2y + 4$$

## Example 3

$$(d) \quad j(x) = x^2 - 2x + 4 \quad x \leq 1$$

$$y = x^2 - 2x + 4$$

$$x = y^2 - 2y + 4$$

$$x = (y - 1)^2 + 3$$

## Example 3

$$(d) \quad j(x) = x^2 - 2x + 4 \quad x \leq 1$$

$$y = x^2 - 2x + 4$$

$$x = y^2 - 2y + 4$$

$$x = (y - 1)^2 + 3$$

$$x - 3 = (y - 1)^2$$

## Example 3

$$(d) \quad j(x) = x^2 - 2x + 4 \quad x \leq 1$$

$$y = x^2 - 2x + 4$$

$$x = y^2 - 2y + 4$$

$$x = (y - 1)^2 + 3$$

$$x - 3 = (y - 1)^2$$

$$\pm \sqrt{x - 3} = y - 1$$

## Example 3

$$(d) \quad j(x) = x^2 - 2x + 4 \quad x \leq 1$$

$$y = x^2 - 2x + 4$$

$$x = y^2 - 2y + 4$$

$$x = (y - 1)^2 + 3$$

$$x - 3 = (y - 1)^2$$

$$\pm \sqrt{x - 3} = y - 1$$

$$1 \pm \sqrt{x - 3} = y$$

## Example 3

$$(d) \quad j(x) = x^2 - 2x + 4 \quad x \leq 1$$

$$y = x^2 - 2x + 4$$

$$x = y^2 - 2y + 4$$

$$x = (y - 1)^2 + 3$$

$$x - 3 = (y - 1)^2$$

$$\pm \sqrt{x - 3} = y - 1$$

$$1 \pm \sqrt{x - 3} = y$$

$$j^{-1}(x) = 1 \pm \sqrt{x - 3}$$



## Example 3

$$(d) \quad j(x) = x^2 - 2x + 4 \quad x \leq 1$$

$$y = x^2 - 2x + 4$$

$$x = y^2 - 2y + 4$$

$$x = (y - 1)^2 + 3$$

$$x - 3 = (y - 1)^2$$

$$\pm \sqrt{x - 3} = y - 1$$

$$1 \pm \sqrt{x - 3} = y$$

$$j^{-1}(x) = 1 \pm \sqrt{x - 3}$$

$$j^{-1}(x) = 1 - \sqrt{x - 3}$$

## Example 3

$$(d) \quad j(x) = x^2 - 2x + 4 \quad x \leq 1 \quad j^{-1}(x) = 1 - \sqrt{x - 3}$$

## Example 3

$$(d) \quad j(x) = x^2 - 2x + 4 \quad x \leq 1 \quad j^{-1}(x) = 1 - \sqrt{x - 3}$$

	Domain	Range
$j(x)$		
$j^{-1}(x)$		

## Example 3

$$(d) \quad j(x) = x^2 - 2x + 4 \quad x \leq 1 \quad j^{-1}(x) = 1 - \sqrt{x - 3}$$

	Domain	Range
$j(x)$	$x \leq 1$	
$j^{-1}(x)$		

## Example 3

$$(d) \quad j(x) = x^2 - 2x + 4 \quad x \leq 1 \quad j^{-1}(x) = 1 - \sqrt{x - 3}$$

	Domain	Range
$j(x)$	$x \leq 1$	
$j^{-1}(x)$		$y \leq 1$

## Example 3

$$(d) \quad j(x) = x^2 - 2x + 4 \quad x \leq 1 \quad j^{-1}(x) = 1 - \sqrt{x - 3}$$

	Domain	Range
$j(x)$	$x \leq 1$	
$j^{-1}(x)$	$x \geq 3$	$y \leq 1$

## Example 3

$$(d) \quad j(x) = x^2 - 2x + 4 \quad x \leq 1 \quad j^{-1}(x) = 1 - \sqrt{x - 3}$$

	Domain	Range
$j(x)$	$x \leq 1$	$y \geq 3$
$j^{-1}(x)$	$x \geq 3$	$y \leq 1$

## Example 3

$$(d) \quad j(x) = x^2 - 2x + 4 \quad x \leq 1 \quad j^{-1}(x) = 1 - \sqrt{x - 3}$$

	Domain	Range
$j(x)$	$x \leq 1$	$y \geq 3$
$j^{-1}(x)$	$x \geq 3$	$y \leq 1$

Domain of  $j^{-1}(x)$ :  $[3, \infty)$



## Example 3

$$(d) \quad j(x) = x^2 - 2x + 4 \quad x \leq 1 \quad j^{-1}(x) = 1 - \sqrt{x - 3}$$

	Domain	Range
$j(x)$	$x \leq 1$	$y \geq 3$
$j^{-1}(x)$	$x \geq 3$	$y \leq 1$

Domain of  $j^{-1}(x)$ :  $[3, \infty)$

Range of  $j^{-1}(x)$ :  $(-\infty, 1]$