

Rational Functions and Their Graphs

What is a Rational Function?

A **rational function** is a function in the form

$$\frac{p(x)}{q(x)}$$

where $p(x)$ and $q(x)$ are polynomials.

Objectives

- 1 Determine the end behavior of a rational function
- 2 Determine the equations of vertical asymptotes and coordinates of holes of a rational function

End Behavior

Recall from Polynomial Functions that **end behavior** refers to the graph's behavior as x approaches ∞ and also $-\infty$.

End Behavior

Recall from Polynomial Functions that **end behavior** refers to the graph's behavior as x approaches ∞ and also $-\infty$.

In other words, what are the output values approaching as our input values get larger (in either direction)?

Asymptotes

With rational functions, 1 of 2 possible scenarios for end behavior of the graph will happen with each function:

Asymptotes

With rational functions, 1 of 2 possible scenarios for end behavior of the graph will happen with each function:

- The function will look like a horizontal line.

Asymptotes

With rational functions, 1 of 2 possible scenarios for end behavior of the graph will happen with each function:

- The function will look like a horizontal line.
- The function will not look like a horizontal line.

Example 1

Determine the end behavior of the function

$$f(x) = \frac{2x}{x^2 - 3}$$

Example 1

Determine the end behavior of the function

$$f(x) = \frac{2x}{x^2 - 3}$$

x	$f(x)$
1,000	0.002
10,000	0.0002
100,000	0.00002
1,000,000	0.000002

Example 1

Determine the end behavior of the function

$$f(x) = \frac{2x}{x^2 - 3}$$

x	$f(x)$	x	$f(x)$
1,000	0.002	-1,000	-0.002
10,000	0.0002	-10,000	-0.0002
100,000	0.00002	-100,000	-0.00002
1,000,000	0.000002	-1,000,000	-0.000002

Example 1

Notice the outputs are getting closer and closer to 0.

Example 1

Notice the outputs are getting closer and closer to 0.

Thus, the end behavior of the rational function

$$f(x) = \frac{2x}{x^2 - 3}$$

is 0.

Example 1

Notice the outputs are getting closer and closer to 0.

Thus, the end behavior of the rational function

$$f(x) = \frac{2x}{x^2 - 3}$$

is 0.

As you zoom out of the function, the graph more and more resembles the graph of $y = 0$.

Example 1

Notice the outputs are getting closer and closer to 0.

Thus, the end behavior of the rational function

$$f(x) = \frac{2x}{x^2 - 3}$$

is 0.

As you zoom out of the function, the graph more and more resembles the graph of $y = 0$.

As far as notation goes, we would say

$$\lim_{x \rightarrow -\infty} f(x) = 0 \quad \text{and} \quad \lim_{x \rightarrow \infty} f(x) = 0$$

End Behavior

Determining end behavior is a result of using polynomial division.

End Behavior

Determining end behavior is a result of using **polynomial division**.

When your end behavior approaches an actual number (such as 0 in the last example), then the rational function has a **horizontal asymptote** as its end behavior.

End Behavior

Determining end behavior is a result of using **polynomial division**.

When your end behavior approaches an actual number (such as 0 in the last example), then the rational function has a **horizontal asymptote** as its end behavior.

This will happen if either

- Degree of numerator $<$ Degree of denominator (horizontal asymptote will be $y = 0$)

End Behavior

Determining end behavior is a result of using **polynomial division**.

When your end behavior approaches an actual number (such as 0 in the last example), then the rational function has a **horizontal asymptote** as its end behavior.

This will happen if either

- Degree of numerator $<$ Degree of denominator (horizontal asymptote will be $y = 0$)
- Degree of numerator $=$ Degree of denominator (horizontal asymptote will be ratio of leading coefficients)

Oblique Asymptotes

If the degree of the numerator $>$ degree of the denominator, the end behavior **will not be a horizontal line**.

Oblique Asymptotes

If the degree of the numerator $>$ degree of the denominator, the end behavior **will not be a horizontal line**.

Instead, it will be an **oblique (or slant) asymptote**.

Oblique Asymptotes

If the degree of the numerator $>$ degree of the denominator, the end behavior **will not be a horizontal line**.

Instead, it will be an **oblique (or slant) asymptote**.

You will need to use **polynomial division** to find the equation of an oblique asymptote.

Example 2

Determine the end behavior of each, then find the equation of the asymptote.

(a) $f(x) = \frac{5x}{x^2 + 1}$

Example 2

Determine the end behavior of each, then find the equation of the asymptote.

(a) $f(x) = \frac{5x}{x^2 + 1}$

- Degree of numerator $<$ Degree of denominator

Example 2

Determine the end behavior of each, then find the equation of the asymptote.

(a) $f(x) = \frac{5x}{x^2 + 1}$

- Degree of numerator $<$ Degree of denominator
- Horizontal asymptote

Example 2

Determine the end behavior of each, then find the equation of the asymptote.

(a) $f(x) = \frac{5x}{x^2 + 1}$

- Degree of numerator $<$ Degree of denominator
- Horizontal asymptote
- $y = 0$

Example 2

Note: If you used polynomial division, you would get

$$f(x) = 0 + \frac{5x}{x^2 + 1}$$

and as $x \rightarrow \pm\infty$,

$$f(x) \rightarrow 0 + 0 = 0$$

Example 2

$$(b) \quad g(x) = \frac{x^2 - 4}{x + 1}$$

Example 2

$$(b) \quad g(x) = \frac{x^2 - 4}{x + 1}$$

- Degree of numerator $>$ Degree of denominator

Example 2

$$(b) \quad g(x) = \frac{x^2 - 4}{x + 1}$$

- Degree of numerator $>$ Degree of denominator
- Oblique asymptote

Example 2 $g(x) = \frac{x^2-4}{x+1}$

$$(x^2 + 0x - 4) \div (x + 1)$$

x	x^2	
1		

Example 2 $g(x) = \frac{x^2-4}{x+1}$

$$(x^2 + 0x - 4) \div (x + 1)$$

	x	
x	x^2	
1		

Example 2 $g(x) = \frac{x^2-4}{x+1}$

$$(x^2 + 0x - 4) \div (x + 1)$$

	x	
x	x^2	
1	x	

Example 2 $g(x) = \frac{x^2-4}{x+1}$

$$(x^2 + 0x - 4) \div (x + 1)$$

		x
x	x^2	$-x$
1	x	

Example 2 $g(x) = \frac{x^2-4}{x+1}$

$$(x^2 + 0x - 4) \div (x + 1)$$

	x	-1
x	x^2	$-x$
1	x	

Example 2 $g(x) = \frac{x^2-4}{x+1}$

$$(x^2 + 0x - 4) \div (x + 1)$$

	x	-1
x	x^2	$-x$
1	x	-1

Example 2 $g(x) = \frac{x^2-4}{x+1}$

$$(x^2 + 0x - 4) \div (x + 1)$$

	x	-1
x	x^2	$-x$
1	x	-1

remainder: -3

Example 2 $g(x) = \frac{x^2-4}{x+1}$

$$(x^2 + 0x - 4) \div (x + 1)$$

	x	-1	
x	x^2	$-x$	
1	x	-1	remainder: -3

$$x - 1 - \frac{3}{x + 1}$$

Example 2 $g(x) = \frac{x^2-4}{x+1}$

$$(x^2 + 0x - 4) \div (x + 1)$$

	x	-1	
x	x^2	$-x$	
1	x	-1	

remainder: -3

$$x - 1 - \frac{3}{x + 1}$$

Slant asymptote: $y = x - 1$

Example 2

$$(c) \quad h(x) = \frac{6x^3 - 3x + 1}{5 - 2x^3}$$

Example 2

$$(c) \quad h(x) = \frac{6x^3 - 3x + 1}{5 - 2x^3}$$

$$h(x) = \frac{6x^3 - 3x + 1}{-2x^3 + 5}$$

Example 2

$$(c) \quad h(x) = \frac{6x^3 - 3x + 1}{5 - 2x^3}$$

$$h(x) = \frac{6x^3 - 3x + 1}{-2x^3 + 5}$$

- Degree of numerator = Degree of denominator

Example 2

$$(c) \quad h(x) = \frac{6x^3 - 3x + 1}{5 - 2x^3}$$

$$h(x) = \frac{6x^3 - 3x + 1}{-2x^3 + 5}$$

- Degree of numerator = Degree of denominator
- Horizontal asymptote

Example 2

$$(c) \quad h(x) = \frac{6x^3 - 3x + 1}{5 - 2x^3}$$

$$h(x) = \frac{6x^3 - 3x + 1}{-2x^3 + 5}$$

- Degree of numerator = Degree of denominator
- Horizontal asymptote
- Ratio of leading coefficients = $\frac{6}{-2} = -3$

Example 2

$$(c) \quad h(x) = \frac{6x^3 - 3x + 1}{5 - 2x^3}$$

$$h(x) = \frac{6x^3 - 3x + 1}{-2x^3 + 5}$$

- Degree of numerator = Degree of denominator
- Horizontal asymptote
- Ratio of leading coefficients = $\frac{6}{-2} = -3$
- $y = -3$

Example 2

$$h(x) = \frac{6x^3 - 3x + 1}{5 - 2x^3}$$

Note: If you perform polynomial division, you get

$$h(x) = -3 + \frac{-3x + 16}{5 - 2x^3}$$

And as $x \rightarrow \pm\infty$,

$$h(x) \rightarrow -3 + 0 = -3$$

Objectives

- 1 Determine the end behavior of a rational function
- 2 Determine the equations of vertical asymptotes and coordinates of holes of a rational function

Domain of Rational Functions

Recall that the **domain** of a rational function is all real numbers
*****EXCEPT***** values that make the **denominator equal 0**.

Domain of Rational Functions

Recall that the **domain** of a rational function is all real numbers *****EXCEPT***** values that make the **denominator equal 0**.

At each of the values that cause the denominator to $= 0$, there will be **only one of two things** there:

Domain of Rational Functions

Recall that the **domain** of a rational function is all real numbers *****EXCEPT***** values that make the **denominator equal 0**.

At each of the values that cause the denominator to $= 0$, there will be **only one of two things** there:

- A vertical asymptote

Domain of Rational Functions

Recall that the **domain** of a rational function is all real numbers *****EXCEPT***** values that make the **denominator equal 0**.

At each of the values that cause the denominator to $= 0$, there will be **only one of two things there**:

- A vertical asymptote
- Or a hole in the graph

Vertical Asymptotes


A **vertical asymptote** is a vertical line that the function will approach, but never intersect.

Vertical Asymptotes

A **vertical asymptote** is a vertical line that the function will approach, but never intersect.

As the graph of the function gets closer to a vertical asymptote, the graph will either skyrocket up towards ∞ or plummet downward towards $-\infty$.

Vertical Asymptotes



Holes in Rational Functions

Most graphing technology will not show holes in graphs upon sight.

How to Tell Your Asymptote From a Hole In the Graph

- 1 **Factor** both numerator and denominator ****completely**** and then **simplify that**.

How to Tell Your Asymptote From a Hole In the Graph

- 1 **Factor** both numerator and denominator ****completely**** and then **simplify that**.
- 2 For **each domain issue** in the denominator:

How to Tell Your Asymptote From a Hole In the Graph

- 1 **Factor** both numerator and denominator ****completely**** and then **simplify that**.
- 2 For **each domain issue** in the denominator:
 - If you get a 0 in the denominator after evaluating the simplified expression, you have a **vertical asymptote**.

How to Tell Your Asymptote From a Hole In the Graph

- 1 **Factor** both numerator and denominator ****completely**** and then **simplify that**.
- 2 For **each domain issue** in the denominator:
 - If you get a 0 in the denominator after evaluating the simplified expression, you have a **vertical asymptote**.
 - Otherwise, there is a hole in the graph there.

How to Tell Your Asymptote From a Hole In the Graph

- 1 **Factor** both numerator and denominator ****completely**** and then **simplify that**.
- 2 For **each domain issue** in the denominator:
 - If you get a 0 in the denominator after evaluating the simplified expression, you have a **vertical asymptote**.
 - Otherwise, there is a hole in the graph there.

To get the y -coordinate of a hole in the graph, plug in the value of x into your simplified expression.

Example 3

Determine the domain of each. Then determine the equations of vertical asymptotes and/or coordinates of any holes.

(a) $f(x) = \frac{2x}{x^2 - 9}$

Example 3

Determine the domain of each. Then determine the equations of vertical asymptotes and/or coordinates of any holes.

$$(a) \quad f(x) = \frac{2x}{x^2 - 9} = \frac{2x}{(x + 3)(x - 3)}$$

Example 3

Determine the domain of each. Then determine the equations of vertical asymptotes and/or coordinates of any holes.

$$(a) \quad f(x) = \frac{2x}{x^2 - 9} = \frac{2x}{(x + 3)(x - 3)}$$

Domain:

Example 3

Determine the domain of each. Then determine the equations of vertical asymptotes and/or coordinates of any holes.

$$(a) \quad f(x) = \frac{2x}{x^2 - 9} = \frac{2x}{(x + 3)(x - 3)}$$

Domain:

$$x + 3 \neq 0$$

$$x - 3 \neq 0$$

Example 3

Determine the domain of each. Then determine the equations of vertical asymptotes and/or coordinates of any holes.

$$(a) \quad f(x) = \frac{2x}{x^2 - 9} = \frac{2x}{(x + 3)(x - 3)}$$

Domain:

$$x + 3 \neq 0$$

$$x \neq -3$$

$$x - 3 \neq 0$$

$$x \neq 3$$

Example 3

Determine the domain of each. Then determine the equations of vertical asymptotes and/or coordinates of any holes.

$$(a) \quad f(x) = \frac{2x}{x^2 - 9} = \frac{2x}{(x + 3)(x - 3)}$$

Domain:

$$x + 3 \neq 0$$

$$x \neq -3$$

$$x - 3 \neq 0$$

$$x \neq 3$$

Domain: $x \neq -3, 3$

Example 3 Domain: $x \neq -3, 3$

$$f(x) = \frac{2x}{(x+3)(x-3)}$$

Example 3 Domain: $x \neq -3, 3$

$$f(x) = \frac{2x}{(x+3)(x-3)}$$

We get 0s in denominator if we evaluate

$$\frac{2x}{(x+3)(x-3)}$$

at $x = -3$ and $x = 3$

Example 3 Domain: $x \neq -3, 3$

$$f(x) = \frac{2x}{(x+3)(x-3)}$$

We get 0s in denominator if we evaluate

$$\frac{2x}{(x+3)(x-3)}$$

at $x = -3$ and $x = 3$

$x = -3$ and $x = 3$ are both **vertical asymptotes**

Example 3 Domain: $x \neq -3, 3$

$$f(x) = \frac{2x}{(x+3)(x-3)}$$

We get 0s in denominator if we evaluate

$$\frac{2x}{(x+3)(x-3)}$$

at $x = -3$ and $x = 3$

$x = -3$ and $x = 3$ are both **vertical asymptotes**

There are no holes in the graph

Example 3

$$(b) \quad g(x) = \frac{x^2 - 6x - 7}{x^2 + 5x + 4}$$

Example 3

$$(b) \quad g(x) = \frac{x^2 - 6x - 7}{x^2 + 5x + 4} = \frac{(x + 1)(x - 7)}{(x + 1)(x + 4)}$$

Example 3

$$(b) \quad g(x) = \frac{x^2 - 6x - 7}{x^2 + 5x + 4} = \frac{(x + 1)(x - 7)}{(x + 1)(x + 4)}$$

Domain:

Example 3

$$(b) \quad g(x) = \frac{x^2 - 6x - 7}{x^2 + 5x + 4} = \frac{(x + 1)(x - 7)}{(x + 1)(x + 4)}$$

Domain: $x \neq -4, -1$

Example 3

$$(b) \quad g(x) = \frac{x^2 - 6x - 7}{x^2 + 5x + 4} = \frac{(x+1)(x-7)}{(x+1)(x+4)} = \frac{x-7}{x+4}$$

Domain: $x \neq -4, -1$

Example 3

$$(b) \quad g(x) = \frac{x^2 - 6x - 7}{x^2 + 5x + 4} = \frac{(x+1)(x-7)}{(x+1)(x+4)} = \frac{x-7}{x+4}$$

Domain: $x \neq -4, -1$

Vertical asymptote at $x = -4$

Example 3

$$(b) \quad g(x) = \frac{x^2 - 6x - 7}{x^2 + 5x + 4} = \frac{(x+1)(x-7)}{(x+1)(x+4)} = \frac{x-7}{x+4}$$

Domain: $x \neq -4, -1$

Vertical asymptote at $x = -4$

Hole in graph at $x = -1$

Example 3

$$(b) \quad g(x) = \frac{x^2 - 6x - 7}{x^2 + 5x + 4} = \frac{(x+1)(x-7)}{(x+1)(x+4)} = \frac{x-7}{x+4}$$

Domain: $x \neq -4, -1$

Vertical asymptote at $x = -4$

Hole in graph at $x = -1$

$$\text{y-coordinate: } \frac{-1-7}{-1+4} = -\frac{8}{3}$$

Example 3

$$(b) \quad g(x) = \frac{x^2 - 6x - 7}{x^2 + 5x + 4} = \frac{(x+1)(x-7)}{(x+1)(x+4)} = \frac{x-7}{x+4}$$

Domain: $x \neq -4, -1$

Vertical asymptote at $x = -4$

Hole in graph at $x = -1$

$$\text{y-coordinate: } \frac{-1-7}{-1+4} = -\frac{8}{3}$$

Hole at $\left(-1, -\frac{8}{3}\right)$

Example 3

$$(c) \quad h(x) = \frac{x^2 - x - 6}{x^2 - 9}$$

Example 3

$$(c) \quad h(x) = \frac{x^2 - x - 6}{x^2 - 9} = \frac{(x - 3)(x + 2)}{(x - 3)(x + 3)}$$

Example 3

$$(c) \quad h(x) = \frac{x^2 - x - 6}{x^2 - 9} = \frac{(x - 3)(x + 2)}{(x - 3)(x + 3)}$$

Domain:

Example 3

$$(c) \quad h(x) = \frac{x^2 - x - 6}{x^2 - 9} = \frac{(x - 3)(x + 2)}{(x - 3)(x + 3)}$$

Domain: $x \neq -3, 3$

Example 3

$$(c) \quad h(x) = \frac{x^2 - x - 6}{x^2 - 9} = \frac{(x - 3)(x + 2)}{(x - 3)(x + 3)} = \frac{x + 2}{x + 3}$$

Domain: $x \neq -3, 3$

Example 3

$$(c) \quad h(x) = \frac{x^2 - x - 6}{x^2 - 9} = \frac{(x - 3)(x + 2)}{(x - 3)(x + 3)} = \frac{x + 2}{x + 3}$$

Domain: $x \neq -3, 3$

Vertical asymptote at $x = -3$

Example 3

$$(c) \quad h(x) = \frac{x^2 - x - 6}{x^2 - 9} = \frac{(x - 3)(x + 2)}{(x - 3)(x + 3)} = \frac{x + 2}{x + 3}$$

Domain: $x \neq -3, 3$

Vertical asymptote at $x = -3$

There is a hole in the graph at $x = 3$

Example 3

$$(c) \quad h(x) = \frac{x^2 - x - 6}{x^2 - 9} = \frac{(x - 3)(x + 2)}{(x - 3)(x + 3)} = \frac{x + 2}{x + 3}$$

Domain: $x \neq -3, 3$

Vertical asymptote at $x = -3$

There is a hole in the graph at $x = 3$

$$y\text{-coordinate: } \frac{3 + 2}{3 + 3} = \frac{5}{6}$$

Example 3

$$(c) \quad h(x) = \frac{x^2 - x - 6}{x^2 - 9} = \frac{(x - 3)(x + 2)}{(x - 3)(x + 3)} = \frac{x + 2}{x + 3}$$

Domain: $x \neq -3, 3$

Vertical asymptote at $x = -3$

There is a hole in the graph at $x = 3$

$$y\text{-coordinate: } \frac{3 + 2}{3 + 3} = \frac{5}{6}$$

Hole at $\left(3, \frac{5}{6}\right)$

Example 3

$$(d) \quad j(x) = \frac{x^2 - x - 6}{x^2 + 9}$$

Example 3

$$(d) \quad j(x) = \frac{x^2 - x - 6}{x^2 + 9} = \frac{(x - 3)(x + 2)}{x^2 + 9}$$

Example 3

$$(d) \quad j(x) = \frac{x^2 - x - 6}{x^2 + 9} = \frac{(x - 3)(x + 2)}{x^2 + 9}$$

Since $x^2 + 9 \neq 0$ for any real number x , domain is All Real Numbers

Example 3

$$(d) \quad j(x) = \frac{x^2 - x - 6}{x^2 + 9} = \frac{(x - 3)(x + 2)}{x^2 + 9}$$

Since $x^2 + 9 \neq 0$ for any real number x , domain is All Real Numbers

Since domain is all reals, there are no vertical asymptotes or holes

Example 3

$$(e) \quad k(x) = \frac{x^2 - x - 6}{x^2 + 4x + 4}$$

Example 3

$$(e) \quad k(x) = \frac{x^2 - x - 6}{x^2 + 4x + 4} = \frac{(x - 3)(x + 2)}{(x + 2)(x + 2)}$$

Example 3

$$(e) \quad k(x) = \frac{x^2 - x - 6}{x^2 + 4x + 4} = \frac{(x - 3)(x + 2)}{(x + 2)(x + 2)}$$

Domain:

Example 3

$$(e) \quad k(x) = \frac{x^2 - x - 6}{x^2 + 4x + 4} = \frac{(x - 3)(x + 2)}{(x + 2)(x + 2)}$$

Domain: $x \neq -2$

Example 3

$$(e) \quad k(x) = \frac{x^2 - x - 6}{x^2 + 4x + 4} = \frac{(x - 3)(x + 2)}{(x + 2)(x + 2)} = \frac{x - 3}{x + 2}$$

Domain: $x \neq -2$

Example 3

$$(e) \quad k(x) = \frac{x^2 - x - 6}{x^2 + 4x + 4} = \frac{(x - 3)(x + 2)}{(x + 2)(x + 2)} = \frac{x - 3}{x + 2}$$

Domain: $x \neq -2$

Vertical asymptote at $x = -2$

Example 3

$$(e) \quad k(x) = \frac{x^2 - x - 6}{x^2 + 4x + 4} = \frac{(x - 3)(x + 2)}{(x + 2)(x + 2)} = \frac{x - 3}{x + 2}$$

Domain: $x \neq -2$

Vertical asymptote at $x = -2$

There is no hole in the graph