

Vectors

Table of Contents


Intro

A **vector** is a mathematical object with both magnitude (length) and direction.

It is represented geometrically by a directed line segment.

Intro

The following shows an example of a vector $\vec{v} = \overrightarrow{PQ}$.



P is the **initial point** and Q is the **terminal point**.


Magnitude

The **magnitude** of a vector (denoted $\|\vec{v}\|$ or $|\vec{v}|$) is the distance between P and Q .

Think Pythagorean Theorem.


Component Form

If we connect the point P to point Q we get the following diagram:



Component Form

If we connect the point P to point Q we get the following diagram:



This can be represented by $\vec{v} = \langle 3, 4 \rangle$ or by $\mathbf{v} = 3\mathbf{i} + 4\mathbf{j}$.

Component Form

To find the component form for $P(x_0, y_0)$ and $Q(x_1, y_1)$, just take the difference in coordinates:

$$\vec{v} = \overrightarrow{PQ} = \langle x_1 - x_0, y_1 - y_0 \rangle$$

Component Form

To find the component form for $P(x_0, y_0)$ and $Q(x_1, y_1)$, just take the difference in coordinates:

$$\vec{v} = \overrightarrow{PQ} = \langle x_1 - x_0, y_1 - y_0 \rangle$$

or

$$(x_1 - x_0)\mathbf{i} + (y_1 - y_0)\mathbf{j}$$

Example 1a

Find the component form and exact magnitude of \overrightarrow{AB} for each.

(a) $A(2, -1) \quad B(-5, 3)$

Example 1a

Find the component form and exact magnitude of \overrightarrow{AB} for each.

$$\begin{aligned} \text{(a)} \quad A(2, -1) \quad B(-5, 3) \\ = \langle -5 - 2, 3 - (-1) \rangle \end{aligned}$$

Example 1a

Find the component form and exact magnitude of \overrightarrow{AB} for each.

(a) $A(2, -1) \quad B(-5, 3)$

$$= \langle -5 - 2, 3 - (-1) \rangle$$

$$= \langle -7, 4 \rangle$$

$$= -7\mathbf{i} + 4\mathbf{j}$$

Example 1a

Find the component form and exact magnitude of \overrightarrow{AB} for each.

$$(a) \quad A(2, -1) \quad B(-5, 3)$$

$$= \langle -5 - 2, 3 - (-1) \rangle$$

$$= \langle -7, 4 \rangle$$

$$= -7\mathbf{i} + 4\mathbf{j}$$

$$\|\overrightarrow{AB}\| = \sqrt{7^2 + 4^2}$$

Example 1a

Find the component form and exact magnitude of \overrightarrow{AB} for each.

$$(a) \quad A(2, -1) \quad B(-5, 3)$$

$$= \langle -5 - 2, 3 - (-1) \rangle$$

$$= \langle -7, 4 \rangle$$

$$= -7\mathbf{i} + 4\mathbf{j}$$

$$\|\overrightarrow{AB}\| = \sqrt{7^2 + 4^2}$$

$$= \sqrt{49 + 16}$$

Example 1a

Find the component form and exact magnitude of \overrightarrow{AB} for each.

$$(a) \quad A(2, -1) \quad B(-5, 3)$$

$$= \langle -5 - 2, 3 - (-1) \rangle$$

$$= \langle -7, 4 \rangle$$

$$= -7\mathbf{i} + 4\mathbf{j}$$

$$\|\overrightarrow{AB}\| = \sqrt{7^2 + 4^2}$$

$$= \sqrt{49 + 16}$$

$$= \sqrt{65}$$

Example 1b

$$(b) \quad A(-3, 2) \quad B(0, 1)$$

Example 1b

$$(b) \quad A(-3, 2) \quad B(0, 1)$$

$$= \langle 0 - (-3), 1 - 2 \rangle$$

Example 1b

$$(b) \quad A(-3, 2) \quad B(0, 1)$$

$$= \langle 0 - (-3), 1 - 2 \rangle$$

$$= \langle 3, -1 \rangle$$

$$= 3\mathbf{i} - \mathbf{j}$$

Example 1b

$$(b) \quad A(-3, 2) \quad B(0, 1)$$

$$= \langle 0 - (-3), 1 - 2 \rangle$$

$$= \langle 3, -1 \rangle$$

$$= 3\mathbf{i} - \mathbf{j}$$

$$\|\overrightarrow{AB}\| = \sqrt{3^2 + 1^2}$$

Example 1b

$$(b) \quad A(-3, 2) \quad B(0, 1)$$

$$= \langle 0 - (-3), 1 - 2 \rangle$$

$$= \langle 3, -1 \rangle$$

$$= 3\mathbf{i} - \mathbf{j}$$

$$\|\overrightarrow{AB}\| = \sqrt{3^2 + 1^2}$$

$$= \sqrt{10}$$

Table of Contents

Vector Addition

We can add vectors by adding the horizontal components, and adding the vertical components.

Think of these as *combining like terms*.

Vector Addition

For instance, when adding $\vec{v} = \langle 3, 4 \rangle$ and $\vec{w} = \langle 1, -2 \rangle$, combine horizontal components and vertical components:

Vector Addition

For instance, when adding $\vec{v} = \langle 3, 4 \rangle$ and $\vec{w} = \langle 1, -2 \rangle$, combine horizontal components and vertical components:

$$\vec{v} + \vec{w} = \langle 3 + 1, 4 + (-2) \rangle$$


Vector Addition

For instance, when adding $\vec{v} = \langle 3, 4 \rangle$ and $\vec{w} = \langle 1, -2 \rangle$, combine horizontal components and vertical components:

$$\begin{aligned}\vec{v} + \vec{w} &= \langle 3 + 1, 4 + (-2) \rangle \\ &= \langle 4, 2 \rangle\end{aligned}$$


Vector Addition

Visually, the previous problem is




Scalar Multiplication

If we multiply our vector by a real number (a **scalar**), we get a new vector.






Scalar Multiplication

If we multiply our vector by a real number (a **scalar**), we get a new vector.




Scalar Multiplication

If we multiply our vector by a real number (a **scalar**), we get a new vector.



Scalar Multiplication


If we multiply our vector by a real number (a **scalar**), we get a new vector.



Scalar multiplication “distributes” the scalar to both horizontal and vertical component.

Vector Subtraction

Vector subtraction $\vec{v} - \vec{w}$ can be thought of as $\vec{v} + (-\vec{w})$ and is illustrated below:



Example 2a

If $\mathbf{v} = 5\mathbf{i} + 4\mathbf{j}$, and $\vec{w} = \langle 6, -9 \rangle$, find each of the following.

(a) $\mathbf{v} + \mathbf{w}$

Example 2a

If $\mathbf{v} = 5\mathbf{i} + 4\mathbf{j}$, and $\vec{w} = \langle 6, -9 \rangle$, find each of the following.

(a) $\mathbf{v} + \mathbf{w}$

$$\mathbf{v} + \mathbf{w} = \langle 5, 4 \rangle + \langle 6, -9 \rangle$$

Example 2a

If $\mathbf{v} = 5\mathbf{i} + 4\mathbf{j}$, and $\vec{w} = \langle 6, -9 \rangle$, find each of the following.

(a) $\mathbf{v} + \mathbf{w}$

$$\begin{aligned}\mathbf{v} + \mathbf{w} &= \langle 5, 4 \rangle + \langle 6, -9 \rangle \\ &= \langle 5 + 6, 4 + (-9) \rangle\end{aligned}$$

Example 2a

If $\mathbf{v} = 5\mathbf{i} + 4\mathbf{j}$, and $\vec{w} = \langle 6, -9 \rangle$, find each of the following.

(a) $\mathbf{v} + \mathbf{w}$

$$\begin{aligned}\mathbf{v} + \mathbf{w} &= \langle 5, 4 \rangle + \langle 6, -9 \rangle \\ &= \langle 5 + 6, 4 + (-9) \rangle \\ &= \langle 11, -5 \rangle\end{aligned}$$

Example 2b

(b) $\mathbf{v} - \mathbf{w}$

Example 2b

(b) $\mathbf{v} - \mathbf{w}$

$$\mathbf{v} - \mathbf{w} = \langle 5, 4 \rangle - \langle 6, -9 \rangle$$

Example 2b

(b) $\mathbf{v} - \mathbf{w}$

$$\begin{aligned}\mathbf{v} - \mathbf{w} &= \langle 5, 4 \rangle - \langle 6, -9 \rangle \\ &= \langle 5 - 6, 4 - (-9) \rangle\end{aligned}$$

Example 2b

(b) $\mathbf{v} - \mathbf{w}$

$$\begin{aligned}\mathbf{v} - \mathbf{w} &= \langle 5, 4 \rangle - \langle 6, -9 \rangle \\ &= \langle 5 - 6, 4 - (-9) \rangle \\ &= \langle -1, 13 \rangle\end{aligned}$$

Example 2c

(c) $6\vec{v}$

Example 2c

(c) $6\vec{v}$

$$6\vec{v} = 6\langle 5, 4 \rangle$$

Example 2c

$$(c) \quad 6\vec{v}$$

$$\begin{aligned} 6\vec{v} &= 6\langle 5, 4 \rangle \\ &= \langle 30, 24 \rangle \end{aligned}$$

Example 2d

$$(d) \quad 2\mathbf{v} + 4\mathbf{w}$$

Example 2d

(d) $2\mathbf{v} + 4\mathbf{w}$

$$2\mathbf{v} + 4\mathbf{w} = 2(5\mathbf{i} + 4\mathbf{j}) + 4(6\mathbf{i} - 9\mathbf{j})$$

Example 2d

$$(d) \quad 2\mathbf{v} + 4\mathbf{w}$$

$$2\mathbf{v} + 4\mathbf{w} = 2(5\mathbf{i} + 4\mathbf{j}) + 4(6\mathbf{i} - 9\mathbf{j})$$

$$= 10\mathbf{i} + 8\mathbf{j} + 24\mathbf{i} - 36\mathbf{j}$$

Example 2d

$$(d) \quad 2\mathbf{v} + 4\mathbf{w}$$

$$2\mathbf{v} + 4\mathbf{w} = 2(5\mathbf{i} + 4\mathbf{j}) + 4(6\mathbf{i} - 9\mathbf{j})$$


$$= 10\mathbf{i} + 8\mathbf{j} + 24\mathbf{i} - 36\mathbf{j}$$

$$= 34\mathbf{i} - 28\mathbf{j}$$

Table of Contents

Unit Vectors

A **unit vector**, \hat{v} , is a vector that has a magnitude of 1.



Notice the unit vector \hat{v} is parallel to \vec{v} .

\mathbf{i} and \mathbf{j} are unit vectors with $\mathbf{i} = \langle 1, 0 \rangle$ and $\mathbf{j} = \langle 0, 1 \rangle$

Unit Vectors

We get \hat{v} by dividing the magnitude of \vec{v} by itself:

$$\hat{v} = \frac{\vec{v}}{\|\vec{v}\|}$$

Unit Vectors

We get \hat{v} by dividing the magnitude of \vec{v} by itself:

$$\hat{v} = \frac{\vec{v}}{\|\vec{v}\|}$$

Since we are dividing the vector (the hypotenuse) by the magnitude, we also divide the x and y components as well:

$$\hat{v} = \left\langle \frac{x}{\|\vec{v}\|}, \frac{y}{\|\vec{v}\|} \right\rangle$$

Example 3a

Find a unit vector for each given vector.

(a) $\mathbf{v} = 5\mathbf{i} - 12\mathbf{j}$

Example 3a

Find a unit vector for each given vector.

(a) $\mathbf{v} = 5\mathbf{i} - 12\mathbf{j}$

$$\|\mathbf{v}\| = \sqrt{5^2 + 12^2}$$

Example 3a

Find a unit vector for each given vector.

(a) $\mathbf{v} = 5\mathbf{i} - 12\mathbf{j}$

$$\begin{aligned}\|\mathbf{v}\| &= \sqrt{5^2 + 12^2} \\ &= 13\end{aligned}$$

Example 3a

Find a unit vector for each given vector.

(a) $\mathbf{v} = 5\mathbf{i} - 12\mathbf{j}$

$$\begin{aligned}\|\mathbf{v}\| &= \sqrt{5^2 + 12^2} \\ &= 13\end{aligned}$$

$$\hat{\mathbf{v}} = \frac{5}{13}\mathbf{i} - \frac{12}{13}\mathbf{j}$$

Example 3b

$$(b) \quad \vec{w} = \langle 4, -3 \rangle$$

Example 3b

$$(b) \quad \vec{w} = \langle 4, -3 \rangle$$

$$|\vec{w}| = \sqrt{4^2 + 3^2}$$

Example 3b

$$(b) \quad \vec{w} = \langle 4, -3 \rangle$$

$$\begin{aligned} |\vec{w}| &= \sqrt{4^2 + 3^2} \\ &= 5 \end{aligned}$$

Example 3b

$$(b) \quad \vec{w} = \langle 4, -3 \rangle$$

$$|\vec{w}| = \sqrt{4^2 + 3^2}$$

$$= 5$$

$$\hat{w} = \left\langle \frac{4}{5}, -\frac{3}{5} \right\rangle$$

Example 3c

(c) $\mathbf{d} = 7\mathbf{i} + \mathbf{j}$

Example 3c

$$(c) \quad \mathbf{d} = 7\mathbf{i} + \mathbf{j}$$

$$|\mathbf{d}| = \sqrt{7^2 + 1^2}$$

Example 3c

$$(c) \quad \mathbf{d} = 7\mathbf{i} + \mathbf{j}$$

$$|\mathbf{d}| = \sqrt{7^2 + 1^2}$$

$$= \sqrt{50}$$

Example 3c

$$(c) \quad \mathbf{d} = 7\mathbf{i} + \mathbf{j}$$

$$|\mathbf{d}| = \sqrt{7^2 + 1^2}$$

$$= \sqrt{50}$$

$$= 5\sqrt{2}$$

Example 3c

$$(c) \quad \mathbf{d} = 7\mathbf{i} + \mathbf{j}$$

$$|\mathbf{d}| = \sqrt{7^2 + 1^2}$$

$$= \sqrt{50}$$

$$= 5\sqrt{2}$$

$$\hat{\mathbf{d}} = \frac{7}{5\sqrt{2}}\mathbf{i} + \frac{1}{5\sqrt{2}}\mathbf{j}$$

Example 3c

$$(c) \quad \mathbf{d} = 7\mathbf{i} + \mathbf{j}$$

$$|\mathbf{d}| = \sqrt{7^2 + 1^2}$$

$$= \sqrt{50}$$

$$= 5\sqrt{2}$$


$$\hat{\mathbf{d}} = \frac{7}{5\sqrt{2}}\mathbf{i} + \frac{1}{5\sqrt{2}}\mathbf{j}$$

$$= \frac{7\sqrt{2}}{10}\mathbf{i} + \frac{\sqrt{2}}{2}\mathbf{j}$$

Table of Contents

Magnitude and Direction

If we place a vector $\langle x, y \rangle$ in the coordinate plane, we can put the initial point at the origin and the terminal point at (x, y) .



Magnitude and Direction

From Trig Functions of Any Angle,

$$x = \|\vec{v}\| \cos \theta \text{ and } y = \|\vec{v}\| \sin \theta$$

.

Magnitude and Direction

From Trig Functions of Any Angle,

$$x = \|\vec{v}\| \cos \theta \text{ and } y = \|\vec{v}\| \sin \theta$$

. Thus,

$$\langle x, y \rangle = \langle \|\vec{v}\| \cos \theta, \|\vec{v}\| \sin \theta \rangle = \|\vec{v}\| \langle \cos \theta, \sin \theta \rangle$$

Example 4

Find the horizontal and vertical component form of the vector whose magnitude and direction angle are given by $|u| = 12$ and $\theta = 150^\circ$.

Example 4

Find the horizontal and vertical component form of the vector whose magnitude and direction angle are given by $|u| = 12$ and $\theta = 150^\circ$.

$$x = 12 \cos 150^\circ \quad y = 12 \sin 150^\circ$$

Example 4

Find the horizontal and vertical component form of the vector whose magnitude and direction angle are given by $|u| = 12$ and $\theta = 150^\circ$.

$$x = 12 \cos 150^\circ \qquad y = 12 \sin 150^\circ$$

$$x = 12 \left(-\frac{\sqrt{3}}{2} \right) \qquad y = 12 \left(\frac{1}{2} \right)$$

Example 4

Find the horizontal and vertical component form of the vector whose magnitude and direction angle are given by $|u| = 12$ and $\theta = 150^\circ$.

$$x = 12 \cos 150^\circ \qquad y = 12 \sin 150^\circ$$

$$x = 12 \left(-\frac{\sqrt{3}}{2} \right) \qquad y = 12 \left(\frac{1}{2} \right)$$

$$x = -6\sqrt{3} \qquad y = 6$$

Example 4

Find the horizontal and vertical component form of the vector whose magnitude and direction angle are given by $|u| = 12$ and $\theta = 150^\circ$.

$$x = 12 \cos 150^\circ \qquad y = 12 \sin 150^\circ$$

$$x = 12 \left(-\frac{\sqrt{3}}{2} \right) \qquad y = 12 \left(\frac{1}{2} \right)$$

$$x = -6\sqrt{3} \qquad y = 6$$

$$\langle -6\sqrt{3}, 6 \rangle$$

Example 5

For $\vec{v} = \langle 3, -3\sqrt{3} \rangle$, find $\|\vec{v}\|$ and θ ($0 \leq \theta < 2\pi$) and write in $\|\vec{v}\|\langle \cos \theta, \sin \theta \rangle$ form.

Example 5

For $\vec{v} = \langle 3, -3\sqrt{3} \rangle$, find $\|\vec{v}\|$ and θ ($0 \leq \theta < 2\pi$) and write in $\|\vec{v}\|\langle \cos \theta, \sin \theta \rangle$ form.

$$\|\vec{v}\| = \sqrt{3^2 + (3\sqrt{3})^2}$$

Example 5

For $\vec{v} = \langle 3, -3\sqrt{3} \rangle$, find $\|\vec{v}\|$ and θ ($0 \leq \theta < 2\pi$) and write in $\|\vec{v}\|\langle \cos \theta, \sin \theta \rangle$ form.

$$\begin{aligned}\|\vec{v}\| &= \sqrt{3^2 + (3\sqrt{3})^2} \\ &= \sqrt{36} = 6\end{aligned}$$

Example 5

For $\vec{v} = \langle 3, -3\sqrt{3} \rangle$, find $\|\vec{v}\|$ and θ ($0 \leq \theta < 2\pi$) and write in $\|\vec{v}\|\langle \cos \theta, \sin \theta \rangle$ form.

$$\theta' = \tan^{-1} \left| \frac{-3\sqrt{3}}{3} \right|$$

Example 5

For $\vec{v} = \langle 3, -3\sqrt{3} \rangle$, find $\|\vec{v}\|$ and θ ($0 \leq \theta < 2\pi$) and write in $\|\vec{v}\|\langle \cos \theta, \sin \theta \rangle$ form.

$$\theta' = \tan^{-1} \left| \frac{-3\sqrt{3}}{3} \right|$$

$$\theta' = 60^\circ$$

Example 5

For $\vec{v} = \langle 3, -3\sqrt{3} \rangle$, find $\|\vec{v}\|$ and θ ($0 \leq \theta < 2\pi$) and write in $\|\vec{v}\|\langle \cos \theta, \sin \theta \rangle$ form.

$$\theta' = \tan^{-1} \left| \frac{-3\sqrt{3}}{3} \right|$$

$$\theta' = 60^\circ$$

$$\theta = 360^\circ - 60^\circ = 300^\circ$$

Example 5

For $\vec{v} = \langle 3, -3\sqrt{3} \rangle$, find $\|\vec{v}\|$ and θ ($0 \leq \theta < 2\pi$) and write in $\|\vec{v}\|\langle \cos \theta, \sin \theta \rangle$ form.

$$\theta' = \tan^{-1} \left| \frac{-3\sqrt{3}}{3} \right|$$

$$\theta' = 60^\circ$$

$$\theta = 360^\circ - 60^\circ = 300^\circ$$

$$= \frac{5\pi}{3}$$

Example 5

For $\vec{v} = \langle 3, -3\sqrt{3} \rangle$, find $\|\vec{v}\|$ and θ ($0 \leq \theta < 2\pi$) and write in $\|\vec{v}\|\langle \cos \theta, \sin \theta \rangle$ form.

$$\theta' = \tan^{-1} \left| \frac{-3\sqrt{3}}{3} \right|$$

$$\theta' = 60^\circ$$

$$\theta = 360^\circ - 60^\circ = 300^\circ$$

$$= \frac{5\pi}{3}$$

$$6 \left\langle \cos \left(\frac{5\pi}{3} \right), \sin \left(\frac{5\pi}{3} \right) \right\rangle$$