

Derivatives

Objectives

- 1 Find the derivative of a function

Difference Quotient

$$\frac{f(x+h) - f(x)}{h}$$

Derivative Definition

The **derivative** is the limit of the difference quotient as h approaches 0.

Derivative Definition

The **derivative** is the limit of the difference quotient as h approaches 0.

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

Derivative Definition

The **derivative** is the limit of the difference quotient as h approaches 0.

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

Other Notations: $\frac{dy}{dx}$ y' \dot{x}

Example 1

Find the derivative of each.

(a) $f(x) = 2x^2 - x + 3$

Example 1

Find the derivative of each.

(a) $f(x) = 2x^2 - x + 3$

$$f(x + h) = 2(x + h)^2 - (x + h) + 3$$

Example 1

Find the derivative of each.

(a) $f(x) = 2x^2 - x + 3$

$$f(x + h) = 2(x + h)^2 - (x + h) + 3$$

$$= 2(x^2 + 2hx + h^2) - x - h + 3$$

Example 1

Find the derivative of each.

(a) $f(x) = 2x^2 - x + 3$

$$f(x + h) = 2(x + h)^2 - (x + h) + 3$$

$$= 2(x^2 + 2hx + h^2) - x - h + 3$$

$$= 2x^2 + 4hx + 2h^2 - x - h + 3$$

Example 1

Find the derivative of each.

(a) $f(x) = 2x^2 - x + 3$

$$f(x+h) = 2(x+h)^2 - (x+h) + 3$$

$$= 2(x^2 + 2hx + h^2) - x - h + 3$$

$$= 2x^2 + 4hx + 2h^2 - x - h + 3$$

$$f(x+h) - f(x) = 2x^2 + 4hx + 2h^2 - x - h + 3 - (2x^2 - x + 3)$$

Example 1

Find the derivative of each.

(a) $f(x) = 2x^2 - x + 3$

$$f(x+h) = 2(x+h)^2 - (x+h) + 3$$

$$= 2(x^2 + 2hx + h^2) - x - h + 3$$

$$= 2x^2 + 4hx + 2h^2 - x - h + 3$$

$$f(x+h) - f(x) = 2x^2 + 4hx + 2h^2 - x - h + 3 - (2x^2 - x + 3)$$

$$= 2x^2 + 4hx + 2h^2 - x - h + 3 - 2x^2 + x - 3$$

Example 1

Find the derivative of each.

(a) $f(x) = 2x^2 - x + 3$

$$f(x+h) = 2(x+h)^2 - (x+h) + 3$$

$$= 2(x^2 + 2hx + h^2) - x - h + 3$$

$$= 2x^2 + 4hx + 2h^2 - x - h + 3$$

$$f(x+h) - f(x) = 2x^2 + 4hx + 2h^2 - x - h + 3 - (2x^2 - x + 3)$$

$$= 2x^2 + 4hx + 2h^2 - x - h + 3 - 2x^2 + x - 3$$

$$= 4hx - h + 2h^2$$

Example 1

$$\frac{f(x+h) - f(x)}{h} = \frac{4hx - h + 2h^2}{h}$$

Example 1

$$\begin{aligned}\frac{f(x+h) - f(x)}{h} &= \frac{4hx - h + 2h^2}{h} \\ &= 4x - 1 + 2h\end{aligned}$$

Example 1

$$\frac{f(x+h) - f(x)}{h} = \frac{4hx - h + 2h^2}{h}$$

$$= 4x - 1 + 2h$$

$$\lim_{h \rightarrow 0} (4x - 1 + 2h)$$

Example 1

$$\frac{f(x+h) - f(x)}{h} = \frac{4hx - h + 2h^2}{h}$$

$$= 4x - 1 + 2h$$

$$\lim_{h \rightarrow 0} (4x - 1 + 2h)$$

$$= 4x - 1 + 2(0)$$

Example 1

$$\frac{f(x+h) - f(x)}{h} = \frac{4hx - h + 2h^2}{h}$$

$$= 4x - 1 + 2h$$

$$\lim_{h \rightarrow 0} (4x - 1 + 2h)$$

$$= 4x - 1 + 2(0)$$

$$= 4x - 1$$

Example 1

$$\frac{f(x+h) - f(x)}{h} = \frac{4hx - h + 2h^2}{h}$$

$$= 4x - 1 + 2h$$

$$\lim_{h \rightarrow 0} (4x - 1 + 2h)$$

$$= 4x - 1 + 2(0)$$

$$= 4x - 1$$

$$f'(x) = 4x - 1$$

Example 1

$$(b) \quad f(x) = -3x^2$$

Example 1

$$(b) \quad f(x) = -3x^2$$

$$f(x + h) = -3(x + h)^2$$

Example 1

$$(b) \quad f(x) = -3x^2$$

$$\begin{aligned} f(x+h) &= -3(x+h)^2 \\ &= -3(x^2 + 2hx + h^2) \end{aligned}$$

Example 1

$$(b) \quad f(x) = -3x^2$$

$$f(x + h) = -3(x + h)^2$$

$$= -3(x^2 + 2hx + h^2)$$

$$= -3x^2 - 6hx - 3h^2$$

Example 1

$$(b) \quad f(x) = -3x^2$$

$$f(x + h) = -3(x + h)^2$$

$$= -3(x^2 + 2hx + h^2)$$

$$= -3x^2 - 6hx - 3h^2$$

$$f(x + h) - f(x) = -3x^2 - 6hx - 3h^2 - (-3x^2)$$

Example 1

$$(b) \quad f(x) = -3x^2$$

$$f(x+h) = -3(x+h)^2$$

$$= -3(x^2 + 2hx + h^2)$$

$$= -3x^2 - 6hx - 3h^2$$

$$f(x+h) - f(x) = -3x^2 - 6hx - 3h^2 - (-3x^2)$$

$$= -3x^2 - 6hx - 3h^2 + 3x^2$$

Example 1

$$(b) \quad f(x) = -3x^2$$

$$f(x+h) = -3(x+h)^2$$

$$= -3(x^2 + 2hx + h^2)$$

$$= -3x^2 - 6hx - 3h^2$$

$$f(x+h) - f(x) = -3x^2 - 6hx - 3h^2 - (-3x^2)$$

$$= -3x^2 - 6hx - 3h^2 + 3x^2$$

$$= -6hx - 3h^2$$

Example 1

$$\frac{f(x+h) - f(x)}{h} = \frac{-6hx - 3h^2}{h}$$

Example 1

$$\begin{aligned}\frac{f(x+h) - f(x)}{h} &= \frac{-6hx - 3h^2}{h} \\ &= -6x - 3h\end{aligned}$$

Example 1

$$\frac{f(x+h) - f(x)}{h} = \frac{-6hx - 3h^2}{h}$$

$$= -6x - 3h$$

$$\lim_{h \rightarrow 0} (-6x - 3h)$$

Example 1

$$\frac{f(x+h) - f(x)}{h} = \frac{-6hx - 3h^2}{h}$$

$$= -6x - 3h$$

$$\lim_{h \rightarrow 0} (-6x - 3h)$$

$$= -6x - 3(0)$$

Example 1

$$\frac{f(x+h) - f(x)}{h} = \frac{-6hx - 3h^2}{h}$$

$$= -6x - 3h$$

$$\lim_{h \rightarrow 0} (-6x - 3h)$$

$$= -6x - 3(0)$$

$$= -6x$$

Example 1

$$\frac{f(x+h) - f(x)}{h} = \frac{-6hx - 3h^2}{h}$$

$$= -6x - 3h$$

$$\lim_{h \rightarrow 0} (-6x - 3h)$$

$$= -6x - 3(0)$$

$$= -6x$$

$$f'(x) = -6x$$

Example 1

$$(c) \quad f(x) = 7x + 9$$

Example 1

$$(c) \quad f(x) = 7x + 9$$

$$f(x + h) = 7(x + h) + 9$$

Example 1

$$(c) \quad f(x) = 7x + 9$$

$$f(x + h) = 7(x + h) + 9$$

$$= 7x + 7h + 9$$

Example 1

$$(c) \quad f(x) = 7x + 9$$

$$f(x + h) = 7(x + h) + 9$$

$$= 7x + 7h + 9$$

$$f(x + h) - f(x) = 7x + 7h + 9 - (7x + 9)$$

Example 1

$$(c) \quad f(x) = 7x + 9$$

$$f(x + h) = 7(x + h) + 9$$

$$= 7x + 7h + 9$$

$$f(x + h) - f(x) = 7x + 7h + 9 - (7x + 9)$$

$$= 7x + 7h + 9 - 7x - 9$$

Example 1

$$(c) \quad f(x) = 7x + 9$$

$$f(x + h) = 7(x + h) + 9$$

$$= 7x + 7h + 9$$

$$f(x + h) - f(x) = 7x + 7h + 9 - (7x + 9)$$

$$= 7x + 7h + 9 - 7x - 9$$

$$= 7h$$

Example 1

$$\frac{f(x+h) - f(x)}{h} = \frac{7h}{h}$$

Example 1

$$\begin{aligned}\frac{f(x+h) - f(x)}{h} &= \frac{7h}{h} \\ &= 7\end{aligned}$$

Example 1

$$\frac{f(x+h) - f(x)}{h} = \frac{7h}{h}$$

$$= 7$$

$$\lim_{h \rightarrow 0} (7)$$

Example 1

$$\frac{f(x+h) - f(x)}{h} = \frac{7h}{h}$$

$$= 7$$

$$\lim_{h \rightarrow 0} (7)$$

$$= 7$$

Example 1

$$\frac{f(x+h) - f(x)}{h} = \frac{7h}{h}$$

$$= 7$$

$$\lim_{h \rightarrow 0} (7)$$

$$= 7$$

$$f'(x) = 7$$

Example 1

$$(d) \quad f(x) = x^2 + 3$$

Example 1

$$(d) \quad f(x) = x^2 + 3$$

$$f(x + h) = (x + h)^2 + 3$$

Example 1

$$(d) \quad f(x) = x^2 + 3$$

$$f(x + h) = (x + h)^2 + 3$$

$$= x^2 + 2hx + h^2 + 3$$

Example 1

$$(d) \quad f(x) = x^2 + 3$$

$$f(x + h) = (x + h)^2 + 3$$

$$= x^2 + 2hx + h^2 + 3$$

$$f(x + h) - f(x) = x^2 + 2hx + h^2 + 3 - (x^2 + 3)$$

Example 1

$$(d) \quad f(x) = x^2 + 3$$

$$f(x + h) = (x + h)^2 + 3$$

$$= x^2 + 2hx + h^2 + 3$$

$$f(x + h) - f(x) = x^2 + 2hx + h^2 + 3 - (x^2 + 3)$$

$$= x^2 + 2hx + h^2 + 3 - x^2 - 3$$

Example 1

$$(d) \quad f(x) = x^2 + 3$$

$$f(x + h) = (x + h)^2 + 3$$

$$= x^2 + 2hx + h^2 + 3$$

$$f(x + h) - f(x) = x^2 + 2hx + h^2 + 3 - (x^2 + 3)$$

$$= x^2 + 2hx + h^2 + 3 - x^2 - 3$$

$$= 2hx + h^2$$

Example 1

$$\frac{f(x+h) - f(x)}{h} = \frac{2hx + h^2}{h}$$

Example 1

$$\begin{aligned}\frac{f(x+h) - f(x)}{h} &= \frac{2hx + h^2}{h} \\ &= 2x + h\end{aligned}$$

Example 1

$$\frac{f(x+h) - f(x)}{h} = \frac{2hx + h^2}{h}$$

$$= 2x + h$$

$$\lim_{h \rightarrow 0} (2x + h)$$

Example 1

$$\frac{f(x+h) - f(x)}{h} = \frac{2hx + h^2}{h}$$

$$= 2x + h$$

$$\lim_{h \rightarrow 0} (2x + h)$$

$$= 2x + 0$$

Example 1

$$\frac{f(x+h) - f(x)}{h} = \frac{2hx + h^2}{h}$$

$$= 2x + h$$

$$\lim_{h \rightarrow 0} (2x + h)$$

$$= 2x + 0$$

$$= 2x$$

Example 1

$$\frac{f(x+h) - f(x)}{h} = \frac{2hx + h^2}{h}$$

$$= 2x + h$$

$$\lim_{h \rightarrow 0} (2x + h)$$

$$= 2x + 0$$

$$= 2x$$

$$f'(x) = 2x$$

Aliases

Derivatives have many aliases:

Aliases

Derivatives have many aliases:

- Instant rate of change.

Aliases

Derivatives have many aliases:

- Instant rate of change.
- Tangent slope to a curve.

Aliases

Derivatives have many aliases:

- Instant rate of change.
- Tangent slope to a curve.
- Limit as secant line becomes tangent line.

In calculus, you will learn shortcuts so you won't always have to do what we did in these notes to find the derivative.