

Measures of Center

Objectives

- 1 Calculate the mean, median, and mode of a dataset
- 2 Calculate the weighted mean of a dataset
- 3 Approximate the mean for a grouped dataset

Measures of Center

In this section, we will look at finding a central location for a dataset.

Measures of Center

In this section, we will look at finding a central location for a dataset.

In some instances, this can give us a good value to expect from that dataset.

The Mean

Mean

The **mean** of a dataset is found by adding all of the data values together and then dividing by the total number of values.

The Mean

Mean

The **mean** of a dataset is found by adding all of the data values together and then dividing by the total number of values.

When most people use the term *average*, they are referring to the mean.

Properties of the Mean

- Sample means drawn from the same population tend to vary less than other measures of center.

Properties of the Mean

- Sample means drawn from the same population tend to vary less than other measures of center.
- The mean of a data set uses every value, unless the mean is a *trimmed mean*.

Properties of the Mean

- Sample means drawn from the same population tend to vary less than other measures of center.
- The mean of a data set uses every value, unless the mean is a *trimmed mean*.
- One extreme value (called an **outlier**) can change the value of the mean drastically.

Mean Formula

Sample mean: \bar{x}

Population mean: μ

Mean Formula

Sample mean: \bar{x}

Population mean: μ

$$\bar{x}, \text{ or } \mu, = \frac{\sum x_i}{n} = \frac{1}{n} \sum x_i$$

The Median

Median

The **median** of a dataset is found by first arranging the data values from least to greatest, then selecting the data value in the middle.

The Median

Median

The **median** of a dataset is found by first arranging the data values from least to greatest, then selecting the data value in the middle.

If there are 2 data values in the middle, the median will be the mean of these two values.

Properties of the Median

The median separates the top 50% of the data from the bottom 50%.

Properties of the Median

The median separates the top 50% of the data from the bottom 50%.

Unlike, the mean, the median typically does not change by large amounts when we include just a few extreme values; in other words, the median is **resistant**.

Properties of the Median

The median separates the top 50% of the data from the bottom 50%.

Unlike, the mean, the median typically does not change by large amounts when we include just a few extreme values; in other words, the median is **resistant**.

The median is denoted by \tilde{x} , however, some technologies uses Med to denote it.

The Mode

Mode

The **mode** of a dataset is the value(s) that occur the most.

The Mode

Mode

The **mode** of a dataset is the value(s) that occur the most.

There may be one mode, no mode, or many modes (2 modes is called **bimodal**).

The Mode

Mode

The **mode** of a dataset is the value(s) that occur the most.

There may be one mode, no mode, or many modes (2 modes is called **bimodal**).

The mode is the only measure of center that is applicable to qualitative data.

Example 1

The dataset below represents the number of complaints I receive each week about my teaching.

8, 2, 3, 7, 4, 4, 1, 9, 7, 5

Calculate the mean, median, and mode of the number of complaints.

Example 1

The total number of complaints is 50. There are 10 complaints listed.

Example 1

The total number of complaints is 50. There are 10 complaints listed.

$$\text{Mean} = 50/10 = 5 \text{ complaints per week.}$$

Example 1

The total number of complaints is 50. There are 10 complaints listed.

$$\text{Mean} = 50/10 = 5 \text{ complaints per week.}$$

Sorted: 1, 2, 3, 4, 4, 5, 7, 7, 8, 9

Example 1

The total number of complaints is 50. There are 10 complaints listed.

$$\text{Mean} = 50/10 = 5 \text{ complaints per week.}$$

Sorted: 1, 2, 3, 4, 4, 5, 7, 7, 8, 9

Median number of complaints per week is 4.5

Example 1

The total number of complaints is 50. There are 10 complaints listed.

$$\text{Mean} = 50/10 = 5 \text{ complaints per week.}$$

Sorted: 1, 2, 3, 4, 4, 5, 7, 7, 8, 9

Median number of complaints per week is 4.5

Mode number of complaints per week are 4 and 7.

Example 2

The next week, I received 400 complaints. Re-calculate the mean, median, and mode now.

Example 2

The next week, I received 400 complaints. Re-calculate the mean, median, and mode now.

1, 2, 3, 4, 4, 5, 7, 7, 8, 9, 400

Example 2

The next week, I received 400 complaints. Re-calculate the mean, median, and mode now.

1, 2, 3, 4, 4, 5, 7, 7, 8, 9, 400

Mean: $450 / 11 = 40.9$ complaints per week.

Example 2

The next week, I received 400 complaints. Re-calculate the mean, median, and mode now.

1, 2, 3, 4, 4, 5, 7, 7, 8, 9, 400

Mean: $450 / 11 = 40.9$ complaints per week.

Median: 5 complaints per week.

Example 2

The next week, I received 400 complaints. Re-calculate the mean, median, and mode now.

1, 2, 3, 4, 4, 5, 7, 7, 8, 9, 400

Mean: $450 / 11 = 40.9$ complaints per week.

Median: 5 complaints per week.

Mode: 4 and 7 complaints per week.

Objectives

- 1 Calculate the mean, median, and mode of a dataset
- 2 Calculate the weighted mean of a dataset
- 3 Approximate the mean for a grouped dataset

Example 3

California has a mean class size of 20.9 students per teacher and Alaska has a mean of 16.8 students per teacher.

If we combine the two states, we might find the mean number of students per teacher to be 18.85 ($0.5 * (20.9 + 16.8)$) but is this result correct? Why or why not?

Example 3

California has a mean class size of 20.9 students per teacher and Alaska has a mean of 16.8 students per teacher.

If we combine the two states, we might find the mean number of students per teacher to be 18.85 ($0.5 * (20.9 + 16.8)$) but is this result correct? Why or why not?

It is not correct because California has a much higher population than Alaska. We would have to find what is known as the **weighted mean**.

Weighted Mean

Sometimes it is necessary to take into account how large each class of a data set is (such as the population of each US state).

Weighted Mean

Sometimes it is necessary to take into account how large each class of a data set is (such as the population of each US state).

If w_i represents the **weight** of each class, then the **weighted mean** can be found via

$$\frac{\sum (x_i \cdot w_i)}{\sum w_i}$$

Weighted Mean

Sometimes it is necessary to take into account how large each class of a data set is (such as the population of each US state).

If w_i represents the **weight** of each class, then the **weighted mean** can be found via

$$\frac{\sum (x_i \cdot w_i)}{\sum w_i}$$

In other words,

- 1 Multiply each data value by its corresponding weight.

Weighted Mean

Sometimes it is necessary to take into account how large each class of a data set is (such as the population of each US state).

If w_i represents the **weight** of each class, then the **weighted mean** can be found via

$$\frac{\sum (x_i \cdot w_i)}{\sum w_i}$$

In other words,

- 1 Multiply each data value by its corresponding weight.
- 2 Add those results.

Weighted Mean

Sometimes it is necessary to take into account how large each class of a data set is (such as the population of each US state).

If w_i represents the **weight** of each class, then the **weighted mean** can be found via

$$\frac{\sum (x_i \cdot w_i)}{\sum w_i}$$

In other words,

- 1 Multiply each data value by its corresponding weight.
- 2 Add those results.
- 3 Then divide that by the total of the weights.

Example 4

You've recently completed a semester. Determine the semester's GPA (A = 4pts, B = 3pts, etc).

Course	Grade	Credit Hours
Statistics	A	4
Advanced Chris Farley	A	3
<i>Airplane!</i> Quotes	B	5
Obnoxious Examples	C	3

Example 4

Grade	Credit Hours	TOTALS
4	4	16
4	3	12
3	5	15
2	3	6
	15	49

Example 4

Grade	Credit Hours	TOTALS
4	4	16
4	3	12
3	5	15
2	3	6
	15	49

Weighted Mean: $49/15 = 3.267$

Example 5

In a statistics course, tests count for 60% of the final grade, homework for 20% and midterm and final exams are 10% each. Suppose you've earned an 87% average on tests, 94% average on homeworks and a 77% average on the exams.

What is your overall percentage?

Example 5

In a statistics course, tests count for 60% of the final grade, homework for 20% and midterm and final exams are 10% each. Suppose you've earned an 87% average on tests, 94% average on homeworks and a 77% average on the exams.

What is your overall percentage?

Assessment	Your Scores	Grade Weights	TOTALS
Tests	0.87	0.60	0.522
Homework	0.94	0.20	0.188
Exams	0.77	0.20	0.154

Example 5

Grade Weights	TOTALS
0.60	0.522
0.20	0.188
0.20	0.154

Grade: $0.864/1 = 86.4\%$

Example 5

Grade Weights	TOTALS
0.60	0.522
0.20	0.188
0.20	0.154
1	0.864

Grade: $0.864/1 = 86.4\%$

Objectives

- 1 Calculate the mean, median, and mode of a dataset
- 2 Calculate the weighted mean of a dataset
- 3 Approximate the mean for a grouped dataset