Other Probability Distributions

Objectives

Solve problems involving geometric probability distributions

2 Solve problems involving Poisson probability distributions

With binomial distributions, we had the following conditions:

- There are a fixed number of *n* repeated independent trials
- Each trial's outcome is either a success or failure
- The probability of success, p, never changes

With binomial distributions, we had the following conditions:

- There are a fixed number of *n* repeated independent trials
- Each trial's outcome is either a success or failure
- The probability of success, p, never changes

We could calculate the probability of obtaining 8 heads out of 10 flips of a coin.

With binomial distributions, we had the following conditions:

- There are a fixed number of *n* repeated independent trials
- Each trial's outcome is either a success or failure
- The probability of success, p, never changes

We could calculate the probability of obtaining 8 heads out of 10 flips of a coin.

One of these outcomes is HHHTHHHTHH.

With geometric distributions, we would interested in the probability of obtaining our first success (such as flipping a tail) after x flips of the coin.

With geometric distributions, we would interested in the probability of obtaining our first success (such as flipping a tail) after x flips of the coin.

To put it another way, a geometric distribution has the following property:

With geometric distributions, we would interested in the probability of obtaining our first success (such as flipping a tail) after x flips of the coin.

To put it another way, a geometric distribution has the following property:

$$\underbrace{FFF\cdots F}_{x-1 \text{ failures}} S$$

Geometric Distributions

The probability of obtaining our first success after x binomial experiments is given by

$$P(X=x)=(1-p)^{x-1}\cdot p$$

A student is given a 10-question multiple choice test in which each question has 5 possible answers. What is the probability that the first question the student guesses correctly on the 4th question?

A student is given a 10-question multiple choice test in which each question has 5 possible answers. What is the probability that the first question the student guesses correctly on the 4th question?

FFFS

A student is given a 10-question multiple choice test in which each question has 5 possible answers. What is the probability that the first question the student guesses correctly on the 4th question?

FFFS

$$P(X = 4) = (1 - 0.2)^3(0.2)$$

A student is given a 10-question multiple choice test in which each question has 5 possible answers. What is the probability that the first question the student guesses correctly on the 4th question?

FFFS

$$P(X = 4) = (1 - 0.2)^{3}(0.2)$$
$$= 0.1024$$


A student is given a 10-question multiple choice test in which each question has 5 possible answers. What is the probability that the first question the student guesses correctly on the 4th question?

FFFS

$$P(X = 4) = (1 - 0.2)^{3}(0.2)$$
$$= 0.1024$$

The student has about an 10.24% chance of first guessing correctly on the 4th question.

Bar Graph of Example 1



Mean and Standard Deviation of Geometric Distributions

The mean (expected value) of a geometric distribution is the reciprocal of the probability of success:

$$E(X)=\frac{1}{p}$$

Mean and Standard Deviation of Geometric Distributions

The mean (expected value) of a geometric distribution is the reciprocal of the probability of success:

$$E(X)=\frac{1}{p}$$

The variance for a geometric distribution is the probability of failure divided by the square of the probability of success, given as

$$\sigma^2 = \frac{1-p}{p^2}$$

Mean and Standard Deviation of Geometric Distributions

The mean (expected value) of a geometric distribution is the reciprocal of the probability of success:

$$E(X) = \frac{1}{p}$$

The variance for a geometric distribution is the probability of failure divided by the square of the probability of success, given as

$$\sigma^2 = \frac{1-p}{p^2}$$

From which the standard deviation, σ is

$$\sigma = \sqrt{\frac{1-p}{p^2}} = \frac{\sqrt{1-p}}{p}$$

(a) A student is given a 10-question multiple choice test in which each question has 5 possible answers. What is the expected number of answers a student can guess before getting one right?

(a) A student is given a 10-question multiple choice test in which each question has 5 possible answers. What is the expected number of answers a student can guess before getting one right?

The expected value is the reciprocal of the probability of success:

(a) A student is given a 10-question multiple choice test in which each question has 5 possible answers. What is the expected number of answers a student can guess before getting one right?

The expected value is the reciprocal of the probability of success:

$$E(X) = \frac{1}{0.2}$$

(a) A student is given a 10-question multiple choice test in which each question has 5 possible answers. What is the expected number of answers a student can guess before getting one right?

The expected value is the reciprocal of the probability of success:

$$E(X) = \frac{1}{0.2}$$
$$= 5$$

(a) A student is given a 10-question multiple choice test in which each question has 5 possible answers. What is the expected number of answers a student can guess before getting one right?

The expected value is the reciprocal of the probability of success:

$$E(X) = \frac{1}{0.2}$$
$$= 5$$

We would expect the student to answer 5 questions before guessing one correctly.

(b) What is the standard deviation of the number of questions needed before the student guesses correctly?

(b) What is the standard deviation of the number of questions needed before the student guesses correctly?

The standard deviation is the square root of the probability of failue divided by the probability of success:

(b) What is the standard deviation of the number of questions needed before the student guesses correctly?

The standard deviation is the square root of the probability of failue divided by the probability of success:

$$\sigma = \frac{\sqrt{1 - 0.2}}{0.2}$$

(b) What is the standard deviation of the number of questions needed before the student guesses correctly?

The standard deviation is the square root of the probability of failue divided by the probability of success:

$$\sigma = \frac{\sqrt{1 - 0.2}}{0.2}$$

$$\approx 4.47$$

(b) What is the standard deviation of the number of questions needed before the student guesses correctly?

The standard deviation is the square root of the probability of failue divided by the probability of success:

$$\sigma = \frac{\sqrt{1 - 0.2}}{0.2}$$

$$\approx 4.47$$

The standard deviation is about 4.47 questions.

Objectives '

Solve problems involving geometric probability distributions

Solve problems involving Poisson probability distributions

Poisson Distribution

A **Poisson probability distribution** is a discrete probability distribution that involves occurrences of some event over a specified interval, such as time, distance, area, volume, etc.

Poisson Distribution

A **Poisson probability distribution** is a discrete probability distribution that involves occurrences of some event over a specified interval, such as time, distance, area, volume, etc.

Poisson probability distributions have the following characteristics:

Poisson Distribution

A **Poisson probability distribution** is a discrete probability distribution that involves occurrences of some event over a specified interval, such as time, distance, area, volume, etc.

Poisson probability distributions have the following characteristics:

 The experiment consists of counting the number of times, x, an event occurs in a given interval.

Poisson Distribution

A **Poisson probability distribution** is a discrete probability distribution that involves occurrences of some event over a specified interval, such as time, distance, area, volume, etc.

Poisson probability distributions have the following characteristics:

- The experiment consists of counting the number of times, x, an event occurs in a given interval.
- The probability of the event occurring is the same for each interval.

Poisson Distribution

A **Poisson probability distribution** is a discrete probability distribution that involves occurrences of some event over a specified interval, such as time, distance, area, volume, etc.

Poisson probability distributions have the following characteristics:

- The experiment consists of counting the number of times, *x*, an event occurs in a given interval.
- The probability of the event occurring is the same for each interval.
- The number of occurrences in one interval is independent of the number of occurrences in other intervals.

Poisson Formula

$$P(X=x) = \frac{\mu^x \cdot e^{-\mu}}{x!}$$

where $\boldsymbol{\mu}$ is the mean number of occurrences of the event over the

intervals and $e \approx 2.71828$

The mean number of dogs adopted from a rescue center per month is 4.

The mean number of dogs adopted from a rescue center per month is 4.

(a) What is the probability that in any given month, 5 dogs will be adopted?

The mean number of dogs adopted from a rescue center per month is 4.

(a) What is the probability that in any given month, 5 dogs will be adopted?

$$P(X=5) = \frac{4^5 \cdot e^{-4}}{5!}$$

The mean number of dogs adopted from a rescue center per month is 4.

(a) What is the probability that in any given month, 5 dogs will be adopted?

$$P(X = 5) = \frac{4^5 \cdot e^{-4}}{5!}$$

$$\approx 0.1563$$

The mean number of dogs adopted from a rescue center per month is 4.

(a) What is the probability that in any given month, 5 dogs will be adopted?

$$P(X = 5) = \frac{4^5 \cdot e^{-4}}{5!}$$

$$\approx 0.1563$$

There is about a 15.63% chance that 5 dogs will be adopted in a given month.

(b) What is the probability that in any given month, 2 or less dogs will be adopted?

(b) What is the probability that in any given month, 2 or less dogs will be adopted?

$$P(X \le 2) = P(X = 0) + P(X = 1) + P(X = 2)$$

(b) What is the probability that in any given month, 2 or less dogs will be adopted?

$$P(X \le 2) = P(X = 0) + P(X = 1) + P(X = 2)$$

 ≈ 0.2381

(b) What is the probability that in any given month, 2 or less dogs will be adopted?

$$P(X \le 2) = P(X = 0) + P(X = 1) + P(X = 2)$$

 ≈ 0.2381

There is about a 23.81% chance that 2 or less dogs will be adopted in a given month.

$$P(X \ge 3) = 1 - P(X \le 2)$$

$$P(X \ge 3) = 1 - P(X \le 2)$$

 $\approx 1 - 0.2381$

$$P(X \ge 3) = 1 - P(X \le 2)$$

$$\approx 1 - 0.2381$$

$$\approx 0.7619$$

(c) What is the probability that at least 3 dogs will be adopted in a given month?

$$P(X \ge 3) = 1 - P(X \le 2)$$

$$\approx 1 - 0.2381$$

$$\approx 0.7619$$

There is about a 76.19% chance that at least 3 dogs will be adopted in a given month.

Mean and Standard Deviation of Poisson Distributions

The **mean** of a Poisson distribution is given to us as μ , which also happens to be the variance as well.

Mean and Standard Deviation of Poisson Distributions

The **mean** of a Poisson distribution is given to us as μ , which also happens to be the variance as well.

Since the standard deviation is the square root of variance, it follows that the standard deviation of a Poisson distribution is also the square root of the mean:

$$\sigma = \sqrt{\mu}$$