

Hypothesis Testing

Regression

Slope of a Line

Recall that the slope of a line tells us how much the y -coordinates change by increasing each x -coordinate by 1.

Slope of a Line

Recall that the slope of a line tells us how much the y -coordinates change by increasing each x -coordinate by 1.

Previously, we looked at calculating regression equations in the form

$$\hat{y} = mx + b$$

which we can use to predict data values both inside and outside our dataset.

Slope of a Line

Recall that the slope of a line tells us how much the y -coordinates change by increasing each x -coordinate by 1.

Previously, we looked at calculating regression equations in the form

$$\hat{y} = mx + b$$

which we can use to predict data values both inside and outside our dataset.

Also, recall that the coefficient of determination, r^2 , tells us the prediction error decrease we get by using the linear regression equation, as opposed to the mean of the y -coordinates.

Using the y -Intercept

However, we might also be able to use the y -intercept of our linear regression equation as a predictor:

$$y = b$$

Using the y -Intercept

However, we might also be able to use the y -intercept of our linear regression equation as a predictor:

$$y = b$$

In this case, the slope of the equation, m , is 0.

Using the y -Intercept

However, we might also be able to use the y -intercept of our linear regression equation as a predictor:

$$y = b$$

In this case, the slope of the equation, m , is 0.

We can then test whether or not the linear regression model we obtained contributes to predicting values of y .

Hypotheses Regarding Slope

For this hypothesis, testing, we are operating under the assumption that the following null hypothesis is true:

The slope of the regression equation does not help in predicting values of y .

Hypotheses Regarding Slope

For this hypothesis, testing, we are operating under the assumption that the following null hypothesis is true:

The slope of the regression equation does not help in predicting values of y .

To put it mathematically:

$$H_0 : m = 0$$

Hypotheses Regarding Slope

For this hypothesis, testing, we are operating under the assumption that the following null hypothesis is true:

The slope of the regression equation does not help in predicting values of y .

To put it mathematically:

$$H_0 : m = 0$$

Our alternative hypothesis will be one of the forms

Left-Tailed	Right-Tailed	Two-Tailed
$m < 0$	$m > 0$	$m \neq 0$

Test Statistic

Since the population standard deviation of our regression slope is likely not to be known, we can estimate it by using the **estimated standard error of the least squares slope**.

Test Statistic

Since the population standard deviation of our regression slope is likely not to be known, we can estimate it by using the **estimated standard error of the least squares slope**.

$$\frac{s}{\sqrt{SS_{xx}}}$$

where

$$SS_{xx} = \sum x^2 - \frac{1}{n} \left(\sum x \right)^2$$

Test Statistic

Since the population standard deviation of our regression slope is likely not to be known, we can estimate it by using the **estimated standard error of the least squares slope**.

$$\frac{s}{\sqrt{SS_{xx}}}$$

where

$$SS_{xx} = \sum x^2 - \frac{1}{n} \left(\sum x \right)^2$$

Our test statistic, t , is thus

$$t = \frac{\hat{m}}{s/\sqrt{SS_{xx}}}$$

where \hat{m} is the slope of our regression equation.

p -Value

The area under the curve either to the left, right, or outside of our test statistic(s), i.e. the p -value, can be found by taking $n - 2$ degrees of freedom with that test statistic.

Confidence Intervals

We can construct confidence intervals as

$$\hat{m} \pm t_{\alpha/2} \left(s / \sqrt{SS_{xx}} \right)$$

Confidence Intervals

We can construct confidence intervals as

$$\hat{m} \pm t_{\alpha/2} \left(s / \sqrt{SS_{xx}} \right)$$

Remember, for one-sided confidence intervals, use t_{α} instead of $t_{\alpha/2}$.

Residuals Review

Recall that the **residual** of a data point is the distance the y -coordinate of that data point is from the corresponding y -coordinate on the regression equation.

Residuals Review

Recall that the **residual** of a data point is the distance the y -coordinate of that data point is from the corresponding y -coordinate on the regression equation.

We usually denote the residual of data point (x_i, y_i) as ϵ_i , and it can be found by

$$\epsilon_i = y_{\text{observed}} - y_{\text{predicted}}$$

Residuals Review

Recall that the **residual** of a data point is the distance the y -coordinate of that data point is from the corresponding y -coordinate on the regression equation.

We usually denote the residual of data point (x_i, y_i) as ϵ_i , and it can be found by

$$\epsilon_i = y_{\text{observed}} - y_{\text{predicted}}$$

When performing hypothesis testing regarding the slope of the regression equation, we operate under the following assumptions:

You Know What Happens When You Assume, Right?

Assumptions:

You Know What Happens When You Assume, Right?

Assumptions:

- The mean of all values of ϵ is 0.

You Know What Happens When You Assume, Right?

Assumptions:

- The mean of all values of ϵ is 0.
- The variance of all values of ϵ is constant for all values of x .

You Know What Happens When You Assume, Right?

Assumptions:

- The mean of all values of ϵ is 0.
- The variance of all values of ϵ is constant for all values of x .
- The distribution of ϵ is normal.

You Know What Happens When You Assume, Right?

Assumptions:

- The mean of all values of ϵ is 0.
- The variance of all values of ϵ is constant for all values of x .
- The distribution of ϵ is normal.
- Values of ϵ associated with any two values of y are independent.