# Binomial Probability Distributions

### **Objectives**

Calculate probabilities of binomial distributions

2 Calculate the mean, variance, and standard deviation of a binomial distribution

A **binomial probability experiment** is an experiment with the following conditions:

A **binomial probability experiment** is an experiment with the following conditions:

• There are a fixed number of *n* repeated independent trials

A **binomial probability experiment** is an experiment with the following conditions:

- There are a fixed number of *n* repeated independent trials
- Each trial's outcome is either a success or failure

A **binomial probability experiment** is an experiment with the following conditions:

- There are a fixed number of *n* repeated independent trials
- Each trial's outcome is either a success or failure
- The probability of success, p, never changes

Number of heads when flipping a coin 3 times:

x	Outcomes	P(X=x)
0	TTT	1/8
1	нтт тнт ттн	3/8
2	ннт нтн тнн	3/8
3	ннн	1/8

Number of heads when flipping a coin 3 times:

x	Outcomes	P(X=x)
0	TTT	1/8
1	нтт тнт ттн	3/8
2	ннт нтн тнн	3/8
3	ннн	1/8

In the outcome for 2 heads, HHT, HTH, and THH are all considered equal.

Number of heads when flipping a coin 3 times:

X	Outcomes	P(X=x)
0	TTT	1/8
1	нтт тнт ттн	3/8
2	ннт нтн тнн	3/8
3	ннн	1/8

In the outcome for 2 heads, HHT, HTH, and THH are all considered equal.

Thus, combinations play a role in binomial probability distributions.

Also note that if two heads are obtained when flipping a coin 3 times, then the remaining flips (1 in this case) must be tails.

Also note that if two heads are obtained when flipping a coin 3 times, then the remaining flips (1 in this case) must be tails.

Heads has a probability of  $\frac{1}{2}$  of occurring and tails has a probability of  $1-\frac{1}{2}$  of occurring.

Also note that if two heads are obtained when flipping a coin 3 times, then the remaining flips (1 in this case) must be tails.

Heads has a probability of  $\frac{1}{2}$  of occurring and tails has a probability of  $1-\frac{1}{2}$  of occurring.

Combining the idea of combinations, the number of successes of each event, and the probability of success and failure, we get the formula for calculating probabilities for binomial distributions:

Also note that if two heads are obtained when flipping a coin 3 times, then the remaining flips (1 in this case) must be tails.

Heads has a probability of  $\frac{1}{2}$  of occurring and tails has a probability of  $1-\frac{1}{2}$  of occurring.

Combining the idea of combinations, the number of successes of each event, and the probability of success and failure, we get the formula for calculating probabilities for binomial distributions:

$$P(X = x) = \binom{n}{x} \cdot p^{x} \cdot (1 - p)^{n - x}$$

Also note that if two heads are obtained when flipping a coin 3 times, then the remaining flips (1 in this case) must be tails.

Heads has a probability of  $\frac{1}{2}$  of occurring and tails has a probability of  $1-\frac{1}{2}$  of occurring.

Combining the idea of combinations, the number of successes of each event, and the probability of success and failure, we get the formula for calculating probabilities for binomial distributions:

$$P(X = x) = \binom{n}{x} \cdot p^{x} \cdot (1 - p)^{n - x}$$
$$= \frac{n!}{x!(n - x)!} \cdot p^{x} \cdot (1 - p)^{n - x}$$

Which is more likely when flipping a coin: obtaining 4 out of 5 tails or obtaining 8 out of 10 tails?

Which is more likely when flipping a coin: obtaining 4 out of 5 tails or obtaining 8 out of 10 tails?

4 out of 5:

Which is more likely when flipping a coin: obtaining 4 out of 5 tails or obtaining 8 out of 10 tails?

4 out of 5:

$$\frac{5!}{4!(5-4)!} \left(\frac{1}{2}\right)^4 \left(\frac{1}{2}\right)^1 = 0.15625$$

Which is more likely when flipping a coin: obtaining 4 out of 5 tails or obtaining 8 out of 10 tails?

4 out of 5:

$$\frac{5!}{4!(5-4)!} \left(\frac{1}{2}\right)^4 \left(\frac{1}{2}\right)^1 = 0.15625$$

8 out of 10:

Which is more likely when flipping a coin: obtaining 4 out of 5 tails or obtaining 8 out of 10 tails?

4 out of 5:

$$\frac{5!}{4!(5-4)!} \left(\frac{1}{2}\right)^4 \left(\frac{1}{2}\right)^1 = 0.15625$$

8 out of 10:

$$\frac{10!}{8!(10-8)!} \left(\frac{1}{2}\right)^8 \left(\frac{1}{2}\right)^2$$

Which is more likely when flipping a coin: obtaining 4 out of 5 tails or obtaining 8 out of 10 tails?

4 out of 5:

$$\frac{5!}{4!(5-4)!} \left(\frac{1}{2}\right)^4 \left(\frac{1}{2}\right)^1 = 0.15625$$

8 out of 10:

$$\frac{10!}{8!(10-8)!} \left(\frac{1}{2}\right)^8 \left(\frac{1}{2}\right)^2 \approx 0.04395$$

Which is more likely when flipping a coin: obtaining 4 out of 5 tails or obtaining 8 out of 10 tails?

4 out of 5:

$$\frac{5!}{4!(5-4)!} \left(\frac{1}{2}\right)^4 \left(\frac{1}{2}\right)^1 = 0.15625$$

8 out of 10:

$$\frac{10!}{8!(10-8)!} \left(\frac{1}{2}\right)^8 \left(\frac{1}{2}\right)^2 \approx 0.04395$$

It is more likely you will obtain 4 heads from 5 flips than 8 heads from 10 flips.

(a) A multiple choice test consists of 10 questions with 5 answer choices per question. What is the probability that a student guesses correctly on 6 questions?

(a) A multiple choice test consists of 10 questions with 5 answer choices per question. What is the probability that a student guesses correctly on 6 questions?

$$P(X=6) = \frac{10!}{6!(10-6)!} \left(\frac{1}{5}\right)^6 \left(\frac{4}{5}\right)^4$$

(a) A multiple choice test consists of 10 questions with 5 answer choices per question. What is the probability that a student guesses correctly on 6 questions?

$$P(X=6) = \frac{10!}{6!(10-6)!} \left(\frac{1}{5}\right)^6 \left(\frac{4}{5}\right)^4$$

$$\approx 0.00551$$

(b) What is the probability a student guesses at least 6 out of 10 questions correctly?

(b) What is the probability a student guesses at least 6 out of 10 questions correctly?

We can either add the probabilities from x = 6 to x = 10, or use a cumulative binomial calculation with the **complement rule** to achieve this as well.

(b) What is the probability a student guesses at least 6 out of 10 questions correctly?

We can either add the probabilities from x = 6 to x = 10, or use a cumulative binomial calculation with the **complement rule** to achieve this as well.

$$P(x \ge 6) = 1 - P(x < 6)$$

(b) What is the probability a student guesses at least 6 out of 10 questions correctly?

We can either add the probabilities from x = 6 to x = 10, or use a cumulative binomial calculation with the **complement rule** to achieve this as well.

$$P(x \ge 6) = 1 - P(x < 6)$$

$$\approx 0.00637$$

### **Objectives**

Calculate probabilities of binomial distributions

2 Calculate the mean, variance, and standard deviation of a binomial distribution