

Probability: OR

Objectives

- 1 Calculate probabilities using the Addition Rule
- 2 Calculate the complement of an event
- 3 Calculate "at least one" probabilities
- 4 Calculate the odds of an event

AND vs. OR

In the last section, we examined probabilities that focused on the word *and*.

AND vs. OR

In the last section, we examined probabilities that focused on the word *and*.

The word *and* meant that we **multiplied** the probabilities.

AND vs. OR

In the last section, we examined probabilities that focused on the word *and*.

The word *and* meant that we **multiplied** the probabilities.

In this section, we will focus on the word *or*, which will mean **adding** probabilities.

Example 1

A fair die is rolled. What is the probability of rolling a 4 or a 5.

Example 1

A fair die is rolled. What is the probability of rolling a 4 or a 5.

Number of outcomes in the sample space: 6

Example 1

A fair die is rolled. What is the probability of rolling a 4 or a 5.

Number of outcomes in the sample space: 6

What we want to happen: roll a 4 or a 5. This can happen in 2 ways.

Example 1

A fair die is rolled. What is the probability of rolling a 4 or a 5.

Number of outcomes in the sample space: 6

What we want to happen: roll a 4 or a 5. This can happen in 2 ways.

$$P(4 \text{ or } 5) = \frac{2}{6}$$

Example 1

A fair die is rolled. What is the probability of rolling a 4 or a 5.

Number of outcomes in the sample space: 6

What we want to happen: roll a 4 or a 5. This can happen in 2 ways.

$$\begin{aligned}P(4 \text{ or } 5) &= \frac{2}{6} \\&= \frac{1}{3}\end{aligned}$$

The Addition Rule

In the previous example, the events “rolling a 4” and “rolling a 5” were *mutually exclusive*.

The Addition Rule

In the previous example, the events “rolling a 4” and “rolling a 5” were *mutually exclusive*.

To find the OR probability of two mutually exclusive events, use the Addition Rule:


The Addition Rule

In the previous example, the events “rolling a 4” and “rolling a 5” were *mutually exclusive*.

To find the OR probability of two mutually exclusive events, use the Addition Rule:

$$P(A \text{ or } B) = P(A) + P(B)$$

Venn Diagram – OR



$$P(A \text{ or } B)$$

Example 2

The table below lists the types and numbers of cars sold at Lemon Autos along with their ages. Find each probability.

	0–2	3–5	6–10	Over 10	Total
Foreign	37	21	12	30	100
Domestic	35	23	11	31	100
Total	72	44	23	61	200

- (a) If a car is randomly selected, what is the probability that the car is 0–2 years old or over 10 years old?

Example 2

The table below lists the types and numbers of cars sold at Lemon Autos along with their ages. Find each probability.

	0–2	3–5	6–10	Over 10	Total
Foreign	37	21	12	30	100
Domestic	35	23	11	31	100
Total	72	44	23	61	200

(a) If a car is randomly selected, what is the probability that the car is 0–2 years old or over 10 years old?

$$P(0-2 \text{ or over } 10) = P(0 - 2) + P(\text{over } 10)$$

Example 2

The table below lists the types and numbers of cars sold at Lemon Autos along with their ages. Find each probability.

	0–2	3–5	6–10	Over 10	Total
Foreign	37	21	12	30	100
Domestic	35	23	11	31	100
Total	72	44	23	61	200

- (a) If a car is randomly selected, what is the probability that the car is 0–2 years old or over 10 years old?

$$\begin{aligned}P(0-2 \text{ or over } 10) &= P(0 - 2) + P(\text{over } 10) \\&= \frac{72}{200} + \frac{61}{200}\end{aligned}$$

Example 2

The table below lists the types and numbers of cars sold at Lemon Autos along with their ages. Find each probability.

	0-2	3-5	6-10	Over 10	Total
Foreign	37	21	12	30	100
Domestic	35	23	11	31	100
Total	72	44	23	61	200

(a) If a car is randomly selected, what is the probability that the car is 0-2 years old or over 10 years old?

$$\begin{aligned}P(0-2 \text{ or over } 10) &= P(0-2) + P(\text{over } 10) \\&= \frac{72}{200} + \frac{61}{200} \\&= \frac{133}{200}\end{aligned}$$

Example 2

	0-2	3-5	6-10	Over 10	Total
Foreign	37	21	12	30	100
Domestic	35	23	11	31	100
Total	72	44	23	61	200

(b) If a car is randomly selected, what is the probability that the car is 3-5 years old or a domestic car?

Example 2

	0-2	3-5	6-10	Over 10	Total
Foreign	37	21	12	30	100
Domestic	35	23	11	31	100
Total	72	44	23	61	200

(b) If a car is randomly selected, what is the probability that the car is 3-5 years old or a domestic car?

$$P(3-5 \text{ or domestic}) = P(3-5) + P(\text{domestic})$$

Example 2

	0-2	3-5	6-10	Over 10	Total
Foreign	37	21	12	30	100
Domestic	35	23	11	31	100
Total	72	44	23	61	200

(b) If a car is randomly selected, what is the probability that the car is 3-5 years old or a domestic car?

$$\begin{aligned}P(3-5 \text{ or domestic}) &= P(3-5) + P(\text{domestic}) \\&= \frac{44}{200} + \frac{100}{200}\end{aligned}$$

Example 2

	0-2	3-5	6-10	Over 10	Total
Foreign	37	21	12	30	100
Domestic	35	23	11	31	100
Total	72	44	23	61	200

(b) If a car is randomly selected, what is the probability that the car is 3-5 years old or a domestic car?

$$\begin{aligned}P(3-5 \text{ or domestic}) &= P(3-5) + P(\text{domestic}) \\&= \frac{44}{200} + \frac{100}{200} \\&= \frac{144}{200}\end{aligned}$$

Example 2

	0-2	3-5	6-10	Over 10	Total
Foreign	37	21	12	30	100
Domestic	35	23	11	31	100
Total	72	44	23	61	200

(b) If a car is randomly selected, what is the probability that the car is 3-5 years old or a domestic car?

$$\begin{aligned}P(3-5 \text{ or domestic}) &= P(3-5) + P(\text{domestic}) \\&= \frac{44}{200} + \frac{100}{200} \\&= \frac{144}{200} \\&= \frac{18}{25}\end{aligned}$$

Example 2

	0-2	3-5	6-10	Over 10	Total
Foreign	37	21	12	30	100
Domestic	35	23	11	31	100
Total	72	44	23	61	200

(b) If a car is randomly selected, what is the probability that the car is 3-5 years old or a domestic car?

$$\begin{aligned}P(3-5 \text{ or domestic}) &= P(3-5) + P(\text{domestic}) \\&= \frac{44}{200} + \frac{100}{200} \\&= \frac{144}{200} \\&= \frac{18}{25} \quad \dots \text{or is it?}\end{aligned}$$

Example 2 $P(3 - 5 \text{ or domestic})$

	0-2	3-5	6-10	Over 10	Total
Foreign	37	21	12	30	100
Domestic	35	23	11	31	100
Total	72	44	23	61	200

Example 2 $P(3 - 5 \text{ or domestic})$

	0-2	3-5	6-10	Over 10	Total
Foreign	37	21	12	30	100
Domestic	35	23	11	31	100
Total	72	44	23	61	200

There are 23 cars that are counted twice: once as a 3-5 year old car and again as a domestic car.

Example 2 $P(3 - 5 \text{ or domestic})$

	0-2	3-5	6-10	Over 10	Total
Foreign	37	21	12	30	100
Domestic	35	23	11	31	100
Total	72	44	23	61	200

There are 23 cars that are counted twice: once as a 3-5 year old car and again as a domestic car.

So, we need to subtract 23 cars from our original total of 144

Example 2 $P(3 - 5 \text{ or domestic})$

	0-2	3-5	6-10	Over 10	Total
Foreign	37	21	12	30	100
Domestic	35	23	11	31	100
Total	72	44	23	61	200

There are 23 cars that are counted twice: once as a 3-5 year old car and again as a domestic car.

So, we need to subtract 23 cars from our original total of 144


$$P(3 - 5 \text{ years old or domestic}) = \frac{121}{200}$$

General Addition Rule

$$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$$


General Addition Rule

$$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$$




General Addition Rule

$$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$$




General Addition Rule

$$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$$




General Addition Rule

$$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$$




Venn Diagram of Example 2b

	0-2	3-5	6-10	Over 10	Total
Foreign	37	21	12	30	100
Domestic	35	23	11	31	100
Total	72	44	23	61	200




Venn Diagram of Example 2b

	0-2	3-5	6-10	Over 10	Total
Foreign	37	21	12	30	100
Domestic	35	23	11	31	100
Total	72	44	23	61	200




Venn Diagram of Example 2b

	0-2	3-5	6-10	Over 10	Total
Foreign	37	21	12	30	100
Domestic	35	23	11	31	100
Total	72	44	23	61	200




Venn Diagram of Example 2b

	0-2	3-5	6-10	Over 10	Total
Foreign	37	21	12	30	100
Domestic	35	23	11	31	100
Total	72	44	23	61	200



Venn Diagram of Example 2b

	0-2	3-5	6-10	Over 10	Total
Foreign	37	21	12	30	100
Domestic	35	23	11	31	100
Total	72	44	23	61	200



$$23 + 21 + 77 = 121$$

Example 3

(a) If a single card is drawn from a standard deck, what is the probability of selecting a 3 or a club?

Example 3

(a) If a single card is drawn from a standard deck, what is the probability of selecting a 3 or a club?

Number of 3s: 4

Example 3

(a) If a single card is drawn from a standard deck, what is the probability of selecting a 3 or a club?

Number of 3s: 4

Number of clubs: 13

Example 3

(a) If a single card is drawn from a standard deck, what is the probability of selecting a 3 or a club?

Number of 3s: 4

Number of clubs: 13

Number of cards that are 3 and clubs: 1

Example 3

(a) If a single card is drawn from a standard deck, what is the probability of selecting a 3 or a club?

Number of 3s: 4

Number of clubs: 13

Number of cards that are 3 and clubs: 1

$$P(3 \text{ or club}) = P(3) + P(\text{club}) - P(3 \text{ and club})$$

Example 3

(a) If a single card is drawn from a standard deck, what is the probability of selecting a 3 or a club?

Number of 3s: 4

Number of clubs: 13

Number of cards that are 3 and clubs: 1

$$\begin{aligned} P(3 \text{ or club}) &= P(3) + P(\text{club}) - P(3 \text{ and club}) \\ &= \frac{4}{52} + \frac{13}{52} - \frac{1}{52} \end{aligned}$$

Example 3

(a) If a single card is drawn from a standard deck, what is the probability of selecting a 3 or a club?

Number of 3s: 4

Number of clubs: 13

Number of cards that are 3 and clubs: 1

$$P(3 \text{ or club}) = P(3) + P(\text{club}) - P(3 \text{ and club})$$

$$= \frac{4}{52} + \frac{13}{52} - \frac{1}{52}$$

$$= \frac{16}{52}$$

Example 3

(a) If a single card is drawn from a standard deck, what is the probability of selecting a 3 or a club?

Number of 3s: 4

Number of clubs: 13

Number of cards that are 3 and clubs: 1

$$P(3 \text{ or club}) = P(3) + P(\text{club}) - P(3 \text{ and club})$$

$$= \frac{4}{52} + \frac{13}{52} - \frac{1}{52}$$

$$= \frac{16}{52}$$

$$= \frac{4}{13}$$

Example 3

(b) What is the probability of drawing a face card or a red card?

Example 3

(b) What is the probability of drawing a face card or a red card?

Number of face cards: 12

Example 3

(b) What is the probability of drawing a face card or a red card?

Number of face cards: 12

Number of red cards: 26

Example 3

(b) What is the probability of drawing a face card or a red card?

Number of face cards: 12

Number of red cards: 26

Number of face cards that are also red: 6

Example 3

(b) What is the probability of drawing a face card or a red card?

Number of face cards: 12

Number of red cards: 26

Number of face cards that are also red: 6

$$P(\text{face or red}) = P(\text{face card}) + P(\text{red card}) - P(\text{face card and red})$$

Example 3

(b) What is the probability of drawing a face card or a red card?

Number of face cards: 12

Number of red cards: 26

Number of face cards that are also red: 6

$$P(\text{face or red}) = P(\text{face card}) + P(\text{red card}) - P(\text{face card and red})$$

$$= \frac{12}{52} + \frac{26}{52} - \frac{6}{52}$$

Example 3

(b) What is the probability of drawing a face card or a red card?

Number of face cards: 12

Number of red cards: 26

Number of face cards that are also red: 6

$$P(\text{face or red}) = P(\text{face card}) + P(\text{red card}) - P(\text{face card and red})$$

$$= \frac{12}{52} + \frac{26}{52} - \frac{6}{52}$$

$$= \frac{32}{52}$$

Example 3

(b) What is the probability of drawing a face card or a red card?

Number of face cards: 12

Number of red cards: 26

Number of face cards that are also red: 6

$$P(\text{face or red}) = P(\text{face card}) + P(\text{red card}) - P(\text{face card and red})$$

$$= \frac{12}{52} + \frac{26}{52} - \frac{6}{52}$$

$$= \frac{32}{52}$$

$$= \frac{8}{13}$$

Objectives

- 1 Calculate probabilities using the Addition Rule
- 2 Calculate the complement of an event
- 3 Calculate "at least one" probabilities
- 4 Calculate the odds of an event

Complements

Complements

The **complement** of an event is the probability the event does *not* happen.

Complements

Complements

The **complement** of an event is the probability the event does *not* happen.

Event: $P(A)$

Complement: $P(A')$

Complements

Complements

The **complement** of an event is the probability the event does *not* happen.

Event: $P(A)$

Complement: $P(A')$

$$P(A) + P(A') = 1$$

Complements

Complements

The **complement** of an event is the probability the event does *not* happen.

Event: $P(A)$

Complement: $P(A')$


$$P(A) + P(A') = 1$$

$$P(A') = 1 - P(A)$$


Objectives

- 1 Calculate probabilities using the Addition Rule
- 2 Calculate the complement of an event
- 3 Calculate "at least one" probabilities
- 4 Calculate the odds of an event

“At Least One” Probability




“At Least One” Probability




At least 1 is 1,

“At Least One” Probability




At least 1 is 1, or 2,

“At Least One” Probability




At least 1 is 1, or 2, or 3,

“At Least One” Probability



At least 1 is 1, or 2, or 3, ... or more.

“At Least One” Probability



At least 1 is 1, or 2, or 3, ... or more.

The complement of *at least one* is **none**.

Example 4

Two dice are rolled. What is the probability of rolling a sum of at least 4?

	1	2	3	4	5	6
1	2	3	4	5	6	7
2	3	4	5	6	7	8
3	4	5	6	7	8	9
4	5	6	7	8	9	10
5	6	7	8	9	10	11
6	7	8	9	10	11	12

Example 4

Two dice are rolled. What is the probability of rolling a sum of at least 4?

	1	2	3	4	5	6
1	2	3	4	5	6	7
2	3	4	5	6	7	8
3	4	5	6	7	8	9
4	5	6	7	8	9	10
5	6	7	8	9	10	11
6	7	8	9	10	11	12

$$P(\text{at least } 4) = 1 - P(\text{less than } 4)$$

Example 4

Two dice are rolled. What is the probability of rolling a sum of at least 4?

	1	2	3	4	5	6
1	2	3	4	5	6	7
2	3	4	5	6	7	8
3	4	5	6	7	8	9
4	5	6	7	8	9	10
5	6	7	8	9	10	11
6	7	8	9	10	11	12

$$\begin{aligned}P(\text{at least } 4) &= 1 - P(\text{less than } 4) \\&= 1 - P(2 \text{ or } 3)\end{aligned}$$

Example 4

Two dice are rolled. What is the probability of rolling a sum of at least 4?

	1	2	3	4	5	6
1	2	3	4	5	6	7
2	3	4	5	6	7	8
3	4	5	6	7	8	9
4	5	6	7	8	9	10
5	6	7	8	9	10	11
6	7	8	9	10	11	12

$$\begin{aligned}P(\text{at least } 4) &= 1 - P(\text{less than } 4) \\&= 1 - P(2 \text{ or } 3) \\&= 1 - \frac{3}{36}\end{aligned}$$

Example 4

Two dice are rolled. What is the probability of rolling a sum of at least 4?

	1	2	3	4	5	6
1	2	3	4	5	6	7
2	3	4	5	6	7	8
3	4	5	6	7	8	9
4	5	6	7	8	9	10
5	6	7	8	9	10	11
6	7	8	9	10	11	12

$$\begin{aligned}P(\text{at least } 4) &= 1 - P(\text{less than } 4) \\&= 1 - P(2 \text{ or } 3) \\&= 1 - \frac{3}{36} = \frac{33}{36}\end{aligned}$$

Example 4

Two dice are rolled. What is the probability of rolling a sum of at least 4?

	1	2	3	4	5	6
1	2	3	4	5	6	7
2	3	4	5	6	7	8
3	4	5	6	7	8	9
4	5	6	7	8	9	10
5	6	7	8	9	10	11
6	7	8	9	10	11	12

$$\begin{aligned}P(\text{at least } 4) &= 1 - P(\text{less than } 4) \\&= 1 - P(2 \text{ or } 3) \\&= 1 - \frac{3}{36} = \frac{33}{36} = \frac{11}{12}\end{aligned}$$

Example 5

A certain blood test can determine the presence of a bloodborne pathogen 97% of the time (that is, if 100 people have the pathogen, the test will confirm true for 97 of them). If 4 people with the pathogen are given the test, find the probability that the test is accurate for at least one of them.

Example 5

A certain blood test can determine the presence of a bloodborne pathogen 97% of the time (that is, if 100 people have the pathogen, the test will confirm true for 97 of them). If 4 people with the pathogen are given the test, find the probability that the test is accurate for at least one of them.

$$P(\text{at least 1 accurate}) = 1 - P(\text{none are accurate})$$

Example 5

A certain blood test can determine the presence of a bloodborne pathogen 97% of the time (that is, if 100 people have the pathogen, the test will confirm true for 97 of them). If 4 people with the pathogen are given the test, find the probability that the test is accurate for at least one of them.

$$\begin{aligned} P(\text{at least 1 accurate}) &= 1 - P(\text{none are accurate}) \\ &= 1 - P(\text{1st inaccurate}) \times P(\text{2nd inaccurate}) \cdots \end{aligned}$$

Example 5

A certain blood test can determine the presence of a bloodborne pathogen 97% of the time (that is, if 100 people have the pathogen, the test will confirm true for 97 of them). If 4 people with the pathogen are given the test, find the probability that the test is accurate for at least one of them.

$$\begin{aligned}P(\text{at least 1 accurate}) &= 1 - P(\text{none are accurate}) \\&= 1 - P(\text{1st inaccurate}) \times P(\text{2nd inaccurate}) \cdots \\&= 1 - (0.03)^4\end{aligned}$$

Example 5

A certain blood test can determine the presence of a bloodborne pathogen 97% of the time (that is, if 100 people have the pathogen, the test will confirm true for 97 of them). If 4 people with the pathogen are given the test, find the probability that the test is accurate for at least one of them.

$$\begin{aligned}P(\text{at least 1 accurate}) &= 1 - P(\text{none are accurate}) \\&= 1 - P(\text{1st inaccurate}) \times P(\text{2nd inaccurate}) \cdots \\&= 1 - (0.03)^4 \\&= 0.99999919\end{aligned}$$

Objectives

- 1 Calculate probabilities using the Addition Rule
- 2 Calculate the complement of an event
- 3 Calculate "at least one" probabilities
- 4 Calculate the odds of an event

Odds Definition

For events A and A' :

Odds Definition

For events A and A' :

Odds in Favor

The **odds in favor** of event A to happen are $\frac{A}{A'}$, or $A : A'$

Odds Definition

For events A and A' :

Odds in Favor

The **odds in favor** of event A to happen are $\frac{A}{A'}$, or $A : A'$

Odds Against

The **odds against** event A to happen are $\frac{A'}{A}$, or $A' : A$

Odds Definition

For events A and A' :

Odds in Favor

The **odds in favor** of event A to happen are $\frac{A}{A'}$, or $A : A'$

Odds Against

The **odds against** event A to happen are $\frac{A'}{A}$, or $A' : A$

Note: Typically when odds are listed, they are the odds against.

Example 6

The probability that the Cleveland Browns win the Super Bowl this year is 20%. What are the odds for and against this?

Example 6

The probability that the Cleveland Browns win the Super Bowl this year is 20%. What are the odds for and against this?

$$P(\text{win}) = 0.2$$

Example 6

The probability that the Cleveland Browns win the Super Bowl this year is 20%. What are the odds for and against this?

$$P(\text{win}) = 0.2$$

$$P(\text{don't win}) = 0.8$$

Example 6

The probability that the Cleveland Browns win the Super Bowl this year is 20%. What are the odds for and against this?

$$P(\text{win}) = 0.2$$

$$P(\text{don't win}) = 0.8$$

$$\text{odds for} = \frac{0.2}{0.8}$$

Example 6

The probability that the Cleveland Browns win the Super Bowl this year is 20%. What are the odds for and against this?

$$P(\text{win}) = 0.2$$

$$P(\text{don't win}) = 0.8$$

$$\text{odds for} = \frac{0.2}{0.8}$$

$$\text{odds for} = \frac{1}{4}$$

Example 6

The probability that the Cleveland Browns win the Super Bowl this year is 20%. What are the odds for and against this?

$$P(\text{win}) = 0.2$$

$$P(\text{don't win}) = 0.8$$

$$\text{odds for} = \frac{0.2}{0.8}$$

$$\text{odds for} = \frac{1}{4}$$

$$\text{odds against} = \frac{4}{1}$$

Example 7

A jar contains red and yellow marbles. The odds against selecting a red marble are 5 to 3. What is the probability of selecting a red marble?

Example 7

A jar contains red and yellow marbles. The odds against selecting a red marble are 5 to 3. What is the probability of selecting a red marble?

$$\frac{\text{yellow}}{\text{red}} = \frac{5}{3}$$

Example 7

A jar contains red and yellow marbles. The odds against selecting a red marble are 5 to 3. What is the probability of selecting a red marble?

$$\frac{\text{yellow}}{\text{red}} = \frac{5}{3} \longrightarrow \text{total marbles} = 8$$

$$P(\text{red marble}) = \frac{3}{8}$$