

Probability: OR

Objectives

- 1 Calculate probabilities using the Addition Rule
- 2 Calculate the complement of an event
- 3 Calculate "at least one" probabilities
- 4 Calculate the odds of an event

AND vs. OR

In the last section, we examined probabilities that focused on the word *and*.

AND vs. OR

In the last section, we examined probabilities that focused on the word *and*.

The word *and* meant that we **multiplied** the probabilities.

AND vs. OR

In the last section, we examined probabilities that focused on the word *and*.

The word *and* meant that we **multiplied** the probabilities.

In this section, we will focus on the word *or*, which will mean **adding** probabilities.

Example 1

A fair die is rolled. What is the probability of rolling a 4 or a 5.

Example 1

A fair die is rolled. What is the probability of rolling a 4 or a 5.

Number of outcomes in the sample space: 6

Example 1

A fair die is rolled. What is the probability of rolling a 4 or a 5.

Number of outcomes in the sample space: 6

What we want to happen: roll a 4 or a 5. This can happen in 2 ways.

Example 1

A fair die is rolled. What is the probability of rolling a 4 or a 5.

Number of outcomes in the sample space: 6

What we want to happen: roll a 4 or a 5. This can happen in 2 ways.

$$P(4 \text{ or } 5) = \frac{2}{6}$$

Example 1

A fair die is rolled. What is the probability of rolling a 4 or a 5.

Number of outcomes in the sample space: 6

What we want to happen: roll a 4 or a 5. This can happen in 2 ways.

$$\begin{aligned} P(4 \text{ or } 5) &= \frac{2}{6} \\ &= \frac{1}{3} \end{aligned}$$

The Addition Rule

In the previous example, the events “rolling a 4” and “rolling a 5” were *mutually exclusive*.

The Addition Rule

In the previous example, the events “rolling a 4” and “rolling a 5” were *mutually exclusive*.

To find the OR probability of two mutually exclusive events, use the Addition Rule:


The Addition Rule

In the previous example, the events “rolling a 4” and “rolling a 5” were *mutually exclusive*.

To find the OR probability of two mutually exclusive events, use the Addition Rule:

$$P(A \text{ or } B) = P(A) + P(B)$$

Venn Diagram – OR



$$P(A \text{ or } B)$$

Example 2

The table below lists the types and numbers of cars sold at Lemon Autos along with their ages. Find each probability.

| | 0–2 | 3–5 | 6–10 | Over 10 | Total |
|-----------------|------------|------------|-------------|----------------|--------------|
| Import | 37 | 21 | 12 | 30 | 100 |
| Domestic | 35 | 23 | 11 | 31 | 100 |
| Total | 72 | 44 | 23 | 61 | 200 |

- (a) If a car is randomly selected, what is the probability that the car is 0–2 years old or over 10 years old?

Example 2

The table below lists the types and numbers of cars sold at Lemon Autos along with their ages. Find each probability.

| | 0–2 | 3–5 | 6–10 | Over 10 | Total |
|----------|-----|-----|------|---------|-------|
| Import | 37 | 21 | 12 | 30 | 100 |
| Domestic | 35 | 23 | 11 | 31 | 100 |
| Total | 72 | 44 | 23 | 61 | 200 |

- (a) If a car is randomly selected, what is the probability that the car is 0–2 years old or over 10 years old?

$$P(0-2 \text{ or over } 10) = P(0 - 2) + P(\text{over } 10)$$

Example 2

The table below lists the types and numbers of cars sold at Lemon Autos along with their ages. Find each probability.

| | 0–2 | 3–5 | 6–10 | Over 10 | Total |
|----------|-----|-----|------|---------|-------|
| Import | 37 | 21 | 12 | 30 | 100 |
| Domestic | 35 | 23 | 11 | 31 | 100 |
| Total | 72 | 44 | 23 | 61 | 200 |

- (a) If a car is randomly selected, what is the probability that the car is 0–2 years old or over 10 years old?

$$\begin{aligned}P(0-2 \text{ or over } 10) &= P(0 - 2) + P(\text{over } 10) \\&= \frac{72}{200} + \frac{61}{200}\end{aligned}$$

Example 2

The table below lists the types and numbers of cars sold at Lemon Autos along with their ages. Find each probability.

| | 0–2 | 3–5 | 6–10 | Over 10 | Total |
|----------|-----|-----|------|---------|-------|
| Import | 37 | 21 | 12 | 30 | 100 |
| Domestic | 35 | 23 | 11 | 31 | 100 |
| Total | 72 | 44 | 23 | 61 | 200 |

- (a) If a car is randomly selected, what is the probability that the car is 0–2 years old or over 10 years old?

$$\begin{aligned}P(0-2 \text{ or over } 10) &= P(0 - 2) + P(\text{over } 10) \\&= \frac{72}{200} + \frac{61}{200} \\&= \frac{133}{200}\end{aligned}$$

Example 2

| | 0-2 | 3-5 | 6-10 | Over 10 | Total |
|----------|-----|-----|------|---------|-------|
| Import | 37 | 21 | 12 | 30 | 100 |
| Domestic | 35 | 23 | 11 | 31 | 100 |
| Total | 72 | 44 | 23 | 61 | 200 |

(b) If a car is randomly selected, what is the probability that the car is 3-5 years old or a domestic car?

Example 2

| | 0-2 | 3-5 | 6-10 | Over 10 | Total |
|----------|-----|-----|------|---------|-------|
| Import | 37 | 21 | 12 | 30 | 100 |
| Domestic | 35 | 23 | 11 | 31 | 100 |
| Total | 72 | 44 | 23 | 61 | 200 |

(b) If a car is randomly selected, what is the probability that the car is 3-5 years old or a domestic car?

$$P(3-5 \text{ or domestic}) = P(3-5) + P(\text{domestic})$$

Example 2

| | 0-2 | 3-5 | 6-10 | Over 10 | Total |
|----------|-----|-----|------|---------|-------|
| Import | 37 | 21 | 12 | 30 | 100 |
| Domestic | 35 | 23 | 11 | 31 | 100 |
| Total | 72 | 44 | 23 | 61 | 200 |

(b) If a car is randomly selected, what is the probability that the car is 3-5 years old or a domestic car?

$$\begin{aligned}P(3-5 \text{ or domestic}) &= P(3-5) + P(\text{domestic}) \\&= \frac{44}{200} + \frac{100}{200}\end{aligned}$$

Example 2

| | 0-2 | 3-5 | 6-10 | Over 10 | Total |
|----------|-----|-----|------|---------|-------|
| Import | 37 | 21 | 12 | 30 | 100 |
| Domestic | 35 | 23 | 11 | 31 | 100 |
| Total | 72 | 44 | 23 | 61 | 200 |

(b) If a car is randomly selected, what is the probability that the car is 3-5 years old or a domestic car?

$$\begin{aligned}P(3-5 \text{ or domestic}) &= P(3-5) + P(\text{domestic}) \\&= \frac{44}{200} + \frac{100}{200} \\&= \frac{144}{200}\end{aligned}$$

Example 2

| | 0-2 | 3-5 | 6-10 | Over 10 | Total |
|----------|-----|-----|------|---------|-------|
| Import | 37 | 21 | 12 | 30 | 100 |
| Domestic | 35 | 23 | 11 | 31 | 100 |
| Total | 72 | 44 | 23 | 61 | 200 |

(b) If a car is randomly selected, what is the probability that the car is 3-5 years old or a domestic car?

$$\begin{aligned}P(3-5 \text{ or domestic}) &= P(3-5) + P(\text{domestic}) \\&= \frac{44}{200} + \frac{100}{200} \\&= \frac{144}{200} \\&= \frac{18}{25}\end{aligned}$$

Example 2

| | 0-2 | 3-5 | 6-10 | Over 10 | Total |
|----------|-----|-----|------|---------|-------|
| Import | 37 | 21 | 12 | 30 | 100 |
| Domestic | 35 | 23 | 11 | 31 | 100 |
| Total | 72 | 44 | 23 | 61 | 200 |

(b) If a car is randomly selected, what is the probability that the car is 3-5 years old or a domestic car?

$$\begin{aligned}P(3-5 \text{ or domestic}) &= P(3-5) + P(\text{domestic}) \\&= \frac{44}{200} + \frac{100}{200} \\&= \frac{144}{200} \\&= \frac{18}{25} \quad \dots \text{or is it?}\end{aligned}$$

Example 2 $P(3 - 5 \text{ or domestic})$

| | 0-2 | 3-5 | 6-10 | Over 10 | Total |
|----------|-----|-----|------|---------|-------|
| Import | 37 | 21 | 12 | 30 | 100 |
| Domestic | 35 | 23 | 11 | 31 | 100 |
| Total | 72 | 44 | 23 | 61 | 200 |

Example 2 $P(3 - 5 \text{ or domestic})$

| | 0-2 | 3-5 | 6-10 | Over 10 | Total |
|----------|-----|-----|------|---------|-------|
| Import | 37 | 21 | 12 | 30 | 100 |
| Domestic | 35 | 23 | 11 | 31 | 100 |
| Total | 72 | 44 | 23 | 61 | 200 |

There are 23 cars that are counted twice: once as a 3-5 year old car and again as a domestic car.

Example 2 $P(3 - 5 \text{ or domestic})$

| | 0-2 | 3-5 | 6-10 | Over 10 | Total |
|----------|-----|-----|------|---------|-------|
| Import | 37 | 21 | 12 | 30 | 100 |
| Domestic | 35 | 23 | 11 | 31 | 100 |
| Total | 72 | 44 | 23 | 61 | 200 |

There are 23 cars that are counted twice: once as a 3-5 year old car and again as a domestic car.

So, we need to subtract 23 cars from our original total of 144

Example 2 $P(3 - 5 \text{ or domestic})$

| | 0-2 | 3-5 | 6-10 | Over 10 | Total |
|----------|-----|-----|------|---------|-------|
| Import | 37 | 21 | 12 | 30 | 100 |
| Domestic | 35 | 23 | 11 | 31 | 100 |
| Total | 72 | 44 | 23 | 61 | 200 |

There are 23 cars that are counted twice: once as a 3-5 year old car and again as a domestic car.

So, we need to subtract 23 cars from our original total of 144

$$P(3 - 5 \text{ years old or domestic}) = \frac{121}{200}$$

General Addition Rule

$$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$$


General Addition Rule

$$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$$




General Addition Rule

$$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$$




General Addition Rule

$$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$$




General Addition Rule

$$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$$




Venn Diagram of Example 2b

| | 0-2 | 3-5 | 6-10 | Over 10 | Total |
|----------|-----|-----|------|---------|-------|
| Import | 37 | 21 | 12 | 30 | 100 |
| Domestic | 35 | 23 | 11 | 31 | 100 |
| Total | 72 | 44 | 23 | 61 | 200 |




Venn Diagram of Example 2b

| | 0-2 | 3-5 | 6-10 | Over 10 | Total |
|----------|-----|-----|------|---------|-------|
| Import | 37 | 21 | 12 | 30 | 100 |
| Domestic | 35 | 23 | 11 | 31 | 100 |
| Total | 72 | 44 | 23 | 61 | 200 |




Venn Diagram of Example 2b

| | 0-2 | 3-5 | 6-10 | Over 10 | Total |
|----------|-----|-----|------|---------|-------|
| Import | 37 | 21 | 12 | 30 | 100 |
| Domestic | 35 | 23 | 11 | 31 | 100 |
| Total | 72 | 44 | 23 | 61 | 200 |




Venn Diagram of Example 2b

| | 0-2 | 3-5 | 6-10 | Over 10 | Total |
|----------|-----|-----|------|---------|-------|
| Import | 37 | 21 | 12 | 30 | 100 |
| Domestic | 35 | 23 | 11 | 31 | 100 |
| Total | 72 | 44 | 23 | 61 | 200 |



Venn Diagram of Example 2b

| | 0-2 | 3-5 | 6-10 | Over 10 | Total |
|----------|-----|-----|------|---------|-------|
| Import | 37 | 21 | 12 | 30 | 100 |
| Domestic | 35 | 23 | 11 | 31 | 100 |
| Total | 72 | 44 | 23 | 61 | 200 |



$$23 + 21 + 77 = 121$$

Example 3

(a) If a single card is drawn from a standard deck, what is the probability of selecting a 3 or a club?

Example 3

(a) If a single card is drawn from a standard deck, what is the probability of selecting a 3 or a club?

Number of 3s: 4

Example 3

(a) If a single card is drawn from a standard deck, what is the probability of selecting a 3 or a club?

Number of 3s: 4

Number of clubs: 13

Example 3

(a) If a single card is drawn from a standard deck, what is the probability of selecting a 3 or a club?

Number of 3s: 4

Number of clubs: 13

Number of cards that are 3 and clubs: 1

Example 3

(a) If a single card is drawn from a standard deck, what is the probability of selecting a 3 or a club?

Number of 3s: 4

Number of clubs: 13

Number of cards that are 3 and clubs: 1

$$P(3 \text{ or club}) = P(3) + P(\text{club}) - P(3 \text{ and club})$$

Example 3

(a) If a single card is drawn from a standard deck, what is the probability of selecting a 3 or a club?

Number of 3s: 4

Number of clubs: 13

Number of cards that are 3 and clubs: 1

$$\begin{aligned} P(3 \text{ or club}) &= P(3) + P(\text{club}) - P(3 \text{ and club}) \\ &= \frac{4}{52} + \frac{13}{52} - \frac{1}{52} \end{aligned}$$

Example 3

(a) If a single card is drawn from a standard deck, what is the probability of selecting a 3 or a club?

Number of 3s: 4

Number of clubs: 13

Number of cards that are 3 and clubs: 1

$$\begin{aligned}P(3 \text{ or club}) &= P(3) + P(\text{club}) - P(3 \text{ and club}) \\&= \frac{4}{52} + \frac{13}{52} - \frac{1}{52} \\&= \frac{16}{52}\end{aligned}$$

Example 3

(a) If a single card is drawn from a standard deck, what is the probability of selecting a 3 or a club?

Number of 3s: 4

Number of clubs: 13

Number of cards that are 3 and clubs: 1

$$P(3 \text{ or club}) = P(3) + P(\text{club}) - P(3 \text{ and club})$$

$$= \frac{4}{52} + \frac{13}{52} - \frac{1}{52}$$

$$= \frac{16}{52}$$

$$= \frac{4}{13}$$

Example 3

(b) What is the probability of drawing a face card or a red card?

Example 3

(b) What is the probability of drawing a face card or a red card?

Number of face cards: 12

Example 3

(b) What is the probability of drawing a face card or a red card?

Number of face cards: 12

Number of red cards: 26

Example 3

(b) What is the probability of drawing a face card or a red card?

Number of face cards: 12

Number of red cards: 26

Number of face cards that are also red: 6

Example 3

(b) What is the probability of drawing a face card or a red card?

Number of face cards: 12

Number of red cards: 26

Number of face cards that are also red: 6

$$P(\text{face or red}) = P(\text{face card}) + P(\text{red card}) - P(\text{face card and red})$$

Example 3

(b) What is the probability of drawing a face card or a red card?

Number of face cards: 12

Number of red cards: 26

Number of face cards that are also red: 6

$$P(\text{face or red}) = P(\text{face card}) + P(\text{red card}) - P(\text{face card and red})$$

$$= \frac{12}{52} + \frac{26}{52} - \frac{6}{52}$$

Example 3

(b) What is the probability of drawing a face card or a red card?

Number of face cards: 12

Number of red cards: 26

Number of face cards that are also red: 6

$$P(\text{face or red}) = P(\text{face card}) + P(\text{red card}) - P(\text{face card and red})$$

$$= \frac{12}{52} + \frac{26}{52} - \frac{6}{52}$$

$$= \frac{32}{52}$$

Example 3

(b) What is the probability of drawing a face card or a red card?

Number of face cards: 12

Number of red cards: 26

Number of face cards that are also red: 6

$$P(\text{face or red}) = P(\text{face card}) + P(\text{red card}) - P(\text{face card and red})$$

$$= \frac{12}{52} + \frac{26}{52} - \frac{6}{52}$$

$$= \frac{32}{52}$$

$$= \frac{8}{13}$$

Objectives

- 1 Calculate probabilities using the Addition Rule
- 2 Calculate the complement of an event
- 3 Calculate "at least one" probabilities
- 4 Calculate the odds of an event

Complements

Complements

The **complement** of an event is the probability the event does *not* happen.

Complements

Complements

The **complement** of an event is the probability the event does *not* happen.

Event: $P(A)$

Complement: $P(A')$

Complements

Complements

The **complement** of an event is the probability the event does *not* happen.

Event: $P(A)$

Complement: $P(A')$

$$P(A) + P(A') = 1$$

Complements

Complements

The **complement** of an event is the probability the event does *not* happen.

Event: $P(A)$

Complement: $P(A')$


$$P(A) + P(A') = 1$$

$$P(A') = 1 - P(A)$$


Objectives

- 1 Calculate probabilities using the Addition Rule
- 2 Calculate the complement of an event
- 3 Calculate "at least one" probabilities
- 4 Calculate the odds of an event

“At Least One” Probability




“At Least One” Probability



At least one means 1 or 2 or 3 or 4 or ...

“At Least One” Probability



At least one means 1 or 2 or 3 or 4 or ...

Rather than add all of the probabilities of those events up, we can instead find the **complement** of *at least one*.

“At Least One” Probability



At least one means 1 or 2 or 3 or 4 or ...

Rather than add all of the probabilities of those events up, we can instead find the **complement** of *at least one*.

The complement of *at least one* is **none**.

“At Least One” Probability



At least one means 1 or 2 or 3 or 4 or ...

Rather than add all of the probabilities of those events up, we can instead find the **complement** of *at least one*.

The complement of *at least one* is **none**.

In general, the complement of *at least n events* is **$n - 1$ events or less**.

Example 4

Two dice are rolled. What is the probability of rolling a sum of at least 4?

| | 1 | 2 | 3 | 4 | 5 | 6 |
|----------|----------|----------|----------|----------|----------|----------|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 |
| 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| 5 | 6 | 7 | 8 | 9 | 10 | 11 |
| 6 | 7 | 8 | 9 | 10 | 11 | 12 |

Example 4

Two dice are rolled. What is the probability of rolling a sum of at least 4?

| | 1 | 2 | 3 | 4 | 5 | 6 |
|---|---|---|---|----|----|----|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 |
| 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| 5 | 6 | 7 | 8 | 9 | 10 | 11 |
| 6 | 7 | 8 | 9 | 10 | 11 | 12 |

$$P(\text{at least } 4) = 1 - P(\text{less than } 4)$$

Example 4

Two dice are rolled. What is the probability of rolling a sum of at least 4?

| | 1 | 2 | 3 | 4 | 5 | 6 |
|---|---|---|---|----|----|----|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 |
| 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| 5 | 6 | 7 | 8 | 9 | 10 | 11 |
| 6 | 7 | 8 | 9 | 10 | 11 | 12 |

$$\begin{aligned}P(\text{at least } 4) &= 1 - P(\text{less than } 4) \\&= 1 - P(2 \text{ or } 3)\end{aligned}$$

Example 4

Two dice are rolled. What is the probability of rolling a sum of at least 4?

| | 1 | 2 | 3 | 4 | 5 | 6 |
|---|---|---|---|----|----|----|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 |
| 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| 5 | 6 | 7 | 8 | 9 | 10 | 11 |
| 6 | 7 | 8 | 9 | 10 | 11 | 12 |

$$\begin{aligned}P(\text{at least } 4) &= 1 - P(\text{less than } 4) \\&= 1 - P(2 \text{ or } 3) \\&= 1 - \frac{3}{36}\end{aligned}$$

Example 4

Two dice are rolled. What is the probability of rolling a sum of at least 4?

| | 1 | 2 | 3 | 4 | 5 | 6 |
|---|---|---|---|----|----|----|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 |
| 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| 5 | 6 | 7 | 8 | 9 | 10 | 11 |
| 6 | 7 | 8 | 9 | 10 | 11 | 12 |

$$\begin{aligned}P(\text{at least } 4) &= 1 - P(\text{less than } 4) \\&= 1 - P(2 \text{ or } 3) \\&= 1 - \frac{3}{36} = \frac{33}{36}\end{aligned}$$

Example 4

Two dice are rolled. What is the probability of rolling a sum of at least 4?

| | 1 | 2 | 3 | 4 | 5 | 6 |
|---|---|---|---|----|----|----|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 |
| 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| 5 | 6 | 7 | 8 | 9 | 10 | 11 |
| 6 | 7 | 8 | 9 | 10 | 11 | 12 |

$$\begin{aligned}P(\text{at least } 4) &= 1 - P(\text{less than } 4) \\&= 1 - P(2 \text{ or } 3) \\&= 1 - \frac{3}{36} = \frac{33}{36} = \frac{11}{12}\end{aligned}$$

Example 5

A certain blood test can determine the presence of a bloodborne pathogen 97% of the time (that is, if 100 people have the pathogen, the test will confirm true for 97 of them). If 4 people with the pathogen are given the test, find the probability that the test is accurate for at least one of them.

Example 5

A certain blood test can determine the presence of a bloodborne pathogen 97% of the time (that is, if 100 people have the pathogen, the test will confirm true for 97 of them). If 4 people with the pathogen are given the test, find the probability that the test is accurate for at least one of them.

$$P(\text{at least 1 accurate}) = 1 - P(\text{none are accurate})$$

Example 5

A certain blood test can determine the presence of a bloodborne pathogen 97% of the time (that is, if 100 people have the pathogen, the test will confirm true for 97 of them). If 4 people with the pathogen are given the test, find the probability that the test is accurate for at least one of them.

$$\begin{aligned} P(\text{at least 1 accurate}) &= 1 - P(\text{none are accurate}) \\ &= 1 - P(\text{1st inaccurate}) \times P(\text{2nd inaccurate}) \cdots \end{aligned}$$

Example 5

A certain blood test can determine the presence of a bloodborne pathogen 97% of the time (that is, if 100 people have the pathogen, the test will confirm true for 97 of them). If 4 people with the pathogen are given the test, find the probability that the test is accurate for at least one of them.

$$\begin{aligned}P(\text{at least 1 accurate}) &= 1 - P(\text{none are accurate}) \\&= 1 - P(\text{1st inaccurate}) \times P(\text{2nd inaccurate}) \cdots \\&= 1 - (0.03)^4\end{aligned}$$

Example 5

A certain blood test can determine the presence of a bloodborne pathogen 97% of the time (that is, if 100 people have the pathogen, the test will confirm true for 97 of them). If 4 people with the pathogen are given the test, find the probability that the test is accurate for at least one of them.

$$\begin{aligned}P(\text{at least 1 accurate}) &= 1 - P(\text{none are accurate}) \\&= 1 - P(\text{1st inaccurate}) \times P(\text{2nd inaccurate}) \cdots \\&= 1 - (0.03)^4 \\&= 0.99999919\end{aligned}$$

Objectives

- 1 Calculate probabilities using the Addition Rule
- 2 Calculate the complement of an event
- 3 Calculate "at least one" probabilities
- 4 Calculate the odds of an event

Odds Definition

For events A and A' :

Odds Definition

For events A and A' :

Odds in Favor

The **odds in favor** of event A to happen are $\frac{A}{A'}$, or $A : A'$

Odds Definition

For events A and A' :

Odds in Favor

The **odds in favor** of event A to happen are $\frac{A}{A'}$, or $A : A'$

Odds Against

The **odds against** event A to happen are $\frac{A'}{A}$, or $A' : A$

Odds Definition

For events A and A' :

Odds in Favor

The **odds in favor** of event A to happen are $\frac{A}{A'}$, or $A : A'$

Odds Against

The **odds against** event A to happen are $\frac{A'}{A}$, or $A' : A$

Note: Typically when odds are listed, they are the odds against.

Example 6

The probability that the Cleveland Browns win the Super Bowl this year is 20%. What are the odds for and against this?

Example 6

The probability that the Cleveland Browns win the Super Bowl this year is 20%. What are the odds for and against this?

$$P(\text{win}) = 0.2$$

Example 6

The probability that the Cleveland Browns win the Super Bowl this year is 20%. What are the odds for and against this?

$$P(\text{win}) = 0.2$$

$$P(\text{don't win}) = 0.8$$

Example 6

The probability that the Cleveland Browns win the Super Bowl this year is 20%. What are the odds for and against this?

$$P(\text{win}) = 0.2$$

$$P(\text{don't win}) = 0.8$$

$$\text{odds for} = \frac{0.2}{0.8}$$

Example 6

The probability that the Cleveland Browns win the Super Bowl this year is 20%. What are the odds for and against this?

$$P(\text{win}) = 0.2$$

$$P(\text{don't win}) = 0.8$$

$$\text{odds for} = \frac{0.2}{0.8}$$

$$\text{odds for} = \frac{1}{4}$$

Example 6

The probability that the Cleveland Browns win the Super Bowl this year is 20%. What are the odds for and against this?

$$P(\text{win}) = 0.2$$

$$P(\text{don't win}) = 0.8$$

$$\text{odds for} = \frac{0.2}{0.8}$$

$$\text{odds for} = \frac{1}{4}$$

$$\text{odds against} = \frac{4}{1}$$

Example 7

A jar contains red and yellow marbles. The odds against selecting a red marble are 5 to 3. What is the probability of selecting a red marble?

Example 7

A jar contains red and yellow marbles. The odds against selecting a red marble are 5 to 3. What is the probability of selecting a red marble?

$$\frac{\text{yellow}}{\text{red}} = \frac{5}{3}$$

Example 7

A jar contains red and yellow marbles. The odds against selecting a red marble are 5 to 3. What is the probability of selecting a red marble?

$$\frac{\text{yellow}}{\text{red}} = \frac{5}{3} \longrightarrow \text{total marbles} = 8$$

$$P(\text{red marble}) = \frac{3}{8}$$