

Speed Counting

Objectives

- 1 Use the Fundamental Counting Rule
- 2 Understand factorial notation
- 3 Find permutations of objects
- 4 Find combinations of objects
- 5 Find probabilities using counting techniques

Example 1

For a special at a restaurant, you can choose between 3 appetizers, 4 entrees, and 2 desserts. If you select one item from each category (appetizer, entree, and dessert), how many different meals can you create?

Example 1

For a special at a restaurant, you can choose between 3 appetizers, 4 entrees, and 2 desserts. If you select one item from each category (appetizer, entree, and dessert), how many different meals can you create?

Let's call the appetizers A, B, and C.

Example 1

For a special at a restaurant, you can choose between 3 appetizers, 4 entrees, and 2 desserts. If you select one item from each category (appetizer, entree, and dessert), how many different meals can you create?

Let's call the appetizers A, B, and C.

Let's call the entrees D, E, F, and G.

Example 1

For a special at a restaurant, you can choose between 3 appetizers, 4 entrees, and 2 desserts. If you select one item from each category (appetizer, entree, and dessert), how many different meals can you create?

Let's call the appetizers A, B, and C.

Let's call the entrees D, E, F, and G.

Let's call the desserts H and I.

Example 1

For a special at a restaurant, you can choose between 3 appetizers, 4 entrees, and 2 desserts. If you select one item from each category (appetizer, entree, and dessert), how many different meals can you create?

Let's call the appetizers A, B, and C.

Let's call the entrees D, E, F, and G.

Let's call the desserts H and I.

With appetizer A: ADH, ADI, AEH, AEI, AFH, AFI, AGH, AGI

Example 1

For a special at a restaurant, you can choose between 3 appetizers, 4 entrees, and 2 desserts. If you select one item from each category (appetizer, entree, and dessert), how many different meals can you create?

Let's call the appetizers A, B, and C.

Let's call the entrees D, E, F, and G.

Let's call the desserts H and I.

With appetizer A: ADH, ADI, AEH, AEI, AFH, AFI, AGH, AGI

With appetizer B: BDH, BDI, BEH, BEI, BFH, BFI, BGH, BGI

Example 1

For a special at a restaurant, you can choose between 3 appetizers, 4 entrees, and 2 desserts. If you select one item from each category (appetizer, entree, and dessert), how many different meals can you create?

Let's call the appetizers A, B, and C.

Let's call the entrees D, E, F, and G.

Let's call the desserts H and I.

With appetizer A: ADH, ADI, AEH, AEI, AFH, AFI, AGH, AGI

With appetizer B: BDH, BDI, BEH, BEI, BFH, BFI, BGH, BGI

With appetizer C: CDH, CDI, CEH, CEI, CFH, CFI, CGH, CGI

Example 1

For a special at a restaurant, you can choose between 3 appetizers, 4 entrees, and 2 desserts. If you select one item from each category (appetizer, entree, and dessert), how many different meals can you create?

Let's call the appetizers A, B, and C.

Let's call the entrees D, E, F, and G.

Let's call the desserts H and I.

With appetizer A: ADH, ADI, AEH, AEI, AFH, AFI, AGH, AGI

With appetizer B: BDH, BDI, BEH, BEI, BFH, BFI, BGH, BGI

With appetizer C: CDH, CDI, CEH, CEI, CFH, CFI, CGH, CGI

For a total of 24 possible different meals.

Fundamental Counting Rule

If event A can occur in a different ways and event B can occur in b different ways, then the total number of ways both events can occur is ab ways.

Fundamental Counting Rule

If event A can occur in a different ways and event B can occur in b different ways, then the total number of ways both events can occur is ab ways.

This can be generalized to multiple events, such as those in
example 1: $3 \times 4 \times 2 = 24$

Objectives

- 1 Use the Fundamental Counting Rule
- 2 Understand factorial notation
- 3 Find permutations of objects
- 4 Find combinations of objects
- 5 Find probabilities using counting techniques

Example 2

A baseball lineup consists of 9 players. How many different lineups using all 9 players on a team exist?

Example 2

A baseball lineup consists of 9 players. How many different lineups using all 9 players on a team exist?

Using the Fundamental Counting Rule with 9 positions:

Example 2

A baseball lineup consists of 9 players. How many different lineups using all 9 players on a team exist?

Using the Fundamental Counting Rule with 9 positions:

1st pos.	2nd pos.	3rd pos.	...	7th pos.	8th pos.	9th pos.

Example 2

A baseball lineup consists of 9 players. How many different lineups using all 9 players on a team exist?

Using the Fundamental Counting Rule with 9 positions:

1st pos.	2nd pos.	3rd pos.	...	7th pos.	8th pos.	9th pos.
9						

Example 2

A baseball lineup consists of 9 players. How many different lineups using all 9 players on a team exist?

Using the Fundamental Counting Rule with 9 positions:

1st pos.	2nd pos.	3rd pos.	...	7th pos.	8th pos.	9th pos.
9	8					

Example 2

A baseball lineup consists of 9 players. How many different lineups using all 9 players on a team exist?

Using the Fundamental Counting Rule with 9 positions:

1st pos.	2nd pos.	3rd pos.	...	7th pos.	8th pos.	9th pos.
9	8	7				

Example 2

A baseball lineup consists of 9 players. How many different lineups using all 9 players on a team exist?

Using the Fundamental Counting Rule with 9 positions:

1st pos.	2nd pos.	3rd pos.	...	7th pos.	8th pos.	9th pos.
9	8	7	...			

Example 2

A baseball lineup consists of 9 players. How many different lineups using all 9 players on a team exist?

Using the Fundamental Counting Rule with 9 positions:

1st pos.	2nd pos.	3rd pos.	...	7th pos.	8th pos.	9th pos.
9	8	7	...	3		

Example 2

A baseball lineup consists of 9 players. How many different lineups using all 9 players on a team exist?

Using the Fundamental Counting Rule with 9 positions:

1st pos.	2nd pos.	3rd pos.	...	7th pos.	8th pos.	9th pos.
9	8	7	...	3	2	

Example 2

A baseball lineup consists of 9 players. How many different lineups using all 9 players on a team exist?

Using the Fundamental Counting Rule with 9 positions:

1st pos.	2nd pos.	3rd pos.	...	7th pos.	8th pos.	9th pos.
9	8	7	...	3	2	1

Example 2

A baseball lineup consists of 9 players. How many different lineups using all 9 players on a team exist?

Using the Fundamental Counting Rule with 9 positions:

1st pos.	2nd pos.	3rd pos.	...	7th pos.	8th pos.	9th pos.
9	8	7	...	3	2	1

$$9 \times 8 \times 7 \times \cdots \times 2 \times 1$$

Example 2

A baseball lineup consists of 9 players. How many different lineups using all 9 players on a team exist?

Using the Fundamental Counting Rule with 9 positions:

1st pos.	2nd pos.	3rd pos.	...	7th pos.	8th pos.	9th pos.
9	8	7	...	3	2	1

$$9 \times 8 \times 7 \times \cdots \times 2 \times 1 = 362,880 \text{ unique lineups}$$

Factorial Notation

Rather than write out all the numbers from 9 to 1 and then multiplying them, mathematicians created **factorial notation** to expedite the process.

Factorial Notation

Rather than write out all the numbers from 9 to 1 and then multiplying them, mathematicians created **factorial notation** to expedite the process.

$$9! = 9 \times 8 \times 7 \times \cdots \times 3 \times 2 \times 1 = 362,880$$

Factorial Notation

Rather than write out all the numbers from 9 to 1 and then multiplying them, mathematicians created **factorial notation** to expedite the process.

$$9! = 9 \times 8 \times 7 \times \cdots \times 3 \times 2 \times 1 = 362,880$$

In general, for a positive integer n ,

$$n! = n(n-1)(n-2) \cdots 3(2)(1)$$

Factorial Notation

Rather than write out all the numbers from 9 to 1 and then multiplying them, mathematicians created **factorial notation** to expedite the process.

$$9! = 9 \times 8 \times 7 \times \cdots \times 3 \times 2 \times 1 = 362,880$$

In general, for a positive integer n ,

$$n! = n(n-1)(n-2) \cdots 3(2)(1)$$

with $0! = 1$

Factorial Growth

Factorial values grow very quickly:

$$2! = 2(1) = 2$$

$$3! = 3(2)(1) = 6$$

$$4! = 4(3)(2)(1) = 24$$

$$5! = 5(4)(3)(2)(1) = 120$$

$$6! = 6(5)(4)(3)(2)(1) = 720$$

$$7! = 7(6)(5)(4)(3)(2)(1) = 5,040$$

Example 3

How many ways are there to arrange 5 books on a shelf?

Example 3

How many ways are there to arrange 5 books on a shelf?

$5! = 120$ different arrangements

Objectives

- 1 Use the Fundamental Counting Rule
- 2 Understand factorial notation
- 3 Find permutations of objects
- 4 Find combinations of objects
- 5 Find probabilities using counting techniques

Example 4

Five people are competing for three prizes: \$1,000, \$500, and \$100. How many different ways can the prizes be awarded?

Example 4

Five people are competing for three prizes: \$1,000, \$500, and \$100. How many different ways can the prizes be awarded?

\$1,000	\$500	\$100

Example 4

Five people are competing for three prizes: \$1,000, \$500, and \$100. How many different ways can the prizes be awarded?

\$1,000	\$500	\$100
5		

Example 4

Five people are competing for three prizes: \$1,000, \$500, and \$100. How many different ways can the prizes be awarded?

\$1,000	\$500	\$100
5	4	

Example 4

Five people are competing for three prizes: \$1,000, \$500, and \$100. How many different ways can the prizes be awarded?

\$1,000	\$500	\$100
5	4	3

Example 4

Five people are competing for three prizes: \$1,000, \$500, and \$100. How many different ways can the prizes be awarded?

\$1,000	\$500	\$100
5	4	3

Using the Fundamental Counting Rule:

$$5 \times 4 \times 3 = 60 \text{ different ways}$$

Takeaways from Example 4

We had more contestants available to win prizes than we had prizes available. We could have had an equal number of contestants and prizes, but we can't have more prizes available than contestants.

Takeaways from Example 4

We had more contestants available to win prizes than we had prizes available. We could have had an equal number of contestants and prizes, but we can't have more prizes available than contestants.

If we had 10,000 contestants and 75 prizes, we would have a lot of multiplying to do.

Takeaways from Example 4

We had more contestants available to win prizes than we had prizes available. We could have had an equal number of contestants and prizes, but we can't have more prizes available than contestants.

If we had 10,000 contestants and 75 prizes, we would have a lot of multiplying to do.

So is there an easy way to do this if that's the case?

Takeaways from Example 4

We had more contestants available to win prizes than we had prizes available. We could have had an equal number of contestants and prizes, but we can't have more prizes available than contestants.

If we had 10,000 contestants and 75 prizes, we would have a lot of multiplying to do.

So is there an easy way to do this if that's the case?

Yes, and that is where **permutations** come into play.

Example 5

How many ways are there to award gold, silver, and bronze medals to 8 contestants?

Example 5

How many ways are there to award gold, silver, and bronze medals to 8 contestants?

Gold	Silver	Bronze

Example 5

How many ways are there to award gold, silver, and bronze medals to 8 contestants?

Gold	Silver	Bronze
8		

Example 5

How many ways are there to award gold, silver, and bronze medals to 8 contestants?

Gold	Silver	Bronze
8	7	

Example 5

How many ways are there to award gold, silver, and bronze medals to 8 contestants?

Gold	Silver	Bronze
8	7	6

Example 5

How many ways are there to award gold, silver, and bronze medals to 8 contestants?

Gold	Silver	Bronze
8	7	6

Using the Fundamental Counting Rule:

$$8 \times 7 \times 6 = 336 \text{ different ways}$$

Example 5

$$8 \times 7 \times 6 \times \cdots \times 2 \times 1$$

Example 5

$$\frac{8 \times 7 \times 6 \times 5 \times 4 \times 3 \times 2 \times 1}{5 \times 4 \times 3 \times 2 \times 1}$$

Example 5

$$8 \times 7 \times 6 = 336$$

Permutations

If there are n items available and we take r at a time, then the total number of permutations is given by

$${}_nP_r = \frac{n!}{(n-r)!}$$

with $n \geq r$

Permutations

If there are n items available and we take r at a time, then the total number of permutations is given by

$${}_nP_r = \frac{n!}{(n-r)!}$$

with $n \geq r$

With permutations, the order in which an item is selected matters.

Knowing to Use Permutations

Permutations

- Offering various prizes

Knowing to Use Permutations

Permutations

- Offering various prizes
- Running a race

Knowing to Use Permutations

Permutations

- Offering various prizes
- Running a race
- Assigning officer positions

Knowing to Use Permutations

Permutations

- Offering various prizes
- Running a race
- Assigning officer positions
- Combination locks and passwords

Example 6

How many ways are there of selecting a president, vice president, secretary, and treasurer out of a pool of 10 candidates?

Example 6

How many ways are there of selecting a president, vice president, secretary, and treasurer out of a pool of 10 candidates?

Order matters.

Example 6

How many ways are there of selecting a president, vice president, secretary, and treasurer out of a pool of 10 candidates?

Order matters.

We have $n = 10$ items to choose from.

Example 6

How many ways are there of selecting a president, vice president, secretary, and treasurer out of a pool of 10 candidates?

Order matters.

We have $n = 10$ items to choose from.

We are filling $r = 4$ positions at a time.

Example 6

How many ways are there of selecting a president, vice president, secretary, and treasurer out of a pool of 10 candidates?

Order matters.

We have $n = 10$ items to choose from.

We are filling $r = 4$ positions at a time.

$${}_{10}P_4 = \frac{10!}{(10 - 4)!}$$

Example 6

How many ways are there of selecting a president, vice president, secretary, and treasurer out of a pool of 10 candidates?

Order matters.

We have $n = 10$ items to choose from.

We are filling $r = 4$ positions at a time.

$$\begin{aligned} {}_{10}P_4 &= \frac{10!}{(10 - 4)!} \\ &= \frac{10!}{6!} \end{aligned}$$

Example 6

How many ways are there of selecting a president, vice president, secretary, and treasurer out of a pool of 10 candidates?

Order matters.

We have $n = 10$ items to choose from.

We are filling $r = 4$ positions at a time.

$$\begin{aligned} {}_{10}P_4 &= \frac{10!}{(10 - 4)!} \\ &= \frac{10!}{6!} \\ &= 5,040 \end{aligned}$$

Objectives

- 1 Use the Fundamental Counting Rule
- 2 Understand factorial notation
- 3 Find permutations of objects
- 4 Find combinations of objects**
- 5 Find probabilities using counting techniques

Combinations

With permutations, order selection mattered, so

ABC, ACB, BAC, BCA, CAB, and CBA

were all different.

Combinations

With permutations, order selection mattered, so

ABC, ACB, BAC, BCA, CAB, and CBA

were all different.

With combinations, selection order does not matter, so there is no distinction among the orderings above. So,

ABC, ACB, BAC, BCA, CAB, and CBA

are all the same.

Combinations

Notice that there are 6, or $3!$, arrangements of the letters A, B, and C.

Combinations

Notice that there are 6, or $3!$, arrangements of the letters A, B, and C.

This can help us develop the formula for finding the number of combinations of n items taken r at a time.

Example 7

Five people are competing for three equal prizes. How many ways can the prizes be awarded?

If order mattered, there would be ${}_5P_3 = 60$ different possibilities:

ABC	ABD	ABE	ACB	ACD	ACE	...
BAC	BAD	BAE	BCA	BCD	BCE	...
⋮	⋮	⋮	⋮	⋮	⋮	...

Example 7

Since ABC is the same as ACB, BAC, BCA, CAB, and CBA in the eyes of combinations, we can divide our permutation result of 60 by $3!$ to get **10 combinations**:

ABC	ABD	ABE	ACD	ACE
ADE	BCD	BCE	BDE	CDE

Combinations

If we have n items available and we take r at a time **without regard to order of selection**, then the total number of possible combinations are

Combinations

If we have n items available and we take r at a time **without regard to order of selection**, then the total number of possible combinations are

$${}_nC_r = \frac{n!}{r!(n-r)!}$$

Combinations

If we have n items available and we take r at a time **without regard to order of selection**, then the total number of possible combinations are

$${}_nC_r = \frac{n!}{r!(n-r)!} = \binom{n}{r}$$

Combinations

If we have n items available and we take r at a time **without regard to order of selection**, then the total number of possible combinations are

$${}_nC_r = \frac{n!}{r!(n-r)!} = \binom{n}{r}$$

Combinations

Combinations

If we have n items available and we take r at a time **without regard to order of selection**, then the total number of possible combinations are

$${}_nC_r = \frac{n!}{r!(n-r)!} = \binom{n}{r}$$

Combinations

- Awarding equal prizes

Combinations

If we have n items available and we take r at a time **without regard to order of selection**, then the total number of possible combinations are

$${}_nC_r = \frac{n!}{r!(n-r)!} = \binom{n}{r}$$

Combinations

- Awarding equal prizes
- Combinations (not the lock though)

Combinations

If we have n items available and we take r at a time **without regard to order of selection**, then the total number of possible combinations are

$${}_nC_r = \frac{n!}{r!(n-r)!} = \binom{n}{r}$$

Combinations

- Awarding equal prizes
- Combinations (not the lock though)
- Committees

Example 8

A committee of 5 is to be formed from a pool of 12 potential candidates. How many committees are possible?

Example 8

A committee of 5 is to be formed from a pool of 12 potential candidates. How many committees are possible?

We have $n = 12$ candidates to choose from

Example 8

A committee of 5 is to be formed from a pool of 12 potential candidates. How many committees are possible?

We have $n = 12$ candidates to choose from

We are selecting $r = 5$ at a time

Example 8

A committee of 5 is to be formed from a pool of 12 potential candidates. How many committees are possible?

We have $n = 12$ candidates to choose from

We are selecting $r = 5$ at a time

$${}_{12}C_5 = \frac{12!}{5!(12 - 5)!}$$

Example 8

A committee of 5 is to be formed from a pool of 12 potential candidates. How many committees are possible?

We have $n = 12$ candidates to choose from

We are selecting $r = 5$ at a time

$$\begin{aligned} {}_{12}C_5 &= \frac{12!}{5!(12-5)!} \\ &= 792 \end{aligned}$$

Example 9

A committee of 5 is to be formed from a pool of 12 potential candidates. The committee is to be made up of 3 managers and 2 accountants. If there are 8 managers and 4 accountants available, how many committees can be formed?

Example 9

A committee of 5 is to be formed from a pool of 12 potential candidates. The committee is to be made up of 3 managers and 2 accountants. If there are 8 managers and 4 accountants available, how many committees can be formed?

We need to make sure 3 of the positions are managers and 2 of the positions are accountants.

Example 9

A committee of 5 is to be formed from a pool of 12 potential candidates. The committee is to be made up of 3 managers and 2 accountants. If there are 8 managers and 4 accountants available, how many committees can be formed?

We need to make sure 3 of the positions are managers and 2 of the positions are accountants.

Thus, we must blend the Fundamental Counting Rule with combinations.

Example 9

Select 3 managers from 8 \times Select 2 accountants from 4

\times

\times

Example 9

Select 3 managers from 8 \times Select 2 accountants from 4

${}_8C_3$

\times

\times

Example 9

Select 3 managers from 8 \times Select 2 accountants from 4

${}_8C_3$

\times

${}_4C_2$

\times

Example 9

Select 3 managers from 8 \times Select 2 accountants from 4

$${}_8C_3$$

 \times

$${}_4C_2$$

$$56$$

 \times

Example 9

Select 3 managers from 8 \times Select 2 accountants from 4

$${}_8C_3$$

 \times

$${}_4C_2$$

$$56$$

 \times

$$6$$

Example 9

Select 3 managers from 8	×	Select 2 accountants from 4
${}_8C_3$	×	${}_4C_2$
56	×	6

There are 336 ways to do this.

Example 10

Starting at point A and only moving right or up, how many paths are there to get to point B ?

Objectives

- 1 Use the Fundamental Counting Rule
- 2 Understand factorial notation
- 3 Find permutations of objects
- 4 Find combinations of objects
- 5 Find probabilities using counting techniques