

## Other Probability Distributions

# Objectives

- 1 Solve problems involving geometric probability distributions
- 2 Solve problems involving hypergeometric probability distributions

# Binomial vs. Geometric Distributions

With binomial distributions, we had the following conditions:

- There are a fixed number of  $n$  repeated independent trials
- Each trial's outcome is either a success or failure
- The probability of success,  $p$ , never changes

# Binomial vs. Geometric Distributions

With binomial distributions, we had the following conditions:

- There are a fixed number of  $n$  repeated independent trials
- Each trial's outcome is either a success or failure
- The probability of success,  $p$ , never changes

We could calculate the probability of obtaining 8 heads out of 10 flips of a coin.

# Binomial vs. Geometric Distributions

With binomial distributions, we had the following conditions:

- There are a fixed number of  $n$  repeated independent trials
- Each trial's outcome is either a success or failure
- The probability of success,  $p$ , never changes

We could calculate the probability of obtaining 8 heads out of 10 flips of a coin.

One of these outcomes is TTHTHHHTTH.

# Binomial vs. Geometric Distributions

With geometric distributions, we would be interested in the probability of obtaining our first success (flipping a tail) after  $x$  flips of the coin.

# Binomial vs. Geometric Distributions

With geometric distributions, we would be interested in the probability of obtaining our first success (flipping a tail) after  $x$  flips of the coin.

To put it another way, a geometric distribution has the following property:

# Binomial vs. Geometric Distributions

With geometric distributions, we would be interested in the probability of obtaining our first success (flipping a tail) after  $x$  flips of the coin.

To put it another way, a geometric distribution has the following property:

$$\underbrace{FFF \dots FS}_{x-1 \text{ failures}}$$



# Geometric Distributions

The probability of obtaining our first success after  $x$  binomial experiments is given by

$$P(X = x) = (1 - p)^{x-1} \cdot p$$


## Example 1

A student is given a 10-question multiple choice test in which each question has 5 possible answers. What is the probability that the first question the student guesses correctly on the 4th question?

FFFS

$$\begin{aligned}P(X = 4) &= (1 - 0.2)^4(0.2) \\ &= 0.08192\end{aligned}$$

# Bar Graph of Example 1



# Mean and Standard Deviation of Geometric Distributions

The mean (expected value) of a geometric distribution is the **reciprocal of the probability of success**:

$$E(X) = \frac{1}{p}$$

# Mean and Standard Deviation of Geometric Distributions

The mean (expected value) of a geometric distribution is the **reciprocal of the probability of success**:

$$E(X) = \frac{1}{p}$$

The variance for a geometric distribution is the probability of failure divided by the square of the probability of success, given as

$$\sigma^2 = \frac{1 - p}{p^2}$$

# Mean and Standard Deviation of Geometric Distributions

The mean (expected value) of a geometric distribution is the **reciprocal of the probability of success**:

$$E(X) = \frac{1}{p}$$

The variance for a geometric distribution is the probability of failure divided by the square of the probability of success, given as

$$\sigma^2 = \frac{1-p}{p^2}$$

From which the standard deviation,  $\sigma$  is

$$\sigma = \sqrt{\frac{1-p}{p^2}} = \frac{\sqrt{1-p}}{p}$$

## Example 2

(a) A student is given a 10-question multiple choice test in which each question has 5 possible answers. What is the expected number of answers a student can guess before getting one right?

## Example 2

(a) A student is given a 10-question multiple choice test in which each question has 5 possible answers. What is the expected number of answers a student can guess before getting one right?

The expected value is the reciprocal of the probability of success:



## Example 2

(a) A student is given a 10-question multiple choice test in which each question has 5 possible answers. What is the expected number of answers a student can guess before getting one right?

The expected value is the reciprocal of the probability of success:

$$E(X) = \frac{1}{0.2}$$

## Example 2

(a) A student is given a 10-question multiple choice test in which each question has 5 possible answers. What is the expected number of answers a student can guess before getting one right?

The expected value is the reciprocal of the probability of success:

$$\begin{aligned} E(X) &= \frac{1}{0.2} \\ &= 5 \end{aligned}$$

## Example 2

(a) A student is given a 10-question multiple choice test in which each question has 5 possible answers. What is the expected number of answers a student can guess before getting one right?

The expected value is the reciprocal of the probability of success:

$$\begin{aligned} E(X) &= \frac{1}{0.2} \\ &= 5 \end{aligned}$$

We would expect the student to answer 5 questions before guessing one correctly.

## Example 2

(b) What is the standard deviation of the number of questions needed before the student guesses correctly?

## Example 2

(b) What is the standard deviation of the number of questions needed before the student guesses correctly?

The standard deviation is the square root of the probability of failure divided by the probability of success:

## Example 2

(b) What is the standard deviation of the number of questions needed before the student guesses correctly?

The standard deviation is the square root of the probability of failure divided by the probability of success:

$$\sigma = \frac{\sqrt{1 - 0.2}}{0.2}$$

## Example 2

(b) What is the standard deviation of the number of questions needed before the student guesses correctly?

The standard deviation is the square root of the probability of failure divided by the probability of success:

$$\sigma = \frac{\sqrt{1 - 0.2}}{0.2}$$
$$\approx 4.47$$

## Example 2

(b) What is the standard deviation of the number of questions needed before the student guesses correctly?

The standard deviation is the square root of the probability of failure divided by the probability of success:

$$\sigma = \frac{\sqrt{1 - 0.2}}{0.2}$$
$$\approx 4.47$$

The standard deviation is about 4.47 questions.



# Objectives

- 1 Solve problems involving geometric probability distributions
- 2 Solve problems involving hypergeometric probability distributions