

Hypothesis Testing

Single Sample Proportion

Testing a Population Proportion

In this section, we will test a claim regarding a population proportion.

Testing a Population Proportion

In this section, we will test a claim regarding a population proportion.

The sample proportion, \hat{p} is given by

$$\hat{p} = \frac{x}{n} = \frac{\text{number of observed successes}}{\text{sample size}}$$

Testing a Population Proportion

In this section, we will test a claim regarding a population proportion.

The sample proportion, \hat{p} is given by

$$\hat{p} = \frac{x}{n} = \frac{\text{number of observed successes}}{\text{sample size}}$$

with sample standard error

$$\sigma_{\hat{p}} = \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}$$

Test Statistic and Critical Value Method

The test statistic of the population proportion, z , can be found by calculating

$$z = \frac{\hat{p} - p}{\sqrt{p(1 - p)/n}}$$

Test Statistic and Critical Value Method

The test statistic of the population proportion, z , can be found by calculating

$$z = \frac{\hat{p} - p}{\sqrt{p(1-p)/n}}$$
$$= \frac{\text{sample proportion} - \text{population proportion}}{\text{standard error}}$$


Test Statistic and Critical Value Method

The test statistic of the population proportion, z , can be found by calculating


$$z = \frac{\hat{p} - p}{\sqrt{p(1-p)/n}}$$
$$= \frac{\text{sample proportion} - \text{population proportion}}{\text{standard error}}$$

Significance Level	Critical Value
$\alpha = 0.01$	± 2.576
$\alpha = 0.05$	± 1.96
$\alpha = 0.10$	± 1.645


Test Statistic and Critical Value Method



Test Statistic and Critical Value Method



Test Statistic and Critical Value Method



p -Value Method

The area of the rejection region is given by α , the significance level.

p -Value Method

The area of the rejection region is given by α , the significance level.

Recall that the p -value is the probability of obtaining sample results as extreme (or more) than one we got.

p -Value Method

The area of the rejection region is given by α , the significance level.

Recall that the p -value is the probability of obtaining sample results as extreme (or more) than one we got.

If our p -value is lower than α , then our results are **statistically significant**, and would not likely occur by chance if the null hypothesis were true.

p -Value Method

The area of the rejection region is given by α , the significance level.


Recall that the p -value is the probability of obtaining sample results as extreme (or more) than one we got.

If our p -value is lower than α , then our results are **statistically significant**, and would not likely occur by chance if the null hypothesis were true.

Reminder, if the p -value $< \alpha$, we reject the null hypothesis.


p -Value Method

Left-tailed




p -Value Method

Left-tailed




p -Value Method




p -Value Method

Left-tailed




Reject null

Left-tailed




p -Value Method

Left-tailed




Reject null

Left-tailed




p -Value Method

Left-tailed



Reject null


Left-tailed



Do Not Reject Null


Confidence Interval Method

Reject Null



Confidence Interval Method

Reject Null



Do Not Reject Null

