

Dot Product and Projection

Table of Contents

Dot Product

For vectors $\vec{v} = \langle v_1, v_2 \rangle$ and $\vec{w} = \langle w_1, w_2 \rangle$, the **dot product** of \vec{v} and \vec{w} is given as

$$\vec{v} \cdot \vec{w} = \langle v_1, v_2 \rangle \cdot \langle w_1, w_2 \rangle$$

Dot Product

For vectors $\vec{v} = \langle v_1, v_2 \rangle$ and $\vec{w} = \langle w_1, w_2 \rangle$, the **dot product** of \vec{v} and \vec{w} is given as

$$\begin{aligned}\vec{v} \cdot \vec{w} &= \langle v_1, v_2 \rangle \cdot \langle w_1, w_2 \rangle \\ &= v_1 w_1 + v_2 w_2\end{aligned}$$

Example 1

Find $\vec{v} \cdot \vec{w}$ if $\vec{v} = \langle 3, 4 \rangle$ and $\vec{w} = \langle 1, -2 \rangle$.

Example 1

Find $\vec{v} \cdot \vec{w}$ if $\vec{v} = \langle 3, 4 \rangle$ and $\vec{w} = \langle 1, -2 \rangle$.

$$\vec{v} \cdot \vec{w} = \langle 3, 4 \rangle \cdot \langle 1, -2 \rangle$$

Example 1

Find $\vec{v} \cdot \vec{w}$ if $\vec{v} = \langle 3, 4 \rangle$ and $\vec{w} = \langle 1, -2 \rangle$.

$$\begin{aligned}\vec{v} \cdot \vec{w} &= \langle 3, 4 \rangle \cdot \langle 1, -2 \rangle \\ &= 3(1) + 4(-2)\end{aligned}$$

Example 1

Find $\vec{v} \cdot \vec{w}$ if $\vec{v} = \langle 3, 4 \rangle$ and $\vec{w} = \langle 1, -2 \rangle$.

$$\vec{v} \cdot \vec{w} = \langle 3, 4 \rangle \cdot \langle 1, -2 \rangle$$

$$= 3(1) + 4(-2)$$

$$= -5$$

Dot Product Properties

Because the dot product produces a scalar, it is often called the **scalar product**.

Dot Product Properties

Because the dot product produces a scalar, it is often called the **scalar product**.

Properties

Dot Product Properties

Because the dot product produces a scalar, it is often called the **scalar product**.

Properties

$$\vec{v} \cdot \vec{w} = \vec{w} \cdot \vec{v}$$

Commutative Property

Dot Product Properties

Because the dot product produces a scalar, it is often called the **scalar product**.

Properties

$$\vec{v} \cdot \vec{w} = \vec{w} \cdot \vec{v}$$

Commutative Property

$$\vec{u} \cdot (\vec{v} + \vec{w}) = \vec{u} \cdot \vec{v} + \vec{u} \cdot \vec{w}$$

Distributive Property

Dot Product Properties

Because the dot product produces a scalar, it is often called the **scalar product**.

Properties

$$\vec{v} \cdot \vec{w} = \vec{w} \cdot \vec{v}$$

Commutative Property

$$\vec{u} \cdot (\vec{v} + \vec{w}) = \vec{u} \cdot \vec{v} + \vec{u} \cdot \vec{w}$$

Distributive Property

$$k(\vec{v}) \cdot \vec{w} = k(\vec{v} \cdot \vec{w}) = \vec{v} \cdot (k\vec{w})$$

Scalar Property

Dot Product Properties

Because the dot product produces a scalar, it is often called the **scalar product**.

Properties

$$\vec{v} \cdot \vec{w} = \vec{w} \cdot \vec{v}$$

Commutative Property

$$\vec{u} \cdot (\vec{v} + \vec{w}) = \vec{u} \cdot \vec{v} + \vec{u} \cdot \vec{w}$$

Distributive Property

$$k(\vec{v}) \cdot \vec{w} = k(\vec{v} \cdot \vec{w}) = \vec{v} \cdot (k\vec{w})$$

Scalar Property

$$\vec{v} \cdot \vec{v} = \|\vec{v}\|^2$$

Relation to Magnitude

Example 2

Show that $\|\vec{v} - \vec{w}\|^2 = \|\vec{v}\|^2 - 2(\vec{v} \cdot \vec{w}) + \|\vec{w}\|^2$

Example 2

Show that $\|\vec{v} - \vec{w}\|^2 = \|\vec{v}\|^2 - 2(\vec{v} \cdot \vec{w}) + \|\vec{w}\|^2$

$$\|\vec{v} - \vec{w}\| = (\vec{v} - \vec{w}) \cdot (\vec{v} - \vec{w})$$

Example 2

Show that $\|\vec{v} - \vec{w}\|^2 = \|\vec{v}\|^2 - 2(\vec{v} \cdot \vec{w}) + \|\vec{w}\|^2$

$$\begin{aligned}\|\vec{v} - \vec{w}\| &= (\vec{v} - \vec{w}) \cdot (\vec{v} - \vec{w}) \\ &= \vec{v} \cdot (\vec{v} - \vec{w}) - \vec{w} \cdot (\vec{v} - \vec{w})\end{aligned}$$

Example 2

Show that $\|\vec{v} - \vec{w}\|^2 = \|\vec{v}\|^2 - 2(\vec{v} \cdot \vec{w}) + \|\vec{w}\|^2$

$$\begin{aligned}\|\vec{v} - \vec{w}\| &= (\vec{v} - \vec{w}) \cdot (\vec{v} - \vec{w}) \\ &= \vec{v} \cdot (\vec{v} - \vec{w}) - \vec{w} \cdot (\vec{v} - \vec{w}) \\ &= \vec{v} \cdot \vec{v} - \vec{v} \cdot \vec{w} - \vec{v} \cdot \vec{w} + \vec{w} \cdot \vec{w}\end{aligned}$$

Example 2

Show that $\|\vec{v} - \vec{w}\|^2 = \|\vec{v}\|^2 - 2(\vec{v} \cdot \vec{w}) + \|\vec{w}\|^2$

$$\begin{aligned}\|\vec{v} - \vec{w}\| &= (\vec{v} - \vec{w}) \cdot (\vec{v} - \vec{w}) \\&= \vec{v} \cdot (\vec{v} - \vec{w}) - \vec{w} \cdot (\vec{v} - \vec{w}) \\&= \vec{v} \cdot \vec{v} - \vec{v} \cdot \vec{w} - \vec{v} \cdot \vec{w} + \vec{w} \cdot \vec{w} \\&= \vec{v} \cdot \vec{v} - 2(\vec{v} \cdot \vec{w}) + \vec{w} \cdot \vec{w}\end{aligned}$$

Example 2




Show that $\|\vec{v} - \vec{w}\|^2 = \|\vec{v}\|^2 - 2(\vec{v} \cdot \vec{w}) + \|\vec{w}\|^2$

$$\begin{aligned}\|\vec{v} - \vec{w}\| &= (\vec{v} - \vec{w}) \cdot (\vec{v} - \vec{w}) \\&= \vec{v} \cdot (\vec{v} - \vec{w}) - \vec{w} \cdot (\vec{v} - \vec{w}) \\&= \vec{v} \cdot \vec{v} - \vec{v} \cdot \vec{w} - \vec{v} \cdot \vec{w} + \vec{w} \cdot \vec{w} \\&= \vec{v} \cdot \vec{v} - 2(\vec{v} \cdot \vec{w}) + \vec{w} \cdot \vec{w} \\&= \|\vec{v}\|^2 - 2(\vec{v} \cdot \vec{w}) + \|\vec{w}\|^2\end{aligned}$$

Table of Contents

Angles Between Vectors

If we draw \vec{v} and \vec{w} with the same initial point, then the angle θ between them is illustrated below:



Dot Product and Angles Between Vectors



Geometrically, the dot product between two vectors is

$$\vec{v} \cdot \vec{w} = \|\vec{v}\| \|\vec{w}\| \cos \theta$$

Dot Product and Angles Between Vectors

Geometrically, the dot product between two vectors is

$$\vec{v} \cdot \vec{w} = \|\vec{v}\| \|\vec{w}\| \cos \theta$$



Dot Product and Angles Between Vectors

By the Law of Cosines, $\|\vec{v} - \vec{w}\|^2 = \|\vec{v}\|^2 + \|\vec{w}\|^2 - 2\|\vec{v}\|\|\vec{w}\|\cos\theta$,
and by Example 2, $\|\vec{v} - \vec{w}\|^2 = \|\vec{v}\|^2 - 2(\vec{v} \cdot \vec{w}) + \|\vec{w}\|^2$.

Dot Product and Angles Between Vectors

By the Law of Cosines, $\|\vec{v} - \vec{w}\|^2 = \|\vec{v}\|^2 + \|\vec{w}\|^2 - 2\|\vec{v}\|\|\vec{w}\|\cos\theta$,
and by Example 2, $\|\vec{v} - \vec{w}\|^2 = \|\vec{v}\|^2 - 2(\vec{v} \cdot \vec{w}) + \|\vec{w}\|^2$.

Equating these and solving for $\vec{v} \cdot \vec{w}$ gives us $\vec{v} \cdot \vec{w} = \|\vec{v}\|\|\vec{w}\|\cos\theta$,
from which

$$\cos\theta = \frac{\vec{v} \cdot \vec{w}}{\|\vec{v}\|\|\vec{w}\|}$$

Dot Product and Angles Between Vectors

By the Law of Cosines, $\|\vec{v} - \vec{w}\|^2 = \|\vec{v}\|^2 + \|\vec{w}\|^2 - 2\|\vec{v}\|\|\vec{w}\|\cos\theta$,
and by Example 2, $\|\vec{v} - \vec{w}\|^2 = \|\vec{v}\|^2 - 2(\vec{v} \cdot \vec{w}) + \|\vec{w}\|^2$.

Equating these and solving for $\vec{v} \cdot \vec{w}$ gives us $\vec{v} \cdot \vec{w} = \|\vec{v}\|\|\vec{w}\|\cos\theta$,
from which

$$\cos\theta = \frac{\vec{v} \cdot \vec{w}}{\|\vec{v}\|\|\vec{w}\|}$$

To find the angle between two vectors, solve for θ :

$$\theta = \cos^{-1} \left(\frac{\vec{v} \cdot \vec{w}}{\|\vec{v}\|\|\vec{w}\|} \right) = \cos^{-1} (\hat{v} \cdot \hat{w})$$

Example 3a

Find the angle between each of the following.

(a) $\vec{v} = \langle 3, -3\sqrt{3} \rangle$ and $\vec{w} = \langle -\sqrt{3}, 1 \rangle$

Example 3a

Find the angle between each of the following.

(a) $\vec{v} = \langle 3, -3\sqrt{3} \rangle$ and $\vec{w} = \langle -\sqrt{3}, 1 \rangle$

$$\vec{v} \cdot \vec{w} = \langle 3, -3\sqrt{3} \rangle \cdot \langle -\sqrt{3}, 1 \rangle$$

Example 3a

Find the angle between each of the following.

(a) $\vec{v} = \langle 3, -3\sqrt{3} \rangle$ and $\vec{w} = \langle -\sqrt{3}, 1 \rangle$

$$\begin{aligned}\vec{v} \cdot \vec{w} &= \langle 3, -3\sqrt{3} \rangle \cdot \langle -\sqrt{3}, 1 \rangle \\ &= -3\sqrt{3} - 3\sqrt{3} = -6\sqrt{3}\end{aligned}$$

Example 3a

Find the angle between each of the following.

(a) $\vec{v} = \langle 3, -3\sqrt{3} \rangle$ and $\vec{w} = \langle -\sqrt{3}, 1 \rangle$

$$\begin{aligned}\vec{v} \cdot \vec{w} &= \langle 3, -3\sqrt{3} \rangle \cdot \langle -\sqrt{3}, 1 \rangle \\ &= -3\sqrt{3} - 3\sqrt{3} = -6\sqrt{3}\end{aligned}$$

$$\|\vec{v}\| = \sqrt{3^2 + (3\sqrt{3})^2} = \sqrt{36}$$

Example 3a

Find the angle between each of the following.

(a) $\vec{v} = \langle 3, -3\sqrt{3} \rangle$ and $\vec{w} = \langle -\sqrt{3}, 1 \rangle$

$$\begin{aligned}\vec{v} \cdot \vec{w} &= \langle 3, -3\sqrt{3} \rangle \cdot \langle -\sqrt{3}, 1 \rangle \\ &= -3\sqrt{3} - 3\sqrt{3} = -6\sqrt{3}\end{aligned}$$

$$\|\vec{v}\| = \sqrt{3^2 + (3\sqrt{3})^2} = \sqrt{36}$$

$$\|\vec{w}\| = \sqrt{(\sqrt{3})^2 + 1^2} = \sqrt{4}$$

Example 3a

Find the angle between each of the following.

(a) $\vec{v} = \langle 3, -3\sqrt{3} \rangle$ and $\vec{w} = \langle -\sqrt{3}, 1 \rangle$

$$\begin{aligned}\vec{v} \cdot \vec{w} &= \langle 3, -3\sqrt{3} \rangle \cdot \langle -\sqrt{3}, 1 \rangle \\ &= -3\sqrt{3} - 3\sqrt{3} = -6\sqrt{3}\end{aligned}$$

$$\|\vec{v}\| = \sqrt{3^2 + (3\sqrt{3})^2} = \sqrt{36}$$

$$\|\vec{w}\| = \sqrt{(\sqrt{3})^2 + 1^2} = \sqrt{4}$$

$$\theta = \arccos \left(\frac{-6\sqrt{3}}{\sqrt{36} \times 4} \right) = \frac{5\pi}{6}$$

Example 3b

(b) $\vec{v} = \langle 2, 2 \rangle$ and $\vec{w} = \langle 5, -5 \rangle$

Example 3b

$$(b) \quad \vec{v} = \langle 2, 2 \rangle \text{ and } \vec{w} = \langle 5, -5 \rangle$$

$$\vec{v} \cdot \vec{w} = 2(5) + 2(-5) = 0$$

Example 3b

$$(b) \quad \vec{v} = \langle 2, 2 \rangle \text{ and } \vec{w} = \langle 5, -5 \rangle$$

$$\vec{v} \cdot \vec{w} = 2(5) + 2(-5) = 0$$

$$\|\vec{v}\| = \sqrt{2^2 + 2^2} = \sqrt{8}$$

Example 3b

$$(b) \quad \vec{v} = \langle 2, 2 \rangle \text{ and } \vec{w} = \langle 5, -5 \rangle$$

$$\vec{v} \cdot \vec{w} = 2(5) + 2(-5) = 0$$

$$\|\vec{v}\| = \sqrt{2^2 + 2^2} = \sqrt{8}$$

$$\|\vec{w}\| = \sqrt{5^2 + 5^2} = \sqrt{50}$$

Example 3b

$$(b) \quad \vec{v} = \langle 2, 2 \rangle \text{ and } \vec{w} = \langle 5, -5 \rangle$$

$$\vec{v} \cdot \vec{w} = 2(5) + 2(-5) = 0$$

$$\|\vec{v}\| = \sqrt{2^2 + 2^2} = \sqrt{8}$$

$$\|\vec{w}\| = \sqrt{5^2 + 5^2} = \sqrt{50}$$

$$\theta = \arccos\left(\frac{0}{\sqrt{8 \times 50}}\right) = \frac{\pi}{2}$$

Example 3c

(c) $\vec{v} = \langle 3, -4 \rangle$ and $\vec{w} = \langle 2, 1 \rangle$

Example 3c

(c) $\vec{v} = \langle 3, -4 \rangle$ and $\vec{w} = \langle 2, 1 \rangle$

$$\vec{v} \cdot \vec{w} = 3(2) + (-4)(1) = 2$$

Example 3c

$$(c) \quad \vec{v} = \langle 3, -4 \rangle \text{ and } \vec{w} = \langle 2, 1 \rangle$$

$$\vec{v} \cdot \vec{w} = 3(2) + (-4)(1) = 2$$

$$\|\vec{v}\| = \sqrt{3^2 + 4^2} = \sqrt{25}$$

Example 3c

$$(c) \quad \vec{v} = \langle 3, -4 \rangle \text{ and } \vec{w} = \langle 2, 1 \rangle$$

$$\vec{v} \cdot \vec{w} = 3(2) + (-4)(1) = 2$$

$$\|\vec{v}\| = \sqrt{3^2 + 4^2} = \sqrt{25}$$

$$\|\vec{w}\| = \sqrt{2^2 + 1^2} = \sqrt{5}$$

Example 3c

$$(c) \quad \vec{v} = \langle 3, -4 \rangle \text{ and } \vec{w} = \langle 2, 1 \rangle$$

$$\vec{v} \cdot \vec{w} = 3(2) + (-4)(1) = 2$$

$$\|\vec{v}\| = \sqrt{3^2 + 4^2} = \sqrt{25}$$

$$\|\vec{w}\| = \sqrt{2^2 + 1^2} = \sqrt{5}$$

$$\theta = \arccos \left(\frac{2}{\sqrt{25 \times 5}} \right) \approx 80^\circ$$

Table of Contents

Orthogonal Projection

Two vectors are **orthogonal** if they meet at a right angle.

If two vectors are orthogonal, then their dot product is 0.



Orthogonal Projection

The **orthogonal projection** of \vec{v} onto \vec{w} is a new vector \vec{p} that is parallel to \vec{w} and has a magnitude of $\|\vec{v}\| \cos \theta$.

Orthogonal Projection

The **orthogonal projection** of \vec{v} onto \vec{w} is a new vector \vec{p} that is parallel to \vec{w} and has a magnitude of $\|\vec{v}\| \cos \theta$.

\vec{p} can be thought of as the “shadow” \vec{v} casts over \vec{w} :



Orthogonal Projection

We need \vec{p} to be parallel to \vec{w} . To do this, we could take the dot product of \vec{p} and \vec{w} .

Orthogonal Projection

We need \vec{p} to be parallel to \vec{w} . To do this, we could take the dot product of \vec{p} and \vec{w} .

However, that would likely change the magnitude of \vec{p} .

Orthogonal Projection

We need \vec{p} to be parallel to \vec{w} . To do this, we could take the dot product of \vec{p} and \vec{w} .

However, that would likely change the magnitude of \vec{p} .

Recall that when finding a unit vector for a given vector, that unit vector has a magnitude of 1 and is parallel to the original vector.

Orthogonal Projection

We need \vec{p} to be parallel to \vec{w} . To do this, we could take the dot product of \vec{p} and \vec{w} .

However, that would likely change the magnitude of \vec{p} .

Recall that when finding a unit vector for a given vector, that unit vector has a magnitude of 1 and is parallel to the original vector.

Thus, we can multiply the magnitude of \vec{p} by a unit vector for \vec{w} .

$$\|\vec{p}\| \hat{w}$$

Orthogonal Projection

We need \vec{p} to be parallel to \vec{w} . To do this, we could take the dot product of \vec{p} and \vec{w} .

However, that would likely change the magnitude of \vec{p} .


Recall that when finding a unit vector for a given vector, that unit vector has a magnitude of 1 and is parallel to the original vector.

Thus, we can multiply the magnitude of \vec{p} by a unit vector for \vec{w} .

$$\|\vec{p}\| \hat{w}$$


This will guarantee that \vec{w} is scaled to \vec{p} and give us the projection of \vec{v} onto \vec{w} .

Orthogonal Projection



Orthogonal Projection


$$\text{proj}_{\vec{w}} \vec{v} = \|\vec{p}\| \hat{w}$$




Orthogonal Projection

$$\text{proj}_{\vec{w}} \vec{v} = \|\vec{p}\| \hat{w}$$

$$= (\|v\| \cos \theta) \left(\frac{\vec{w}}{\|w\|} \right)$$



Orthogonal Projection




$$\text{proj}_{\vec{w}} \vec{v} = \|\vec{p}\| \hat{w}$$

$$= (\|\vec{v}\| \cos \theta) \left(\frac{\vec{w}}{\|\vec{w}\|} \right)$$

$$= \|\vec{v}\| \left(\frac{\vec{v} \cdot \vec{w}}{\|\vec{v}\| \|\vec{w}\|} \right) \left(\frac{\vec{w}}{\|\vec{w}\|} \right)$$

Orthogonal Projection




$$\text{proj}_{\vec{w}} \vec{v} = \|\vec{p}\| \hat{w}$$

$$= (\|\vec{v}\| \cos \theta) \left(\frac{\vec{w}}{\|\vec{w}\|} \right)$$

$$= \|\vec{v}\| \left(\frac{\vec{v} \cdot \vec{w}}{\|\vec{v}\| \|\vec{w}\|} \right) \left(\frac{\vec{w}}{\|\vec{w}\|} \right)$$

$$= \left(\frac{\vec{v} \cdot \vec{w}}{\|\vec{w}\| \|\vec{w}\|} \right) \vec{w}$$

Orthogonal Projection



$$\text{proj}_{\vec{w}} \vec{v} = \|\vec{p}\| \hat{w}$$

$$= (\|\vec{v}\| \cos \theta) \left(\frac{\vec{w}}{\|\vec{w}\|} \right)$$

$$= \|\vec{v}\| \left(\frac{\vec{v} \cdot \vec{w}}{\|\vec{v}\| \|\vec{w}\|} \right) \left(\frac{\vec{w}}{\|\vec{w}\|} \right)$$

$$= \left(\frac{\vec{v} \cdot \vec{w}}{\|\vec{w}\| \|\vec{w}\|} \right) \vec{w}$$

$$\text{proj}_{\vec{w}} \vec{v} = \left(\frac{\vec{v} \cdot \vec{w}}{\|\vec{w}\|^2} \right) \vec{w}$$

Example 4a

Let $\vec{v} = \langle 1, 8 \rangle$ and $\vec{w} = \langle -1, 2 \rangle$. Find each of the following.

(a) $\text{proj}_{\vec{w}} \vec{v}$

Example 4a

Let $\vec{v} = \langle 1, 8 \rangle$ and $\vec{w} = \langle -1, 2 \rangle$. Find each of the following.

(a) $\text{proj}_{\vec{w}} \vec{v}$

$$\text{proj}_{\vec{w}} \vec{v} = \left(\frac{\vec{v} \cdot \vec{w}}{\|\vec{w}\|^2} \right) \vec{w}$$

Example 4a

Let $\vec{v} = \langle 1, 8 \rangle$ and $\vec{w} = \langle -1, 2 \rangle$. Find each of the following.

(a) $\text{proj}_{\vec{w}} \vec{v}$

$$\text{proj}_{\vec{w}} \vec{v} = \left(\frac{\vec{v} \cdot \vec{w}}{\|\vec{w}\|^2} \right) \vec{w}$$

$$\vec{v} \cdot \vec{w} = 1(-1) + 8(2) = 15$$

Example 4a

Let $\vec{v} = \langle 1, 8 \rangle$ and $\vec{w} = \langle -1, 2 \rangle$. Find each of the following.

(a) $\text{proj}_{\vec{w}} \vec{v}$

$$\text{proj}_{\vec{w}} \vec{v} = \left(\frac{\vec{v} \cdot \vec{w}}{\|\vec{w}\|^2} \right) \vec{w}$$

$$\vec{v} \cdot \vec{w} = 1(-1) + 8(2) = 15 \qquad \|\vec{w}\|^2 = \left(\sqrt{1^2 + 2^2} \right)^2 = 5$$

Example 4a

Let $\vec{v} = \langle 1, 8 \rangle$ and $\vec{w} = \langle -1, 2 \rangle$. Find each of the following.

(a) $\text{proj}_{\vec{w}} \vec{v}$

$$\text{proj}_{\vec{w}} \vec{v} = \left(\frac{\vec{v} \cdot \vec{w}}{\|\vec{w}\|^2} \right) \vec{w}$$

$$\vec{v} \cdot \vec{w} = 1(-1) + 8(2) = 15 \qquad \|\vec{w}\|^2 = \left(\sqrt{1^2 + 2^2} \right)^2 = 5$$

$$\text{proj}_{\vec{w}} \vec{v} = \frac{15}{5} \langle -1, 2 \rangle$$

Example 4a

Let $\vec{v} = \langle 1, 8 \rangle$ and $\vec{w} = \langle -1, 2 \rangle$. Find each of the following.

(a) $\text{proj}_{\vec{w}} \vec{v}$

$$\text{proj}_{\vec{w}} \vec{v} = \left(\frac{\vec{v} \cdot \vec{w}}{\|\vec{w}\|^2} \right) \vec{w}$$

$$\vec{v} \cdot \vec{w} = 1(-1) + 8(2) = 15 \qquad \|\vec{w}\|^2 = \left(\sqrt{1^2 + 2^2} \right)^2 = 5$$

$$\begin{aligned} \text{proj}_{\vec{w}} \vec{v} &= \frac{15}{5} \langle -1, 2 \rangle \\ &= 3 \langle -1, 2 \rangle \end{aligned}$$

Example 4a

Let $\vec{v} = \langle 1, 8 \rangle$ and $\vec{w} = \langle -1, 2 \rangle$. Find each of the following.

(a) $\text{proj}_{\vec{w}} \vec{v}$

$$\text{proj}_{\vec{w}} \vec{v} = \left(\frac{\vec{v} \cdot \vec{w}}{\|\vec{w}\|^2} \right) \vec{w}$$

$$\vec{v} \cdot \vec{w} = 1(-1) + 8(2) = 15 \qquad \|\vec{w}\|^2 = \left(\sqrt{1^2 + 2^2} \right)^2 = 5$$

$$\text{proj}_{\vec{w}} \vec{v} = \frac{15}{5} \langle -1, 2 \rangle$$

$$= 3 \langle -1, 2 \rangle$$

$$= \langle -3, 6 \rangle$$

Example 4b

Let $\vec{v} = \langle 1, 8 \rangle$ and $\vec{w} = \langle -1, 2 \rangle$. Find each of the following.

(b) $\text{proj}_{\vec{v}} \vec{w}$

Example 4b

Let $\vec{v} = \langle 1, 8 \rangle$ and $\vec{w} = \langle -1, 2 \rangle$. Find each of the following.

(b) $\text{proj}_{\vec{v}} \vec{w}$

$$\text{proj}_{\vec{v}} \vec{w} = \left(\frac{\vec{w} \cdot \vec{v}}{\|\vec{v}\|^2} \right) \vec{v}$$

Example 4b

Let $\vec{v} = \langle 1, 8 \rangle$ and $\vec{w} = \langle -1, 2 \rangle$. Find each of the following.

(b) $\text{proj}_{\vec{v}} \vec{w}$

$$\text{proj}_{\vec{v}} \vec{w} = \left(\frac{\vec{w} \cdot \vec{v}}{\|\vec{v}\|^2} \right) \vec{v}$$

$$\vec{w} \cdot \vec{v} = 1(-1) + 8(2) = 15$$

Example 4b

Let $\vec{v} = \langle 1, 8 \rangle$ and $\vec{w} = \langle -1, 2 \rangle$. Find each of the following.

(b) $\text{proj}_{\vec{v}} \vec{w}$

$$\text{proj}_{\vec{v}} \vec{w} = \left(\frac{\vec{w} \cdot \vec{v}}{\|\vec{v}\|^2} \right) \vec{v}$$

$$\vec{w} \cdot \vec{v} = 1(-1) + 8(2) = 15 \qquad \|\vec{v}\|^2 = \left(\sqrt{1^2 + 8^2} \right)^2 = 65$$

Example 4b

Let $\vec{v} = \langle 1, 8 \rangle$ and $\vec{w} = \langle -1, 2 \rangle$. Find each of the following.

(b) $\text{proj}_{\vec{v}} \vec{w}$

$$\text{proj}_{\vec{v}} \vec{w} = \left(\frac{\vec{w} \cdot \vec{v}}{\|\vec{v}\|^2} \right) \vec{v}$$

$$\vec{w} \cdot \vec{v} = 1(-1) + 8(2) = 15 \quad \|\vec{v}\|^2 = \left(\sqrt{1^2 + 8^2} \right)^2 = 65$$

$$\text{proj}_{\vec{v}} \vec{w} = \frac{3}{13} \langle 1, 8 \rangle$$

Example 4b

Let $\vec{v} = \langle 1, 8 \rangle$ and $\vec{w} = \langle -1, 2 \rangle$. Find each of the following.

(b) $\text{proj}_{\vec{v}} \vec{w}$

$$\text{proj}_{\vec{v}} \vec{w} = \left(\frac{\vec{w} \cdot \vec{v}}{\|\vec{v}\|^2} \right) \vec{v}$$

$$\vec{w} \cdot \vec{v} = 1(-1) + 8(2) = 15 \quad \|\vec{v}\|^2 = \left(\sqrt{1^2 + 8^2} \right)^2 = 65$$

$$\begin{aligned} \text{proj}_{\vec{v}} \vec{w} &= \frac{3}{13} \langle 1, 8 \rangle \\ &= \left\langle \frac{3}{13}, \frac{24}{13} \right\rangle \end{aligned}$$

Table of Contents

Work

In physics, if a constant force F is exerted over a distance (really, *displacement*) d , the work W done by the force is given by $W = Fd$.

Work


In physics, if a constant force F is exerted over a distance (really, *displacement*) d , the work W done by the force is given by $W = Fd$.

However, if the force being applied is not in the direction of motion, we can use the dot product to find the work done.

Work

In physics, if a constant force F is exerted over a distance (really, *displacement*) d , the work W done by the force is given by $W = Fd$.

However, if the force being applied is not in the direction of motion, we can use the dot product to find the work done.



$$W = Fd = \|\vec{F}\| \|\vec{d}\| \cos \theta = \vec{F} \cdot \vec{d}$$

Example 5

Mr. Bain exerts a force of 230 pounds to pull a stack of rocks a distance of 50 ft. over level ground. If the rope makes a 30° angle with the horizontal, how much work did Mr. Bain do pulling the rocks? Assume Mr. Bain exerts the force of 230 pounds at a 30° angle for the duration of the 50 feet.

Example 5

Mr. Bain exerts a force of 230 pounds to pull a stack of rocks a distance of 50 ft. over level ground. If the rope makes a 30° angle with the horizontal, how much work did Mr. Bain do pulling the rocks? Assume Mr. Bain exerts the force of 230 pounds at a 30° angle for the duration of the 50 feet.

$$W = \|F\| \|d\| \cos \theta$$

Example 5

Mr. Bain exerts a force of 230 pounds to pull a stack of rocks a distance of 50 ft. over level ground. If the rope makes a 30° angle with the horizontal, how much work did Mr. Bain do pulling the rocks? Assume Mr. Bain exerts the force of 230 pounds at a 30° angle for the duration of the 50 feet.

$$\begin{aligned} W &= \|F\| \|d\| \cos \theta \\ &= (230 \text{ pounds})(50 \text{ ft}) \cos 30^\circ \end{aligned}$$

Example 5

Mr. Bain exerts a force of 230 pounds to pull a stack of rocks a distance of 50 ft. over level ground. If the rope makes a 30° angle with the horizontal, how much work did Mr. Bain do pulling the rocks? Assume Mr. Bain exerts the force of 230 pounds at a 30° angle for the duration of the 50 feet.

$$\begin{aligned} W &= \|F\| \|d\| \cos \theta \\ &= (230 \text{ pounds})(50 \text{ ft}) \cos 30^\circ \\ &= 11,500 \text{ foot-pounds} \left(\frac{\sqrt{3}}{2} \right) \end{aligned}$$

Example 5

Mr. Bain exerts a force of 230 pounds to pull a stack of rocks a distance of 50 ft. over level ground. If the rope makes a 30° angle with the horizontal, how much work did Mr. Bain do pulling the rocks? Assume Mr. Bain exerts the force of 230 pounds at a 30° angle for the duration of the 50 feet.

$$\begin{aligned} W &= \|F\| \|d\| \cos \theta \\ &= (230 \text{ pounds})(50 \text{ ft}) \cos 30^\circ \\ &= 11,500 \text{ foot-pounds} \left(\frac{\sqrt{3}}{2} \right) \\ &= 5,750\sqrt{3} (\approx 9,959.3) \text{ foot-pounds} \end{aligned}$$