

## Polar Form of Conics

# Objectives

- 1 Analyze the graphs of conic sections in polar form.

# Intro

Given a fixed line  $L$ , a point  $F$  not on  $L$ , and a positive number  $e$ , a **conic section** is the set of all points  $P$  such that

$$\frac{\text{the distance from } P \text{ to } F}{\text{the distance from } P \text{ to } L} = e$$


Given a fixed line  $L$ , a point  $F$  not on  $L$ , and a positive number  $e$ , a **conic section** is the set of all points  $P$  such that

$$\frac{\text{the distance from } P \text{ to } F}{\text{the distance from } P \text{ to } L} = e$$

The line  $L$  is called the **directrix** of the conic section, the point  $F$  is called a **focus** of the conic section, and the constant  $e$  is called the **eccentricity** of the conic section.

# Eccentricity, Focus, and Directrix Line

The conic section has eccentricity  $e$ , a focus  $F$  at the origin and directrix line  $x = -d$ :



# General Equation

From which we get

$$e = \frac{\text{the distance from } P \text{ to } F}{\text{the distance from } P \text{ to } L} = \frac{r}{d + r \cos \theta} = e$$

# General Equation

From which we get

$$e = \frac{\text{the distance from } P \text{ to } F}{\text{the distance from } P \text{ to } L} = \frac{r}{d + r \cos \theta} = e$$

So that  $r = e(d + r \cos \theta)$ , and solving for  $r$  gives us

# General Equation

From which we get

$$e = \frac{\text{the distance from } P \text{ to } F}{\text{the distance from } P \text{ to } L} = \frac{r}{d + r \cos \theta} = e$$

So that  $r = e(d + r \cos \theta)$ , and solving for  $r$  gives us

$$r = ed + er \cos \theta$$



# General Equation

From which we get

$$e = \frac{\text{the distance from } P \text{ to } F}{\text{the distance from } P \text{ to } L} = \frac{r}{d + r \cos \theta} = e$$

So that  $r = e(d + r \cos \theta)$ , and solving for  $r$  gives us

$$r = ed + er \cos \theta$$

$$r - er \cos \theta = ed$$

# General Equation

From which we get

$$e = \frac{\text{the distance from } P \text{ to } F}{\text{the distance from } P \text{ to } L} = \frac{r}{d + r \cos \theta} = e$$

So that  $r = e(d + r \cos \theta)$ , and solving for  $r$  gives us

$$r = ed + er \cos \theta$$

$$r - er \cos \theta = ed$$

$$r(1 - e \cos \theta) = ed$$

# General Equation

From which we get

$$e = \frac{\text{the distance from } P \text{ to } F}{\text{the distance from } P \text{ to } L} = \frac{r}{d + r \cos \theta} = e$$

So that  $r = e(d + r \cos \theta)$ , and solving for  $r$  gives us

$$r = ed + er \cos \theta$$

$$r - er \cos \theta = ed$$

$$r(1 - e \cos \theta) = ed$$

$$r = \frac{ed}{1 - e \cos \theta}$$

## Example 1

Examine the graphs of each of the following for different values of  $d$ , but with  $e = 1$ .

$$(a) \quad r = \frac{ed}{1 + e \cos \theta}$$

## Example 1

Examine the graphs of each of the following for different values of  $d$ , but with  $e = 1$ .

$$(a) \quad r = \frac{ed}{1 + e \cos \theta} \quad \longrightarrow \quad r = \frac{d}{1 + \cos \theta}$$

## Example 1

Examine the graphs of each of the following for different values of  $d$ , but with  $e = 1$ .

$$(a) \quad r = \frac{ed}{1 + e \cos \theta} \quad \longrightarrow \quad r = \frac{d}{1 + \cos \theta}$$

- Parabola (opens left or right)

## Example 1

Examine the graphs of each of the following for different values of  $d$ , but with  $e = 1$ .

$$(a) \quad r = \frac{ed}{1 + e \cos \theta} \quad \longrightarrow \quad r = \frac{d}{1 + \cos \theta}$$

- Parabola (opens left or right)
- Vertex at  $\left(\frac{1}{2}d, 0\right)$

## Example 1

Examine the graphs of each of the following for different values of  $d$ , but with  $e = 1$ .

$$(a) \quad r = \frac{ed}{1 + e \cos \theta} \quad \longrightarrow \quad r = \frac{d}{1 + \cos \theta}$$

- Parabola (opens left or right)
- Vertex at  $\left(\frac{1}{2}d, 0\right)$
- $d > 0$  opens left



## Example 1

Examine the graphs of each of the following for different values of  $d$ , but with  $e = 1$ .

$$(a) \quad r = \frac{ed}{1 + e \cos \theta} \quad \longrightarrow \quad r = \frac{d}{1 + \cos \theta}$$

- Parabola (opens left or right)
- Vertex at  $\left(\frac{1}{2}d, 0\right)$
- $d > 0$  opens left
- $d < 0$  opens right

## Example 1

Examine the graphs of each of the following for different values of  $d$ , but with  $e = 1$ .

$$(a) \quad r = \frac{ed}{1 + e \cos \theta} \quad \longrightarrow \quad r = \frac{d}{1 + \cos \theta}$$

- Parabola (opens left or right)
- Vertex at  $\left(\frac{1}{2}d, 0\right)$
- $d > 0$  opens left
- $d < 0$  opens right
- $y$ -intercepts at  $(0, \pm d)$

## Example 1

$$(b) \quad r = \frac{ed}{1 - e \cos \theta}$$

## Example 1

$$(b) \quad r = \frac{ed}{1 - e \cos \theta} \quad \longrightarrow \quad r = \frac{d}{1 - \cos \theta}$$

## Example 1

$$(b) \quad r = \frac{ed}{1 - e \cos \theta} \quad \longrightarrow \quad r = \frac{d}{1 - \cos \theta}$$

- Parabola (opens left or right)

## Example 1

$$(b) \quad r = \frac{ed}{1 - e \cos \theta} \quad \longrightarrow \quad r = \frac{d}{1 - \cos \theta}$$

- Parabola (opens left or right)
- Vertex at  $\left(-\frac{1}{2}d, 0\right)$

## Example 1

$$(b) \quad r = \frac{ed}{1 - e \cos \theta} \quad \longrightarrow \quad r = \frac{d}{1 - \cos \theta}$$

- Parabola (opens left or right)
- Vertex at  $\left(-\frac{1}{2}d, 0\right)$
- $d > 0$  opens right

## Example 1

$$(b) \quad r = \frac{ed}{1 - e \cos \theta} \quad \longrightarrow \quad r = \frac{d}{1 - \cos \theta}$$

- Parabola (opens left or right)
- Vertex at  $\left(-\frac{1}{2}d, 0\right)$
- $d > 0$  opens right
- $d < 0$  opens left



## Example 1

$$(b) \quad r = \frac{ed}{1 - e \cos \theta} \quad \longrightarrow \quad r = \frac{d}{1 - \cos \theta}$$

- Parabola (opens left or right)
- Vertex at  $\left(-\frac{1}{2}d, 0\right)$
- $d > 0$  opens right
- $d < 0$  opens left
- $y$ -intercepts at  $(0, \pm d)$

## Example 1

$$(c) \quad r = \frac{ed}{1 + e \sin \theta}$$

## Example 1

$$(c) \quad r = \frac{ed}{1 + e \sin \theta} \quad \longrightarrow \quad r = \frac{d}{1 + \sin \theta}$$

## Example 1

$$(c) \quad r = \frac{ed}{1 + e \sin \theta} \quad \longrightarrow \quad r = \frac{d}{1 + \sin \theta}$$

- Parabola (opens up or down)

## Example 1

$$(c) \quad r = \frac{ed}{1 + e \sin \theta} \quad \longrightarrow \quad r = \frac{d}{1 + \sin \theta}$$

- Parabola (opens up or down)
- Vertex at  $\left(0, \frac{1}{2}d\right)$

## Example 1

$$(c) \quad r = \frac{ed}{1 + e \sin \theta} \quad \longrightarrow \quad r = \frac{d}{1 + \sin \theta}$$

- Parabola (opens up or down)
- Vertex at  $\left(0, \frac{1}{2}d\right)$
- $d > 0$  opens down

## Example 1

$$(c) \quad r = \frac{ed}{1 + e \sin \theta} \quad \longrightarrow \quad r = \frac{d}{1 + \sin \theta}$$

- Parabola (opens up or down)
- Vertex at  $\left(0, \frac{1}{2}d\right)$
- $d > 0$  opens down
- $d < 0$  opens up

## Example 1

$$(c) \quad r = \frac{ed}{1 + e \sin \theta} \quad \longrightarrow \quad r = \frac{d}{1 + \sin \theta}$$

- Parabola (opens up or down)
- Vertex at  $\left(0, \frac{1}{2}d\right)$
- $d > 0$  opens down
- $d < 0$  opens up
- $x$ -intercepts at  $(\pm d, 0)$



## Example 1

$$(d) \quad r = \frac{ed}{1 - e \sin \theta}$$

## Example 1

$$(d) \quad r = \frac{ed}{1 - e \sin \theta} \quad \longrightarrow \quad r = \frac{d}{1 - \sin \theta}$$

## Example 1

$$(d) \quad r = \frac{ed}{1 - e \sin \theta} \quad \longrightarrow \quad r = \frac{d}{1 - \sin \theta}$$

- Parabola (opens up or down)

## Example 1

$$(d) \quad r = \frac{ed}{1 - e \sin \theta} \quad \longrightarrow \quad r = \frac{d}{1 - \sin \theta}$$

- Parabola (opens up or down)
- Vertex at  $\left(0, -\frac{1}{2}d\right)$

## Example 1

$$(d) \quad r = \frac{ed}{1 - e \sin \theta} \quad \longrightarrow \quad r = \frac{d}{1 - \sin \theta}$$

- Parabola (opens up or down)
- Vertex at  $\left(0, -\frac{1}{2}d\right)$
- $d > 0$  opens up

## Example 1

$$(d) \quad r = \frac{ed}{1 - e \sin \theta} \quad \longrightarrow \quad r = \frac{d}{1 - \sin \theta}$$

- Parabola (opens up or down)
- Vertex at  $\left(0, -\frac{1}{2}d\right)$
- $d > 0$  opens up
- $d < 0$  opens down

## Example 1

$$(d) \quad r = \frac{ed}{1 - e \sin \theta} \quad \longrightarrow \quad r = \frac{d}{1 - \sin \theta}$$

- Parabola (opens up or down)
- Vertex at  $\left(0, -\frac{1}{2}d\right)$
- $d > 0$  opens up
- $d < 0$  opens down
- $x$ -intercepts at  $(\pm d, 0)$

## Follow-up to Example 1

Notice each of the previous graphs in Example 1 were parabolas. This is the case when  $e = 1$ . The directrix lines were either  $x = \pm d$  or  $y = \pm d$ , and the focal diameter is  $2d$ .



## Example 2

Examine the graphs of each of the following for different values of  $d$ , but with  $0 < e < 1$  and  $e > 1$ .

(a) 
$$r = \frac{ed}{1 + e \cos \theta}$$

## Example 2

Examine the graphs of each of the following for different values of  $d$ , but with  $0 < e < 1$  and  $e > 1$ .

$$(a) \quad r = \frac{ed}{1 + e \cos \theta}$$

For  $0 < e < 1$ :

## Example 2

Examine the graphs of each of the following for different values of  $d$ , but with  $0 < e < 1$  and  $e > 1$ .

$$(a) \quad r = \frac{ed}{1 + e \cos \theta}$$

For  $0 < e < 1$ :

Ellipse (wide)

## Example 2

$$(a) \quad r = \frac{ed}{1 + e \cos \theta}$$

## Example 2

$$(a) \quad r = \frac{ed}{1 + e \cos \theta}$$

For  $e > 1$ :

## Example 2

$$(a) \quad r = \frac{ed}{1 + e \cos \theta}$$

For  $e > 1$ :

Hyperbola (opening left and right)

## Example 2

Examine the graphs of each of the following for different values of  $d$ , but with  $0 < e < 1$  and  $e > 1$ .

$$(b) \quad r = \frac{ed}{1 - e \cos \theta}$$

## Example 2

Examine the graphs of each of the following for different values of  $d$ , but with  $0 < e < 1$  and  $e > 1$ .

$$(b) \quad r = \frac{ed}{1 - e \cos \theta}$$

For  $0 < e < 1$ :



## Example 2

Examine the graphs of each of the following for different values of  $d$ , but with  $0 < e < 1$  and  $e > 1$ .

$$(b) \quad r = \frac{ed}{1 - e \cos \theta}$$

For  $0 < e < 1$ :

Ellipse (wide)

## Example 2

$$(b) \quad r = \frac{ed}{1 - e \cos \theta}$$

## Example 2

$$(b) \quad r = \frac{ed}{1 - e \cos \theta}$$

For  $e > 1$ :

## Example 2

$$(b) \quad r = \frac{ed}{1 - e \cos \theta}$$

For  $e > 1$ :

Hyperbola (opening left and right)

## Example 2

Examine the graphs of each of the following for different values of  $d$ , but with  $0 < e < 1$  and  $e > 1$ .

$$(c) \quad r = \frac{ed}{1 + e \sin \theta}$$

## Example 2

Examine the graphs of each of the following for different values of  $d$ , but with  $0 < e < 1$  and  $e > 1$ .

$$(c) \quad r = \frac{ed}{1 + e \sin \theta}$$

For  $0 < e < 1$ :

## Example 2

Examine the graphs of each of the following for different values of  $d$ , but with  $0 < e < 1$  and  $e > 1$ .

$$(c) \quad r = \frac{ed}{1 + e \sin \theta}$$

For  $0 < e < 1$ :

Ellipse (tall)

## Example 2

$$(c) \quad r = \frac{ed}{1 + e \sin \theta}$$



## Example 2

$$(c) \quad r = \frac{ed}{1 + e \sin \theta}$$

For  $e > 1$ :

## Example 2

$$(c) \quad r = \frac{ed}{1 + e \sin \theta}$$

For  $e > 1$ :

Hyperbola (opening up and down)

## Example 2

Examine the graphs of each of the following for different values of  $d$ , but with  $0 < e < 1$  and  $e > 1$ .

$$(d) \quad r = \frac{ed}{1 - e \sin \theta}$$

## Example 2

Examine the graphs of each of the following for different values of  $d$ , but with  $0 < e < 1$  and  $e > 1$ .

$$(d) \quad r = \frac{ed}{1 - e \sin \theta}$$

For  $0 < e < 1$ :

## Example 2

Examine the graphs of each of the following for different values of  $d$ , but with  $0 < e < 1$  and  $e > 1$ .

$$(d) \quad r = \frac{ed}{1 - e \sin \theta}$$

For  $0 < e < 1$ :

Ellipse (tall)

## Example 2

$$(d) \quad r = \frac{ed}{1 - e \sin \theta}$$

## Example 2

$$(d) \quad r = \frac{ed}{1 - e \sin \theta}$$

For  $e > 1$ :

## Example 2

$$(d) \quad r = \frac{ed}{1 - e \sin \theta}$$

For  $e > 1$ :

Hyperbola (opening up and down)



## Properties From Example 2

In the previous example, the graphs in which  $0 < e < 1$  were ellipses.

## Properties From Example 2

In the previous example, the graphs in which  $0 < e < 1$  were ellipses.

Major axis length is  $\frac{2ed}{1 - e^2}$

## Properties From Example 2

In the previous example, the graphs in which  $0 < e < 1$  were ellipses.

Major axis length is  $\frac{2ed}{1 - e^2}$

Minor axis length is  $\frac{2ed}{\sqrt{1 - e^2}}$ .

## Properties From Example 2

If  $e > 1$ , the graph is a hyperbola.

## Properties From Example 2

If  $e > 1$ , the graph is a hyperbola.

Transverse axis length  $\frac{2ed}{e^2 - 1}$

## Properties From Example 2

If  $e > 1$ , the graph is a hyperbola.

Transverse axis length  $\frac{2ed}{e^2 - 1}$

Conjugate axis length  $\frac{2ed}{\sqrt{e^2 - 1}}$ .

## Example 3

Identify the conic for each.

(a) 
$$r = \frac{4}{1 - \sin \theta}$$

## Example 3

Identify the conic for each.

$$(a) \quad r = \frac{4}{1 - \sin \theta}$$

$e = 1 \longrightarrow$  Parabola (opens up).



## Example 3

Identify the conic for each.

$$(a) \quad r = \frac{4}{1 - \sin \theta}$$

$e = 1 \longrightarrow$  Parabola (opens up).

Vertex:  $(0, -2)$

## Example 3

Identify the conic for each.

$$(a) \quad r = \frac{4}{1 - \sin \theta}$$

$e = 1 \longrightarrow$  Parabola (opens up).

Vertex:  $(0, -2)$

Goes through  $(\pm 4, 0)$

## Example 3

$$(b) \quad r = \frac{12}{3 - \cos \theta}$$

## Example 3

$$(b) \quad r = \frac{12}{3 - \cos \theta} \quad \longrightarrow \quad r = \frac{4}{1 - 1/3 \cos \theta}$$

## Example 3

$$(b) \quad r = \frac{12}{3 - \cos \theta} \quad \longrightarrow \quad r = \frac{4}{1 - 1/3 \cos \theta}$$

$$e = \frac{1}{3} \longrightarrow \text{Ellipse (wide)}$$

## Example 3

$$(b) \quad r = \frac{12}{3 - \cos \theta} \quad \longrightarrow \quad r = \frac{4}{1 - 1/3 \cos \theta}$$

$$e = \frac{1}{3} \longrightarrow \text{Ellipse (wide)}$$

$$\frac{1}{3}d = 4 \longrightarrow d = 12$$

## Example 3

$$(b) \quad r = \frac{12}{3 - \cos \theta} \quad \longrightarrow \quad r = \frac{4}{1 - 1/3 \cos \theta}$$

$$e = \frac{1}{3} \longrightarrow \text{Ellipse (wide)}$$

$$\frac{1}{3}d = 4 \longrightarrow d = 12$$

$$\text{Major axis length: } \frac{2(1/3)(12)}{1 - (1/3)^2} = 9$$

## Example 3

$$(b) \quad r = \frac{12}{3 - \cos \theta} \quad \longrightarrow \quad r = \frac{4}{1 - 1/3 \cos \theta}$$

$$e = \frac{1}{3} \longrightarrow \text{Ellipse (wide)}$$

$$\frac{1}{3}d = 4 \longrightarrow d = 12$$

$$\text{Major axis length: } \frac{2(1/3)(12)}{1 - (1/3)^2} = 9$$

$$\text{Minor axis length: } \frac{2(1/3)(12)}{\sqrt{1 - (1/3)^2}} = 6\sqrt{3}$$



## Example 3

$$(c) \quad r = \frac{6}{1 + 2 \sin \theta}$$

## Example 3

$$(c) \quad r = \frac{6}{1 + 2 \sin \theta}$$

$e = 2$ : Hyperbola (opens up and down)

## Example 3

$$(c) \quad r = \frac{6}{1 + 2 \sin \theta}$$

$e = 2$ : Hyperbola (opens up and down)

$$2d = 6 \longrightarrow d = 3$$

## Example 3

$$(c) \quad r = \frac{6}{1 + 2 \sin \theta}$$

$e = 2$ : Hyperbola (opens up and down)

$$2d = 6 \longrightarrow d = 3$$

$$\text{Transverse axis length: } \frac{2(2)(3)}{2^2 - 1} = 4$$

## Example 3

$$(c) \quad r = \frac{6}{1 + 2 \sin \theta}$$

$e = 2$ : Hyperbola (opens up and down)

$$2d = 6 \longrightarrow d = 3$$

$$\text{Transverse axis length: } \frac{2(2)(3)}{2^2 - 1} = 4$$

$$\text{Conjugate axis length: } \frac{2(2)(3)}{\sqrt{2^2 - 1}} = 4\sqrt{3}$$

# Polar Form of Rotated Conics

For constants  $\ell > 0$ ,  $e \geq 0$ , and  $\phi$ , the graph of

$$r = \frac{\ell}{1 - e \cos(\theta - \phi)}$$

is a conic section with eccentricity  $e$  and one focus at  $(0, 0)$ .

# Polar Form of Rotated Conics

If  $e = 0$ , the graph is a circle centered at  $(0, 0)$  with radius  $\ell$ .

# Polar Form of Rotated Conics

If  $e \neq 0$ , the conic has a focus at  $(0, 0)$  and the directrix contains the point with polar coordinates  $(-d, \phi)$  where  $d = \frac{\ell}{e}$ .

- If  $0 < e < 1$ , graph is an ellipse with major axis length  $\frac{2ed}{1 - e^2}$  and minor axis length  $\frac{2ed}{\sqrt{1 - e^2}}$ .



# Polar Form of Rotated Conics

If  $e \neq 0$ , the conic has a focus at  $(0, 0)$  and the directrix contains the point with polar coordinates  $(-d, \phi)$  where  $d = \frac{\ell}{e}$ .

- If  $0 < e < 1$ , graph is an ellipse with major axis length  $\frac{2ed}{1 - e^2}$  and minor axis length  $\frac{2ed}{\sqrt{1 - e^2}}$ .
- If  $e = 1$ , graph is a parabola with focal diameter  $2d$ .

# Polar Form of Rotated Conics

If  $e \neq 0$ , the conic has a focus at  $(0, 0)$  and the directrix contains the point with polar coordinates  $(-d, \phi)$  where  $d = \frac{\ell}{e}$ .

- If  $0 < e < 1$ , graph is an ellipse with major axis length  $\frac{2ed}{1 - e^2}$  and minor axis length  $\frac{2ed}{\sqrt{1 - e^2}}$ .
- If  $e = 1$ , graph is a parabola with focal diameter  $2d$ .
- If  $e > 1$ , graph is a hyperbola with transverse axis length  $\frac{2ed}{e^2 - 1}$  and conjugate axis length  $\frac{2ed}{\sqrt{e^2 - 1}}$ .