

Inverse Trig Functions

Objectives

- 1 Find the values of inverse trig functions
- 2 Find the values of the compositions of inverse trig functions
- 3 Write an algebraic expression for the compositions of inverse trig functions

Trig Functions and the Horizontal Line Test


Recall that for a function to have an **inverse**, any horizontal line can only hit the function at most once.

Trig Functions and the Horizontal Line Test

Recall that for a function to have an **inverse**, any horizontal line can only hit the function at most once.

This creates a problem with periodic functions such as $y = \sin x$.

Graph of $y = \sin x$ and $y = \frac{1}{2}$




Passing the Horizontal Line Test

If we limit the graph to a smaller domain, then the Horizontal Line Test will demonstrate the function $y = \sin x$ will have an inverse:

Passing the Horizontal Line Test

If we limit the graph to a smaller domain, then the Horizontal Line Test will demonstrate the function $y = \sin x$ will have an inverse:



Graph of $y = \sin^{-1} x$


The inverse of $y = \sin x$ is

$$y = \sin^{-1} x \quad \text{or} \quad y = \arcsin x$$

Graph of $y = \sin^{-1} x$

The inverse of $y = \sin x$ is

$$y = \sin^{-1} x \quad \text{or} \quad y = \arcsin x$$



Domains and Ranges for Inverse Trig Functions

Inverse Function	Domain (Input)	Range (Output)
-------------------------	-----------------------	-----------------------

Domains and Ranges for Inverse Trig Functions

Inverse Function	Domain (Input)	Range (Output)
$y = \sin^{-1} x$	$[-1, 1]$	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$

Domains and Ranges for Inverse Trig Functions

Inverse Function	Domain (Input)	Range (Output)
$y = \sin^{-1} x$	$[-1, 1]$	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$
$y = \cos^{-1} x$	$[-1, 1]$	$[0, \pi]$

Domains and Ranges for Inverse Trig Functions

Inverse Function	Domain (Input)	Range (Output)
$y = \sin^{-1} x$	$[-1, 1]$	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$
$y = \cos^{-1} x$	$[-1, 1]$	$[0, \pi]$
$y = \tan^{-1} x$	$(-\infty, \infty)$	$\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$

Domains and Ranges for Inverse Trig Functions

Inverse Function	Domain (Input)	Range (Output)
$y = \sin^{-1} x$	$[-1, 1]$	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$
$y = \cos^{-1} x$	$[-1, 1]$	$[0, \pi]$
$y = \tan^{-1} x$	$(-\infty, \infty)$	$\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$
$y = \csc^{-1} x$	$(-\infty, -1] \cup [1, \infty)$	$\left[-\frac{\pi}{2}, 0\right) \cup \left(0, \frac{\pi}{2}\right]$

Domains and Ranges for Inverse Trig Functions

Inverse Function	Domain (Input)	Range (Output)
$y = \sin^{-1} x$	$[-1, 1]$	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$
$y = \cos^{-1} x$	$[-1, 1]$	$[0, \pi]$
$y = \tan^{-1} x$	$(-\infty, \infty)$	$\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$
$y = \csc^{-1} x$	$(-\infty, -1] \cup [1, \infty)$	$\left[-\frac{\pi}{2}, 0\right) \cup \left(0, \frac{\pi}{2}\right]$
$y = \sec^{-1} x$	$(-\infty, -1] \cup [1, \infty)$	$\left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$

Domains and Ranges for Inverse Trig Functions

Inverse Function	Domain (Input)	Range (Output)
$y = \sin^{-1} x$	$[-1, 1]$	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$
$y = \cos^{-1} x$	$[-1, 1]$	$[0, \pi]$
$y = \tan^{-1} x$	$(-\infty, \infty)$	$\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$
$y = \csc^{-1} x$	$(-\infty, -1] \cup [1, \infty)$	$\left[-\frac{\pi}{2}, 0\right) \cup \left(0, \frac{\pi}{2}\right]$
$y = \sec^{-1} x$	$(-\infty, -1] \cup [1, \infty)$	$\left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$
$y = \cot^{-1} x$	$(-\infty, \infty)$	$(0, \pi)$

Example 1


Find the exact value of each.

(a) $\sin^{-1}\left(\frac{\sqrt{2}}{2}\right)$

Example 1

Find the exact value of each.


(a) $\sin^{-1}\left(\frac{\sqrt{2}}{2}\right)$



Example 1

Find the exact value of each.

(a) $\sin^{-1}\left(\frac{\sqrt{2}}{2}\right)$




$$\theta = \sin^{-1}\left(\frac{\sqrt{2}}{2}\right)$$

Example 1

Find the exact value of each.

(a) $\sin^{-1}\left(\frac{\sqrt{2}}{2}\right)$




$$\theta = \sin^{-1}\left(\frac{\sqrt{2}}{2}\right)$$

$$\theta = 45^\circ$$

Example 1

Find the exact value of each.

(a) $\sin^{-1}\left(\frac{\sqrt{2}}{2}\right)$



$$\theta = \sin^{-1}\left(\frac{\sqrt{2}}{2}\right)$$

$$\theta = 45^\circ$$

$$\theta = \frac{\pi}{4}$$

Example 1


Find the exact value of each.

$$(b) \quad \cos^{-1} \left(-\frac{\sqrt{3}}{2} \right)$$

Example 1

Find the exact value of each.


(b) $\cos^{-1}\left(-\frac{\sqrt{3}}{2}\right)$



Example 1

Find the exact value of each.

(b) $\cos^{-1}\left(-\frac{\sqrt{3}}{2}\right)$




$$\theta = \cos^{-1}\left(\frac{-\sqrt{3}}{2}\right)$$

Example 1

Find the exact value of each.

(b) $\cos^{-1}\left(-\frac{\sqrt{3}}{2}\right)$




$$\theta = \cos^{-1}\left(-\frac{\sqrt{3}}{2}\right)$$

$$\theta = -150^\circ$$

Example 1

Find the exact value of each.

(b) $\cos^{-1}\left(-\frac{\sqrt{3}}{2}\right)$



$$\theta = \cos^{-1}\left(-\frac{\sqrt{3}}{2}\right)$$

$$\theta = -150^\circ$$

$$\theta = -\frac{5\pi}{6}$$

Example 1


Find the exact value of each.

(c) $\arctan(\sqrt{3})$

Example 1

Find the exact value of each.


(c) $\arctan(\sqrt{3})$



Example 1

Find the exact value of each.

(c) $\arctan(\sqrt{3})$




$$\theta = \arctan(\sqrt{3})$$

Example 1

Find the exact value of each.

(c) $\arctan(\sqrt{3})$




$$\theta = \arctan(\sqrt{3})$$

$$\theta = 60^\circ$$

Example 1

Find the exact value of each.

(c) $\arctan(\sqrt{3})$



$$\theta = \arctan(\sqrt{3})$$

$$\theta = 60^\circ$$

$$\theta = \frac{\pi}{3}$$

Example 1


Find the exact value of each.

(d) $\sec^{-1}(2)$

Example 1

Find the exact value of each.


(d) $\sec^{-1}(2)$



Example 1

Find the exact value of each.

(d) $\sec^{-1}(2)$




$$\theta = \sec^{-1}(2)$$

Example 1

Find the exact value of each.

(d) $\sec^{-1}(2)$




$$\theta = \sec^{-1}(2)$$

$$\theta = 60^\circ$$

Example 1

Find the exact value of each.

(d) $\sec^{-1}(2)$



$$\theta = \sec^{-1}(2)$$

$$\theta = 60^\circ$$

$$\theta = \frac{\pi}{3}$$

Objectives

- 1 Find the values of inverse trig functions
- 2 Find the values of the compositions of inverse trig functions
- 3 Write an algebraic expression for the compositions of inverse trig functions

Compositions of Inverse Trig Functions

When dealing with problems like these, it helps to sketch a right triangle, much like in the previous example.

You may need to use the Pythagorean Theorem.

Example 2


Find the exact values of each (if possible).

(a) $\cos\left(\cos^{-1}\left(\frac{1}{3}\right)\right)$

Example 2

Find the exact values of each (if possible).


(a) $\cos\left(\cos^{-1}\left(\frac{1}{3}\right)\right)$



Example 2

Find the exact values of each (if possible).

(a) $\cos\left(\cos^{-1}\left(\frac{1}{3}\right)\right)$




$$\theta = \cos^{-1}\left(\frac{1}{3}\right)$$

Example 2

Find the exact values of each (if possible).

(a) $\cos\left(\cos^{-1}\left(\frac{1}{3}\right)\right)$




$$\theta = \cos^{-1}\left(\frac{1}{3}\right)$$

$$\cos \theta = \frac{x}{r}$$

Example 2

Find the exact values of each (if possible).

(a) $\cos\left(\cos^{-1}\left(\frac{1}{3}\right)\right)$



$$\theta = \cos^{-1}\left(\frac{1}{3}\right)$$

$$\cos \theta = \frac{x}{r}$$


$$\cos \theta = \frac{1}{3}$$

Example 2

$$(b) \quad \sin \left(\arcsin \left(-\frac{3}{7} \right) \right)$$


Example 2

$$(b) \quad \sin \left(\arcsin \left(-\frac{3}{7} \right) \right)$$



Example 2


$$(b) \quad \sin \left(\arcsin \left(-\frac{3}{7} \right) \right)$$



$$\theta = \arcsin \left(-\frac{3}{7} \right)$$

Example 2

$$(b) \quad \sin \left(\arcsin \left(-\frac{3}{7} \right) \right)$$




$$\theta = \arcsin \left(-\frac{3}{7} \right)$$

$$\sin \theta = \frac{y}{r}$$

Example 2

$$(b) \quad \sin \left(\arcsin \left(-\frac{3}{7} \right) \right)$$



$$\theta = \arcsin \left(-\frac{3}{7} \right)$$

$$\sin \theta = \frac{y}{r}$$


$$\sin \theta = -\frac{3}{7}$$

Example 2

$$(c) \quad \cos \left(\arccos \left(\frac{5}{4} \right) \right)$$


Example 2

$$(c) \quad \cos \left(\arccos \left(\frac{5}{4} \right) \right)$$



Example 2


$$(c) \quad \cos \left(\arccos \left(\frac{5}{4} \right) \right)$$



$$\theta = \arccos \left(\frac{5}{4} \right)$$

Example 2

$$(c) \quad \cos \left(\arccos \left(\frac{5}{4} \right) \right)$$




$$\theta = \arccos \left(\frac{5}{4} \right)$$

Domain error.

Example 2

(c) $\cos\left(\arccos\left(\frac{5}{4}\right)\right)$



$$\theta = \arccos\left(\frac{5}{4}\right)$$

Domain error.


\emptyset

Example 2

$$(d) \quad \cos \left(\arctan \left(\frac{5}{12} \right) \right)$$

Example 2


$$(d) \quad \cos\left(\arctan\left(\frac{5}{12}\right)\right)$$



Example 2


$$(d) \quad \cos \left(\arctan \left(\frac{5}{12} \right) \right)$$

$$\theta = \arctan \left(\frac{5}{12} \right)$$



Example 2

$$(d) \quad \cos \left(\arctan \left(\frac{5}{12} \right) \right)$$




$$\theta = \arctan \left(\frac{5}{12} \right)$$

$$12^2 + 5^2 = r^2$$

Example 2

$$(d) \quad \cos \left(\arctan \left(\frac{5}{12} \right) \right)$$




$$\theta = \arctan \left(\frac{5}{12} \right)$$

$$12^2 + 5^2 = r^2$$

$$r = \sqrt{169} = 13$$

Example 2

$$(d) \quad \cos \left(\arctan \left(\frac{5}{12} \right) \right)$$



$$\theta = \arctan \left(\frac{5}{12} \right)$$

$$12^2 + 5^2 = r^2$$

$$r = \sqrt{169} = 13$$


$$\cos \theta = \frac{12}{13}$$

Example 2

$$(e) \quad \sin \left(\arctan \left(\frac{3}{4} \right) \right)$$

Example 2


$$(e) \quad \sin \left(\arctan \left(\frac{3}{4} \right) \right)$$



Example 2


(e) $\sin \left(\arctan \left(\frac{3}{4} \right) \right)$

$$\theta = \arctan \left(\frac{3}{4} \right)$$



Example 2

$$(e) \quad \sin \left(\arctan \left(\frac{3}{4} \right) \right)$$




$$\theta = \arctan \left(\frac{3}{4} \right)$$

$$3^2 + 4^2 = r^2$$

Example 2

$$(e) \quad \sin \left(\arctan \left(\frac{3}{4} \right) \right)$$




$$\theta = \arctan \left(\frac{3}{4} \right)$$

$$3^2 + 4^2 = r^2$$

$$r = \sqrt{25} = 5$$

Example 2

$$(e) \quad \sin \left(\arctan \left(\frac{3}{4} \right) \right)$$



$$\theta = \arctan \left(\frac{3}{4} \right)$$

$$3^2 + 4^2 = r^2$$

$$r = \sqrt{25} = 5$$


$$\sin \theta = \frac{3}{5}$$

Example 2

$$(f) \quad \cot \left(\arcsin \left(-\frac{1}{3} \right) \right)$$


Example 2

$$(f) \quad \cot \left(\arcsin \left(-\frac{1}{3} \right) \right)$$



Example 2


$$(f) \quad \cot \left(\arcsin \left(-\frac{1}{3} \right) \right)$$



$$\theta = \arcsin \left(-\frac{1}{3} \right)$$

Example 2

$$(f) \quad \cot \left(\arcsin \left(-\frac{1}{3} \right) \right)$$




$$\theta = \arcsin \left(-\frac{1}{3} \right)$$

$$x^2 + 1^2 = 3^2$$

Example 2

$$(f) \quad \cot \left(\arcsin \left(-\frac{1}{3} \right) \right)$$




$$\theta = \arcsin \left(-\frac{1}{3} \right)$$

$$x^2 + 1^2 = 3^2$$

$$x = \sqrt{8} = 2\sqrt{2}$$

Example 2

$$(f) \quad \cot \left(\arcsin \left(-\frac{1}{3} \right) \right)$$



$$\theta = \arcsin \left(-\frac{1}{3} \right)$$


$$x^2 + 1^2 = 3^2$$

$$x = \sqrt{8} = 2\sqrt{2}$$

$$\cot \theta = \frac{x}{y}$$

Example 2

$$(f) \quad \cot \left(\arcsin \left(-\frac{1}{3} \right) \right)$$



$$\theta = \arcsin \left(-\frac{1}{3} \right)$$

$$x^2 + 1^2 = 3^2$$

$$x = \sqrt{8} = 2\sqrt{2}$$

$$\cot \theta = \frac{x}{y}$$


$$\cot \theta = \frac{2\sqrt{2}}{-1} = -2\sqrt{2}$$

Example 2

$$(g) \quad \sec \left(\arccos \left(\frac{3}{4} \right) \right)$$


Example 2

$$(g) \quad \sec \left(\arccos \left(\frac{3}{4} \right) \right)$$



Example 2


$$(g) \quad \sec \left(\arccos \left(\frac{3}{4} \right) \right)$$



$$\theta = \arccos \left(\frac{3}{4} \right)$$

Example 2

$$(g) \quad \sec \left(\arccos \left(\frac{3}{4} \right) \right)$$




$$\theta = \arccos \left(\frac{3}{4} \right)$$

$$\sec \theta = \frac{r}{x}$$

Example 2

$$(g) \quad \sec \left(\arccos \left(\frac{3}{4} \right) \right)$$



$$\theta = \arccos \left(\frac{3}{4} \right)$$

$$\sec \theta = \frac{r}{x}$$

$$\sec \theta = \frac{4}{3}$$

Objectives

- 1 Find the values of inverse trig functions
- 2 Find the values of the compositions of inverse trig functions
- 3 Write an algebraic expression for the compositions of inverse trig functions

Algebraic Inverse Trig

In calculus, inverse trigonometric functions will be expressed algebraically (i.e. in terms of x).

Set up a right triangle and use the **Pythagorean Theorem** to write the missing side in terms of x .

Example 3


Write each as an algebraic expression of x .

(a) $\cos(\sin^{-1} x)$

Example 3

Write each as an algebraic expression of x .


(a) $\cos(\sin^{-1} x)$



Example 3

Write each as an algebraic expression of x .

(a) $\cos(\sin^{-1} x)$




$$\theta = \arcsin(x)$$

Example 3

Write each as an algebraic expression of x .

(a) $\cos(\sin^{-1} x)$




$$\theta = \arcsin(x)$$

$$a^2 + x^2 = 1^2$$

Example 3

Write each as an algebraic expression of x .

(a) $\cos(\sin^{-1} x)$



$$\theta = \arcsin(x)$$


$$a^2 + x^2 = 1^2$$

$$a = \sqrt{1 - x^2}$$

Example 3

Write each as an algebraic expression of x .

(a) $\cos(\sin^{-1} x)$



$$\theta = \arcsin(x)$$

$$a^2 + x^2 = 1^2$$


$$a = \sqrt{1 - x^2}$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}}$$

Example 3

Write each as an algebraic expression of x .

(a) $\cos(\sin^{-1} x)$



$$\theta = \arcsin(x)$$

$$a^2 + x^2 = 1^2$$

$$a = \sqrt{1-x^2}$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}}$$


$$\cos \theta = \frac{\sqrt{1-x^2}}{1} = \sqrt{1-x^2}$$

Example 3

(b) $\sec(\tan^{-1} x)$


Example 3

(b) $\sec(\tan^{-1} x)$



Example 3


(b) $\sec(\tan^{-1} x)$



$$\theta = \arctan(x)$$

Example 3

(b) $\sec(\tan^{-1} x)$




$$\theta = \arctan(x)$$

$$1^2 + x^2 = c^2$$

Example 3

(b) $\sec(\tan^{-1} x)$




$$\theta = \arctan(x)$$

$$1^2 + x^2 = c^2$$

$$c = \sqrt{1 + x^2}$$

Example 3

(b) $\sec(\tan^{-1} x)$



$$\theta = \arctan(x)$$


$$1^2 + x^2 = c^2$$

$$c = \sqrt{1 + x^2}$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}}$$

Example 3

(b) $\sec(\tan^{-1} x)$



$$\theta = \arctan(x)$$

$$1^2 + x^2 = c^2$$

$$c = \sqrt{1 + x^2}$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}}$$


$$\sec \theta = \frac{\sqrt{1 + x^2}}{1} = \sqrt{1 + x^2}$$

Example 3

(c) $\tan(\sin^{-1}(2x))$


Example 3

(c) $\tan(\sin^{-1}(2x))$



Example 3


(c) $\tan(\sin^{-1}(2x))$



$$\theta = \arcsin(2x)$$

Example 3

(c) $\tan(\sin^{-1}(2x))$




$$\theta = \arcsin(2x)$$

$$a^2 + (2x)^2 = 1^2$$

Example 3

(c) $\tan(\sin^{-1}(2x))$




$$\theta = \arcsin(2x)$$

$$a^2 + (2x)^2 = 1^2$$

$$a^2 + 4x^2 = 1$$

Example 3

(c) $\tan(\sin^{-1}(2x))$



$$\theta = \arcsin(2x)$$


$$a^2 + (2x)^2 = 1^2$$

$$a^2 + 4x^2 = 1$$

$$a = \sqrt{1 - 4x^2}$$


Example 3

(c) $\tan(\sin^{-1}(2x))$



Example 3


(c) $\tan(\sin^{-1}(2x))$



$$\tan \theta = \frac{\text{opp}}{\text{adj}}$$

Example 3

(c) $\tan(\sin^{-1}(2x))$




$$\tan \theta = \frac{\text{opp}}{\text{adj}}$$

$$\tan \theta = \frac{2x}{\sqrt{1 - 4x^2}}$$

Example 3

(c) $\tan(\sin^{-1}(2x))$



$$\tan \theta = \frac{\text{opp}}{\text{adj}}$$

$$\tan \theta = \frac{2x}{\sqrt{1 - 4x^2}}$$


$$\tan \theta = \frac{2x\sqrt{1 - 4x^2}}{1 - 4x^2}$$

Example 3

$$(d) \quad \sin \left(\cot^{-1} \left(\frac{x}{5} \right) \right)$$


Example 3

(d) $\sin \left(\cot^{-1} \left(\frac{x}{5} \right) \right)$



Example 3


(d) $\sin \left(\cot^{-1} \left(\frac{x}{5} \right) \right)$



$$\theta = \cot^{-1} \left(\frac{x}{5} \right)$$

Example 3

(d) $\sin \left(\cot^{-1} \left(\frac{x}{5} \right) \right)$




$$\theta = \cot^{-1} \left(\frac{x}{5} \right)$$

$$5^2 + x^2 = c^2$$

Example 3

(d) $\sin \left(\cot^{-1} \left(\frac{x}{5} \right) \right)$




$$\theta = \cot^{-1} \left(\frac{x}{5} \right)$$

$$5^2 + x^2 = c^2$$

$$c = \sqrt{25 + x^2}$$


Example 3

(d) $\sin \left(\cot^{-1} \left(\frac{x}{5} \right) \right)$



Example 3


(d) $\sin \left(\cot^{-1} \left(\frac{x}{5} \right) \right)$



$$\sec \theta = \frac{\text{hyp}}{\text{adj}}$$

Example 3

(d) $\sin \left(\cot^{-1} \left(\frac{x}{5} \right) \right)$



$$\sec \theta = \frac{\text{hyp}}{\text{adj}}$$

$$\sec \theta = \frac{\sqrt{25 + x^2}}{x}$$