

# Parametric Equations

# Objectives

- 1 Sketch a parametric curve.
- 2 Rewrite an equation by eliminating the parameter.


Up until now, we have looked at functions that define  $x$  in terms of  $y$  (or  $y$  in terms of  $x$  for inverse functions).

Up until now, we have looked at functions that define  $x$  in terms of  $y$  (or  $y$  in terms of  $x$  for inverse functions).

In this section, we will look at parametric functions: ones in which  $x$  and  $y$  are defined by a **parameter**, such as  $t$ .

# Intro

For instance, the plot below could show the path a bug might take (starting at  $O$ ) while walking on a table:



The independent variable ( $t$  in this case) is called a **parameter**.

# Intro

The independent variable ( $t$  in this case) is called a **parameter**.

The system of equations

$$\begin{cases} x = f(t) \\ y = g(t) \end{cases}$$

is called a **parametrization** of the curve.

The independent variable ( $t$  in this case) is called a **parameter**.

The system of equations

$$\begin{cases} x = f(t) \\ y = g(t) \end{cases}$$

is called a **parametrization** of the curve.

*Note:* The curve itself is a set of points and is devoid of any orientation.



## Example 1

Sketch the curve described by

$$\begin{cases} x = t^2 - 3 \\ y = 2t - 1 \end{cases} \quad \text{for } t \geq -2$$

## Example 1

Sketch the curve described by

$$\begin{cases} x = t^2 - 3 \\ y = 2t - 1 \end{cases} \quad \text{for } t \geq -2$$

$t$	$x(t)$	$y(t)$	$(x(t), y(t))$
-2	1	-5	$(1, -5)$

## Example 1

Sketch the curve described by

$$\begin{cases} x = t^2 - 3 \\ y = 2t - 1 \end{cases} \quad \text{for } t \geq -2$$

$t$	$x(t)$	$y(t)$	$(x(t), y(t))$
-2	1	-5	$(1, -5)$
-1	-2	-3	$(-2, -3)$

## Example 1

Sketch the curve described by

$$\begin{cases} x = t^2 - 3 \\ y = 2t - 1 \end{cases} \quad \text{for } t \geq -2$$

$t$	$x(t)$	$y(t)$	$(x(t), y(t))$
-2	1	-5	(1, -5)
-1	-2	-3	(-2, -3)
0	-3	-1	(-3, -1)

## Example 1

Sketch the curve described by

$$\begin{cases} x = t^2 - 3 \\ y = 2t - 1 \end{cases} \quad \text{for } t \geq -2$$

$t$	$x(t)$	$y(t)$	$(x(t), y(t))$
-2	1	-5	(1, -5)
-1	-2	-3	(-2, -3)
0	-3	-1	(-3, -1)
1	-2	1	(-2, 1)

## Example 1

Sketch the curve described by

$$\begin{cases} x = t^2 - 3 \\ y = 2t - 1 \end{cases} \quad \text{for } t \geq -2$$

$t$	$x(t)$	$y(t)$	$(x(t), y(t))$
-2	1	-5	(1, -5)
-1	-2	-3	(-2, -3)
0	-3	-1	(-3, -1)
1	-2	1	(-2, 1)
2	1	3	(1, 3)

## Example 1

Sketch the curve described by

$$\begin{cases} x = t^2 - 3 \\ y = 2t - 1 \end{cases} \quad \text{for } t \geq -2$$


$t$	$x(t)$	$y(t)$	$(x(t), y(t))$
-2	1	-5	(1, -5)
-1	-2	-3	(-2, -3)
0	-3	-1	(-3, -1)
1	-2	1	(-2, 1)
2	1	3	(1, 3)
3	6	5	(6, 5)

# Example 1

Sketch the curve described by

$$\begin{cases} x = t^2 - 3 \\ y = 2t - 1 \end{cases} \quad \text{for } t \geq -2$$

$t$	$x(t)$	$y(t)$	$(x(t), y(t))$
-2	1	-5	(1, -5)
-1	-2	-3	(-2, -3)
0	-3	-1	(-3, -1)
1	-2	1	(-2, 1)
2	1	3	(1, 3)
3	6	5	(6, 5)





# Objectives

- 1 Sketch a parametric curve.
- 2 Rewrite an equation by eliminating the parameter.

# Eliminating the Parameter

We can eliminate the parameter  $t$  by solving one of the equations for  $t$  and substituting it into the other.

## Example 2

Eliminate the parameter in Example 1 and write the equation using only  $x$  and  $y$ .

$$\begin{cases} x = t^2 - 3 \\ y = 2t - 1 \end{cases} \quad \text{for } t \geq -2$$

## Example 2

Eliminate the parameter in Example 1 and write the equation using only  $x$  and  $y$ .

$$\begin{cases} x = t^2 - 3 \\ y = 2t - 1 \end{cases} \quad \text{for } t \geq -2$$

$$y = 2t - 1$$

## Example 2

Eliminate the parameter in Example 1 and write the equation using only  $x$  and  $y$ .

$$\begin{cases} x = t^2 - 3 \\ y = 2t - 1 \end{cases} \quad \text{for } t \geq -2$$

$$y = 2t - 1$$

$$y + 1 = 2t$$

## Example 2

Eliminate the parameter in Example 1 and write the equation using only  $x$  and  $y$ .

$$\begin{cases} x = t^2 - 3 \\ y = 2t - 1 \end{cases} \quad \text{for } t \geq -2$$

$$y = 2t - 1$$

$$y + 1 = 2t$$

$$t = \frac{y + 1}{2}$$

## Example 2

Eliminate the parameter in Example 1 and write the equation using only  $x$  and  $y$ .

$$\begin{cases} x = t^2 - 3 \\ y = 2t - 1 \end{cases} \quad \text{for } t \geq -2$$

$$y = 2t - 1$$

$$y + 1 = 2t$$

$$t = \frac{y + 1}{2}$$

$$x = t^2 - 3$$

## Example 2

$$x = t^2 - 3$$



## Example 2

$$x = t^2 - 3$$

$$x = \left( \frac{y+1}{2} \right)^2 - 3$$

## Example 2

$$x = t^2 - 3$$

$$x = \left( \frac{y+1}{2} \right)^2 - 3$$

$$x + 3 = \frac{(y+1)^2}{4}$$

## Example 2

$$x = t^2 - 3$$

$$x = \left( \frac{y+1}{2} \right)^2 - 3$$

$$x + 3 = \frac{(y+1)^2}{4}$$

$$4(x+3) = (y+1)^2$$

## Example 2

$$x = t^2 - 3$$

$$x = \left( \frac{y+1}{2} \right)^2 - 3$$

$$x + 3 = \frac{(y+1)^2}{4}$$

$$4(x+3) = (y+1)^2$$

$$t = \frac{y+1}{2}$$

## Example 2

$$x = t^2 - 3$$

$$x = \left( \frac{y+1}{2} \right)^2 - 3$$

$$x + 3 = \frac{(y+1)^2}{4}$$

$$4(x+3) = (y+1)^2$$

$$t = \frac{y+1}{2}$$

$$\frac{y+1}{2} \geq -2$$

## Example 2

$$x = t^2 - 3$$

$$x = \left( \frac{y+1}{2} \right)^2 - 3$$

$$x + 3 = \frac{(y+1)^2}{4}$$

$$4(x+3) = (y+1)^2$$

$$t = \frac{y+1}{2}$$

$$\frac{y+1}{2} \geq -2$$

$$y \geq -5$$

## Example 2

$$4(x + 3) = (y + 1)^2, \quad y \geq -5$$

## Example 3a

Sketch each of the following curves.


$$(a) \quad \begin{cases} x = t^3 \\ y = 2t^2 \end{cases} \quad \text{for } -1 \leq t \leq 1$$



## Example 3a

Sketch each of the following curves.


$$(a) \quad \begin{cases} x = t^3 \\ y = 2t^2 \end{cases} \quad \text{for } -1 \leq t \leq 1$$



## Example 3a

Sketch each of the following curves.

(a) 
$$\begin{cases} x = t^3 \\ y = 2t^2 \end{cases} \quad \text{for } -1 \leq t \leq 1$$




$$x = t^3$$

## Example 3a

Sketch each of the following curves.

$$(a) \quad \begin{cases} x = t^3 \\ y = 2t^2 \end{cases} \quad \text{for } -1 \leq t \leq 1$$




$$x = t^3$$

$$t = \sqrt[3]{x}$$

## Example 3a

Sketch each of the following curves.

$$(a) \quad \begin{cases} x = t^3 \\ y = 2t^2 \end{cases} \quad \text{for } -1 \leq t \leq 1$$



$$x = t^3$$


$$t = \sqrt[3]{x}$$

$$y = 2(\sqrt[3]{x})^2$$

## Example 3a

Sketch each of the following curves.

$$(a) \quad \begin{cases} x = t^3 \\ y = 2t^2 \end{cases} \quad \text{for } -1 \leq t \leq 1$$



$$x = t^3$$

$$t = \sqrt[3]{x}$$


$$y = 2(\sqrt[3]{x})^2$$

$$y = 2 \cdot \sqrt[3]{x^2}$$

## Example 3a

Sketch each of the following curves.

(a)  $\begin{cases} x = t^3 \\ y = 2t^2 \end{cases} \text{ for } -1 \leq t \leq 1$



$$x = t^3$$

$$t = \sqrt[3]{x}$$

$$y = 2(\sqrt[3]{x})^2$$

$$y = 2 \cdot \sqrt[3]{x^2}$$


$$y = 2x^{2/3}$$

## Example 3b

$$(b) \quad \begin{cases} x &= 2e^{-t} \\ y &= e^{-2t} \end{cases} \quad \text{for } t \geq 0$$

## Example 3b

$$(b) \quad \begin{cases} x = 2e^{-t} \\ y = e^{-2t} \end{cases} \quad \text{for } t \geq 0$$






## Example 3b


$$(b) \quad \begin{cases} x = 2e^{-t} \\ y = e^{-2t} \end{cases} \quad \text{for } t \geq 0$$

$$x = 2e^{-t}$$



## Example 3b


$$(b) \quad \begin{cases} x = 2e^{-t} \\ y = e^{-2t} \end{cases} \quad \text{for } t \geq 0$$



$$x = 2e^{-t}$$
$$e^{-t} = \frac{x}{2}$$

## Example 3b

$$(b) \quad \begin{cases} x = 2e^{-t} \\ y = e^{-2t} \end{cases} \quad \text{for } t \geq 0$$




$$x = 2e^{-t}$$

$$e^{-t} = \frac{x}{2}$$

$$y = (e^{-t})^2$$

## Example 3b

$$(b) \quad \begin{cases} x = 2e^{-t} \\ y = e^{-2t} \end{cases} \quad \text{for } t \geq 0$$



$$x = 2e^{-t}$$


$$e^{-t} = \frac{x}{2}$$

$$y = (e^{-t})^2$$

$$y = \left(\frac{x}{2}\right)^2$$

## Example 3b

$$(b) \quad \begin{cases} x = 2e^{-t} \\ y = e^{-2t} \end{cases} \quad \text{for } t \geq 0$$



$$x = 2e^{-t}$$

$$e^{-t} = \frac{x}{2}$$

$$y = (e^{-t})^2$$

$$y = \left(\frac{x}{2}\right)^2$$


$$y = \frac{x^2}{4}$$

## Example 3c

$$(c) \quad \begin{cases} x = \sin t \\ y = \csc t \end{cases} \quad \text{for } 0 < t < \pi$$


## Example 3c

$$(c) \quad \begin{cases} x = \sin t \\ y = \csc t \end{cases} \quad \text{for } 0 < t < \pi$$



## Example 3c

$$(c) \quad \begin{cases} x = \sin t \\ y = \csc t \end{cases} \quad \text{for } 0 < t < \pi$$




$$x = \sin t$$



## Example 3c

$$(c) \quad \begin{cases} x = \sin t \\ y = \csc t \end{cases} \quad \text{for } 0 < t < \pi$$




$$x = \sin t$$

$$y = \csc t$$

## Example 3c

$$(c) \quad \begin{cases} x = \sin t \\ y = \csc t \end{cases} \quad \text{for } 0 < t < \pi$$




$$x = \sin t$$

$$y = \csc t$$

$$y = \frac{1}{\sin t}$$

## Example 3c

$$(c) \quad \begin{cases} x = \sin t \\ y = \csc t \end{cases} \quad \text{for } 0 < t < \pi$$



$$x = \sin t$$

$$y = \csc t$$

$$y = \frac{1}{\sin t}$$


$$y = \frac{1}{x}$$

## Example 3d

$$(d) \quad \begin{cases} x = 1 + 3 \cos t \\ y = 2 \sin t \end{cases} \quad \text{for } 0 \leq t \leq \frac{3\pi}{2}$$


## Example 3d

$$(d) \quad \begin{cases} x = 1 + 3 \cos t \\ y = 2 \sin t \end{cases} \quad \text{for } 0 \leq t \leq \frac{3\pi}{2}$$



## Example 3d


$$(d) \quad \begin{cases} x = 1 + 3 \cos t \\ y = 2 \sin t \end{cases} \quad \text{for } 0 \leq t \leq \frac{3\pi}{2}$$



$$x = 1 + 3 \cos t$$

## Example 3d

$$(d) \quad \begin{cases} x = 1 + 3 \cos t \\ y = 2 \sin t \end{cases} \quad \text{for } 0 \leq t \leq \frac{3\pi}{2}$$




$$x = 1 + 3 \cos t$$

$$\frac{x - 1}{3} = \cos t$$

## Example 3d

$$(d) \quad \begin{cases} x = 1 + 3 \cos t \\ y = 2 \sin t \end{cases} \quad \text{for } 0 \leq t \leq \frac{3\pi}{2}$$



$$x = 1 + 3 \cos t$$


$$\frac{x - 1}{3} = \cos t$$

$$y = 2 \sin t$$



## Example 3d

$$(d) \quad \begin{cases} x = 1 + 3 \cos t \\ y = 2 \sin t \end{cases} \quad \text{for } 0 \leq t \leq \frac{3\pi}{2}$$




$$x = 1 + 3 \cos t$$

$$\frac{x - 1}{3} = \cos t$$

$$y = 2 \sin t$$


$$\frac{y}{2} = \sin t$$

## Example 3d



$$\cos^2 t + \sin^2 t = 1$$


## Example 3d



$$\cos^2 t + \sin^2 t = 1$$

$$\left(\frac{x-1}{3}\right)^2 + \left(\frac{y}{2}\right)^2 = 1$$

## Example 3d



$$\cos^2 t + \sin^2 t = 1$$

$$\left(\frac{x-1}{3}\right)^2 + \left(\frac{y}{2}\right)^2 = 1$$

$$\frac{(x-1)^2}{9} + \frac{y^2}{4} = 1$$

# Parametrizations of Common Curves

- For  $y = f(x)$ , as  $x$  runs through some interval  $I$ , let  $x = t$  and  $y = f(t)$  and let  $t$  run through  $I$ .

# Parametrizations of Common Curves

- For  $y = f(x)$ , as  $x$  runs through some interval  $I$ , let  $x = t$  and  $y = f(t)$  and let  $t$  run through  $I$ .
- For  $x = g(y)$ , as  $y$  runs through some interval  $I$ , let  $x = g(t)$  and  $y = t$  and let  $t$  run through  $I$ .

# Parametrizations of Common Curves

- For  $y = f(x)$ , as  $x$  runs through some interval  $I$ , let  $x = t$  and  $y = f(t)$  and let  $t$  run through  $I$ .
- For  $x = g(y)$ , as  $y$  runs through some interval  $I$ , let  $x = g(t)$  and  $y = t$  and let  $t$  run through  $I$ .
- For a directed line segment with initial point  $(x_0, y_0)$  and terminal point  $(x_1, y_1)$  let  $x = x_0 + (x_1 - x_0)t$  and let  $y = y_0 + (y_1 - y_0)t$  for  $0 \leq t \leq 1$ .

# Parametrizations of Common Curves

- For  $y = f(x)$ , as  $x$  runs through some interval  $I$ , let  $x = t$  and  $y = f(t)$  and let  $t$  run through  $I$ .
- For  $x = g(y)$ , as  $y$  runs through some interval  $I$ , let  $x = g(t)$  and  $y = t$  and let  $t$  run through  $I$ .
- For a directed line segment with initial point  $(x_0, y_0)$  and terminal point  $(x_1, y_1)$  let  $x = x_0 + (x_1 - x_0)t$  and let  $y = y_0 + (y_1 - y_0)t$  for  $0 \leq t \leq 1$ .
- For an ellipse in the form  $\frac{(x-h)^2}{a^2} + \frac{(y-k)^2}{b^2} = 1$ , let  $x = h + a \cos t$  and  $y = k + a \sin t$  for  $0 \leq t < 2\pi$ .



## Example 4

Find a parametrization for each of the following.

(a)  $y = x^2$  from  $x = -3$  to  $x = 2$

## Example 4

Find a parametrization for each of the following.

(a)  $y = x^2$  from  $x = -3$  to  $x = 2$

$$x = t \quad \text{and} \quad y = t^2 \quad \text{for} \quad -3 \leq t \leq 2$$

## Example 4

Find a parametrization for each of the following.

(a)  $y = x^2$  from  $x = -3$  to  $x = 2$

$$x = t \quad \text{and} \quad y = t^2 \quad \text{for} \quad -3 \leq t \leq 2$$

(b)  $y = f^{-1}(x)$  where  $f(x) = x^5 + 2x + 1$

## Example 4

Find a parametrization for each of the following.

(a)  $y = x^2$  from  $x = -3$  to  $x = 2$

$$x = t \quad \text{and} \quad y = t^2 \quad \text{for} \quad -3 \leq t \leq 2$$

(b)  $y = f^{-1}(x)$  where  $f(x) = x^5 + 2x + 1$

$$y = x^5 + 2x + 1$$

## Example 4

Find a parametrization for each of the following.

(a)  $y = x^2$  from  $x = -3$  to  $x = 2$

$$x = t \quad \text{and} \quad y = t^2 \quad \text{for} \quad -3 \leq t \leq 2$$

(b)  $y = f^{-1}(x)$  where  $f(x) = x^5 + 2x + 1$

$$y = x^5 + 2x + 1$$

$$x = y^5 + 2y + 1$$

## Example 4

Find a parametrization for each of the following.

(a)  $y = x^2$  from  $x = -3$  to  $x = 2$

$$x = t \quad \text{and} \quad y = t^2 \quad \text{for} \quad -3 \leq t \leq 2$$

(b)  $y = f^{-1}(x)$  where  $f(x) = x^5 + 2x + 1$

$$y = x^5 + 2x + 1$$

$$x = y^5 + 2y + 1$$

$$y = t \quad x = t^5 + 2t + 1 \quad \text{for} \quad -\infty < t < \infty$$

## Example 4c

- (c) The line segment which starts at  $(2, -3)$  and ends at  $(1, 5)$

## Example 4c

- (c) The line segment which starts at  $(2, -3)$  and ends at  $(1, 5)$

$$x_1 - x_0 = 1 - 2$$



## Example 4c

(c) The line segment which starts at  $(2, -3)$  and ends at  $(1, 5)$

$$\begin{aligned}x_1 - x_0 &= 1 - 2 \\ &= -1\end{aligned}$$

## Example 4c

- (c) The line segment which starts at  $(2, -3)$  and ends at  $(1, 5)$

$$x_1 - x_0 = 1 - 2$$

$$= -1$$

$$x = x_0 + (x_1 - x_0)t$$

## Example 4c

(c) The line segment which starts at  $(2, -3)$  and ends at  $(1, 5)$

$$x_1 - x_0 = 1 - 2$$

$$= -1$$

$$x = x_0 + (x_1 - x_0)t$$

$$x = 2 + (-1)t$$

## Example 4c

(c) The line segment which starts at  $(2, -3)$  and ends at  $(1, 5)$

$$x_1 - x_0 = 1 - 2$$

$$= -1$$

$$x = x_0 + (x_1 - x_0)t$$

$$x = 2 + (-1)t$$

$$x = 2 - t$$

$$y_1 - y_0 = 5 - (-3)$$

## Example 4c

(c) The line segment which starts at  $(2, -3)$  and ends at  $(1, 5)$

$$x_1 - x_0 = 1 - 2$$

$$= -1$$

$$x = x_0 + (x_1 - x_0)t$$

$$x = 2 + (-1)t$$

$$x = 2 - t$$

$$y_1 - y_0 = 5 - (-3)$$

$$= 8$$

## Example 4c

(c) The line segment which starts at  $(2, -3)$  and ends at  $(1, 5)$

$$x_1 - x_0 = 1 - 2$$

$$= -1$$

$$x = x_0 + (x_1 - x_0)t$$

$$x = 2 + (-1)t$$

$$x = 2 - t$$

$$y_1 - y_0 = 5 - (-3)$$

$$= 8$$

$$y = y_0 + (y_1 - y_0)t$$

## Example 4c

(c) The line segment which starts at  $(2, -3)$  and ends at  $(1, 5)$

$$x_1 - x_0 = 1 - 2$$

$$= -1$$

$$x = x_0 + (x_1 - x_0)t$$

$$x = 2 + (-1)t$$

$$x = 2 - t$$

$$y_1 - y_0 = 5 - (-3)$$

$$= 8$$

$$y = y_0 + (y_1 - y_0)t$$

$$y = -3 + 8t$$

## Example 4c

(c) The line segment which starts at  $(2, -3)$  and ends at  $(1, 5)$

$$x_1 - x_0 = 1 - 2$$

$$= -1$$

$$x = x_0 + (x_1 - x_0)t$$

$$x = 2 + (-1)t$$

$$x = 2 - t$$

$$y_1 - y_0 = 5 - (-3)$$

$$= 8$$

$$y = y_0 + (y_1 - y_0)t$$

$$y = -3 + 8t$$

$$\text{for } 0 \leq t \leq 1$$



## Example 4

(d) The circle  $x^2 + 2x + y^2 - 4y = 4$

## Example 4

(d) The circle  $x^2 + 2x + y^2 - 4y = 4$

$$\frac{(x+1)^2}{9} + \frac{(y-2)^2}{9} = 1$$

## Example 4

(d) The circle  $x^2 + 2x + y^2 - 4y = 4$

$$\frac{(x+1)^2}{9} + \frac{(y-2)^2}{9} = 1$$

$$x = -1 + 3 \cos t \quad y = 2 + 3 \sin t \quad \text{for } 0 \leq t < 2\pi$$

## Example 4

(e) The left half of the ellipse  $\frac{x^2}{4} + \frac{y^2}{9} = 1$

## Example 4

(e) The left half of the ellipse  $\frac{x^2}{4} + \frac{y^2}{9} = 1$

$$x = 2 \cos t$$

## Example 4

(e) The left half of the ellipse  $\frac{x^2}{4} + \frac{y^2}{9} = 1$

$$x = 2 \cos t$$

$$y = 3 \sin t$$

## Example 4

(e) The left half of the ellipse  $\frac{x^2}{4} + \frac{y^2}{9} = 1$

$$x = 2 \cos t$$

$$y = 3 \sin t$$

$$\text{for } \frac{\pi}{2} \leq t \leq \frac{3\pi}{2}$$