

Math CS 121 HW 5

Zih-Yu Hsieh

November 3, 2025

1

Question 1. 5.2.27:

Let X be a Poisson random variable with parameter λ . Show that the maximum of $\mathbb{P}(X = i)$ occurs at $[\lambda]$, where $[\lambda]$ is the greatest integer less than or equal to λ .

Hint: Let p be the probability mass function of X . Prove that

$$p(i) = \frac{\lambda}{i} p(i-1)$$

Use this to find the values of i at which p is increasing and the values of i at which it is decreasing.

Proof. Recall that for random variable $X \sim \text{Poi}(\lambda)$ (where $\lambda > 0$), then for each $i \in \mathbb{N}$, one has $p(i) = e^{-\lambda} \cdot \frac{\lambda^i}{i!}$. So, for any index $i > 0$, the following relation is true:

$$p(i) = e^{-\lambda} \cdot \frac{\lambda^i}{i!} = e^{-\lambda} \cdot \frac{\lambda^{i-1}}{(i-1)!} \cdot \frac{\lambda}{i} = p(i-1) \cdot \frac{\lambda}{i} \quad (1)$$

So, for all positive integer $i \leq \lambda$, since $\frac{\lambda}{i} \geq 1$, it satisfies $p(i) = p(i-1) \cdot \frac{\lambda}{i} \geq p(i-1)$; conversely, for $i > \lambda$, since $\frac{\lambda}{i} < 1$, it satisfies $p(i) = p(i-1) \cdot \frac{\lambda}{i} < p(i-1)$.

Now, consider integer $j = [\lambda]$ (the largest integer satisfying $k \leq \lambda$). To derive the maximality of $p(j)$, there are two cases:

- Since for any integer $i \leq j$, $1 \leq j \leq \lambda$, so $p(i) \geq p(i-1)$, hence inductively one can prove that $p(i) \leq p(j)$.
- Similarly, for any integer $i > j$, since j is the largest integer satisfying $k \leq \lambda$, then $i > j$ implies that $i > \lambda$, hence we have $p(i) < p(i-1)$ instead. So again, inductively one can also prove that $p(i) < p(j)$.

Since for all $i \in \mathbb{N}$ one must have $p(i) \leq p(j)$ (from the two cases above), then one must have $p(j) = p([\lambda])$ being the maximum value of the probability mass function. Hence, p has maximum occurring at $[\lambda]$. $\square$

2 ND

Question 2. 5.3.24:

Twelve hundred eggs, of which 200 are rotten, are distributed randomly in 100 cartons, each containing a dozen eggs. These cartons are then sold to a restaurant. How many cartons should we expect the chef of the restaurant to open before finding one without rotten eggs?

Proof.

□

Question 3. 5.Rev.14:

Past experience shows that 30% of the customers entering Harry's Clothing Store will make a purchase. Of the customers who make a purchase, 85% use credit cards. Let X be the number of the next six customers who enter the store, make a purchase, and use a credit card. Find the probability mass function, the expected value, and the variance of X .

Proof. First, it's only possible to have $X \in \{0, 1, 2, 3, 4, 5, 6\}$ (since it represents the number of customers "within" the next 6 customers, that make a purchase and using a credit card).

Let Y denotes the random variable of the number of the next six customers who enter the store and make a purchase (similarly, $Y \in \{0, 1, 2, 3, 4, 5, 6\}$). Since there is a probability of $p = 0.3$ that the entered customer will make a purchase, and each of the customer's probability of making a purchase should be independent of each other, hence $Y \sim \text{Bin}(6, p = 0.3)$. Which, for any $i \in \{0, 1, 2, 3, 4, 5, 6\}$, $\mathbb{P}(Y = i) = \binom{6}{i} p^i \cdot (1-p)^{6-i} = \frac{1}{10^6} \binom{6}{i} 3^i \cdot 7^{6-i}$. Also, given $i \neq j$, then the event $Y = i$ and $Y = j$ are disjoint.

Now, if we fix the number of customers who purchased, say $Y = j$, then notice that for each customer in these j customers who purchased, each independently has probability $q = 0.85$ of using credit card to pay. Hence, the random variable X_j (denoting random variable X given that $Y = j$) has $X_j \sim \text{Bin}(j, q = 0.85)$, hence for any $i \leq j$, we have $\mathbb{P}(X_j = i) = \mathbb{P}(X = i | Y = j) = \binom{j}{i} q^i (1-q)^{j-i} = \frac{1}{100^j} \binom{j}{i} \cdot 85^i \cdot 15^{j-i}$. Also, for $i > j$, we have $\mathbb{P}(X_j = i) = \mathbb{P}(X = i | Y = j) = 0$, since the number of people who purchased and used credit card ($X = i$) must be at most the same as the number of people who made a purchase $Y = j$, so $i > j$ is impossible.

With these information, given any number $i \in \{0, 1, 2, 3, 4, 5, 6\}$, we have $\mathbb{P}(X = i)$ given as follow:

$$\mathbb{P}(X = i) = \sum_{j=0}^6 \mathbb{P}(X = i | Y = j) \cdot \mathbb{P}(Y = j) = \sum_{j=0}^6 \mathbb{P}(X_j = i) \cdot \mathbb{P}(Y = j) = \sum_{j=i}^6 \mathbb{P}(X_j = i) \cdot \mathbb{P}(Y = j) \quad (2)$$

(Note: the last part of the equality is due to the fact that $i > j$ implies $\mathbb{P}(X_j = i) = 0$). Which, the term further refines to:

$$\mathbb{P}(X = i) = \sum_{j=i}^6 \mathbb{P}(X_j = i) \cdot \mathbb{P}(Y = j) = \sum_{j=i}^6 \left(\frac{1}{100^j} \binom{j}{i} \cdot 85^i \cdot 15^{j-i} \right) \cdot \left(\frac{1}{10^6} \binom{6}{j} 3^j \cdot 7^{6-j} \right) \quad (3)$$

calculate these terms numerically, we get the following as the Probability Mass Function:

$$\mathbb{P}(X = 0) \approx 0.1710, \quad \mathbb{P}(X = 1) \approx 0.3511, \quad \mathbb{P}(X = 2) \approx 0.3005, \quad \mathbb{P}(X = 3) \approx 0.1371 \quad (4)$$

$$\mathbb{P}(X = 4) \approx 0.0352, \quad \mathbb{P}(X = 5) \approx 0.0048, \quad \mathbb{P}(X = 6) \approx 0.0003 \quad (5)$$

With the probability mass function, we can calculate the expected value $\mathbb{E}[X]$ and variance $\text{Var}(X)$ (which we'll use $\text{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2$, and calculate $\mathbb{E}[X^2]$ the second moment beforehand).

The First Moment/expected value $\mathbb{E}[X]$, and Second Moment $\mathbb{E}[X^2]$ are given as follow numerically:

$$\mathbb{E}[X] = \sum_{i=0}^6 i \cdot \mathbb{P}(X = i) = 1.53, \quad \mathbb{E}[X^2] = \sum_{i=0}^6 i^2 \cdot \mathbb{P}(X = i) = 3.48075 \quad (6)$$

Hence, we have expected value $\mathbb{E}[X] = 1.53$, while variance $\text{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2 = 1.13985$. $\square$

4

Question 4. 5.Rev.24:

Experience shows that 75% of certain kinds of seeds germinate when planted under normal conditions. Determine the minimum number of seeds to be planted so that the chance of at least five of them germinating is more than 90%.

Proof. First, fix any $n \in \mathbb{N}$ that represents the number of seeds planted. Let X_n be the random variable, representing the number of seeds germinating out of the n planted seeds. Then, with each seed having a $p = 0.75$ probability of germinating (and the germinating probability is supposedly independent for each seed), then $X_n \sim \text{Bin}(n, p = 0.75)$ (since each success probability is independent, and it's either being germinated or not).

Then, to find the minimum n such that the probability of at least five of them germinating being more than 90%, the first requirement is $n \geq 5$ (since less than 5 seeds being planted, it's impossible to have at least 5 germinating). Which, notice that this is given by $\mathbb{P}(X_n \geq 5)$, which the complement is given by $\mathbb{P}(X_n < 5) = \mathbb{P}(X_n \leq 4)$. Which, the requirement that $\mathbb{P}(X_n \geq 5) > 0.9$ (more than 90% germinating), is equivalent to $\mathbb{P}(X_n \leq 4) = 1 - \mathbb{P}(X_n \geq 5) < 0.1$.

Now, for any $n \geq 5$, the probability $\mathbb{P}(X_n \leq 4)$ is given by:

$$\mathbb{P}(X_n \leq 4) = \sum_{k=0}^4 \mathbb{P}(X_n = k) = \sum_{k=0}^4 \binom{n}{k} \cdot p^k \cdot (1-p)^{n-k} = \sum_{k=0}^4 \binom{n}{k} \cdot \left(\frac{3}{4}\right)^k \cdot \left(\frac{1}{4}\right)^{n-k} = \frac{1}{4^n} \sum_{k=0}^4 \binom{n}{k} \cdot 3^k \quad (7)$$

Which, numerically we get that for $n = 8$, $\mathbb{P}(X_8 \leq 4) \approx 0.1138 > 0.1$, while for $n = 9$, $\mathbb{P}(X_9 \leq 4) \approx 0.0489 < 0.1$. Hence, $n = 9$ is the smallest number of seeds to be planted to have $\mathbb{P}(X_n \leq 4) < 0.1$, which is also the smallest number of seeds to be planted so that $\mathbb{P}(X_n \geq 5) = 1 - \mathbb{P}(X_n \leq 4) > 0.9$. $\square$

5

Question 5. *An urn contains N green balls and M red ones. Balls are taken out 1-by-1 and the procedure is stopped when a red ball first appears. Let X be the number of green balls taken out. Determine analytically the expected value of X if:*

- a) *Removed balls are returned back (sampling with replacement)*
- b) *Removed balls are gone (sampling without replacement)*

Provide the actual numerical solution when $N = 7$, $M = 5$.

Hint: One of the answers above is going to be ugly, that's just how it is...

Proof.

- a) When the balls are returned each time, since there are total of $N + M$ balls, with N being green balls, then the probability of getting green balls in each take is $p = \frac{N}{N+M}$, while getting red balls in each take is $q = \frac{M}{N+M} = 1 - p$.

Given any $i \in \mathbb{N}$, for the event $X = i$ to happen, we need to get i green balls consecutively, while on the $(i+1)^{\text{th}}$ draw, we need to get red ball to end the process. Hence, $\mathbb{P}(X = i) = p^i \cdot q^1 = \left(\frac{N}{N+M}\right)^i \cdot \frac{M}{N+M}$. So, the expected value of X , $\mathbb{E}[X]$ is given as follow:

$$\mathbb{E}[X] = \sum_{i=0}^{\infty} i \cdot \mathbb{P}(X = i) = \sum_{i=0}^{\infty} \left(\frac{N}{N+M}\right)^i \cdot \frac{M}{N+M} = \frac{M}{N+M} \cdot \frac{1}{1 - \frac{N}{N+M}} = \frac{M}{N+M} \cdot \frac{N+M}{M} = 1 \quad (8)$$

So, the expected number of green balls being drawn is $\mathbb{E}[X] = 1$.

- b) If the removed balls are not putting back to the urn, then people can draw at most N green balls out of the urn, or $X \in \{0, 1, \dots, N\}$. For any $i \in \{0, 1, \dots, n\}$, in case for the event $X = i$ to happen, the person needs to consecutively draw i green balls out of the urn, and on the $(i+1)^{\text{th}}$ move one needs to draw a red ball.

Here, we'll consider the probability of drawing $0 \leq i \leq N$ consecutive green balls (without considering how to end with red balls) using induction.

For $i = 1$, the probability of drawing out $i = 1$ green ball is given by $\frac{N}{N+M}$. For $i = 2$, the probability of drawing out 2 green balls consecutively is given by $\frac{N}{N+M} \cdot \frac{N-1}{N+M-1}$ (since the second green ball is picked from $N - 1$ remaining green balls, while there are $N + M - 1$ balls in total remaining).

Inductively, suppose for given integer $0 \leq k < i$, the probability of drawing out k green balls consecutively is given by $\frac{N!}{(N-k)!} \cdot \frac{(N+M-k)!}{(N+M)!}$, then for the case of $k = i$, since the probability of drawing the first $(i-1)$ green balls is given by $\frac{N!}{(N-(i-1))!} \cdot \frac{(N+M-(i-1))!}{(N+M)!}$, while there are $N - (i-1)$ green balls, and $N + M - (i-1)$ total balls remaining, hence drawing the i^{th} green ball now has a probability of $\frac{N-(i-1)}{N+M-(i-1)}$. Hence, the probability of drawing $k = i$ consecutive green balls, is given by $\left(\frac{N!}{(N-(i-1))!} \cdot \frac{(N+M-(i-1))!}{(N+M)!}\right) \cdot \frac{N-(i-1)}{N+M-(i-1)} = \frac{N!}{(N-i)!} \cdot \frac{(N+M-i)!}{(N+M)!}$.

Finally, if given integer $0 \leq i \leq N$, we know the event $X = i$ happens if i consecutive green balls are drawn (which has probability $\frac{N!}{(N-i)!} \cdot \frac{(N+M-i)!}{(N+M)!}$), while the $(i+1)^{\text{th}}$ ball drawn is red. Since after the first i picks, there are M red balls leftover, while only $N + M - i$ total balls remaining, the probability of drawing a red ball on the $(i+1)^{\text{th}}$ term is $\frac{M}{N+M-i}$. Hence, probability of $X = i$ is given by:

$$\mathbb{P}(X = i) = \frac{N!}{(N-i)!} \cdot \frac{(N+M-i)!}{(N+M)!} \cdot \frac{M}{N+M-i} \quad (9)$$

As a result, the corresponding expected value for X is given as follow:

$$\mathbb{E}[X] = \sum_{i=0}^N i \cdot \mathbb{P}(X = i) = \sum_{i=0}^N i \cdot \frac{N!}{(N-i)!} \cdot \frac{(N+M-i)!}{(N+M)!} \cdot \frac{M}{N+M-i} \quad (10)$$

After all the above calculation, here are the numerical result for $N = 7$, $M = 5$:

- a) If it's a sample with replacement, then $\mathbb{E}[X] = 1$ without ambiguity.
b) If it's a sample without replacement, then $\mathbb{E}[X]$ using the formula above, is given as follow:

$$\mathbb{E}[X]_{N=7, M=5} = \sum_{i=0}^7 i \cdot \frac{7!}{(7-i)!} \cdot \frac{(12-i)!}{12!} \cdot \frac{5}{12-i} \approx 1.1667 \quad (11)$$

□

Question 6. A student answers a multiple-choice test completely randomly. What is the probability he will pass (get 50% or more correct) if:

- a) The test has 10 questions with 4 choices for each question.
- b) Find the smallest number of questions with 4 choices for each such that the probability of passing through random guessing is less than 1%.
- c) The test has 10 questions with 4 choices each. There are 10 students in the class. Assuming each student guesses all the answers, what is the probability that all 10 students do not pass.

Proof. If the student guesses the multiple-choice test completely randomly, then each of the question's probability of getting right / wrong is independent from the other questions. Also, since each question is multiple choice (suppose there's only one right answer among the choices), if there are N total choices, then the probability of getting the question right is $\frac{1}{N}$.

- a) For total of 10 questions, with each question having 4 choices. Then, since getting a question itself right has probability $p = \frac{1}{4} = 0.25$, and there are total of 10 questions, let X being the random variable of the number of question this student got right out of 10 questions, then $X \sim \text{Bin}(10, p = 0.25)$ (total of 10 questions, each has an independent probability of 0.25 of getting right).

In case to pass, it needs the variable $X \geq 5$ (since we need at least $10 \cdot 0.5 = 5$ questions, or at least 50% of the questions right to pass). So, we want to find $\mathbb{P}(X \geq 5)$, it is given by:

$$\mathbb{P}(X \leq 5) = \sum_{k=5}^{10} \mathbb{P}(X = k) = \sum_{k=5}^{10} \binom{10}{k} \cdot p^k (1-p)^{10-k} \quad (12)$$

$$= \sum_{k=5}^{10} \binom{10}{k} \cdot \frac{1}{4^k} \cdot \left(\frac{3}{4}\right)^{10-k} = \frac{1}{4^{10}} \sum_{k=5}^{10} \binom{10}{k} \cdot 3^{10-k} \quad (13)$$

Which, $\mathbb{P}(X \geq 5) \approx 0.0781$, hence the probability of passing by random guessing is about 0.0781.

- b) Here, if the total number of questions is labeled as $n \in \mathbb{N}$, then let X_n denotes the random variable of the number of questions this student got right out of n questions, we have $X_n \sim \text{Bin}(n, p = 0.25)$ (since there are still 4 choices per question, so the probability p of answering each question right stays the same).

Which, in case to pass the test, it requires more than 50% of the questions to be right, therefore the student must have $X_n \geq \lceil \frac{n}{2} \rceil$ (where $\lceil \frac{n}{2} \rceil$ denotes the smallest number that's greater than or equal to $\frac{n}{2}$). So, the probability of passing the test is given by $\mathbb{P}(X_n \geq \lceil \frac{n}{2} \rceil)$, which has its complement given by $\mathbb{P}(X_n < \lceil \frac{n}{2} \rceil) = \mathbb{P}(X_n \leq \lceil \frac{n}{2} \rceil - 1)$. Again, using the probability of binomial distribution, it is given by:

$$\mathbb{P}(X_n \geq \lceil \frac{n}{2} \rceil) = \sum_{k=\lceil \frac{n}{2} \rceil}^n \mathbb{P}(X_n = k) = \sum_{k=\lceil \frac{n}{2} \rceil}^n \binom{n}{k} p^k (1-p)^{n-k} \quad (14)$$

$$= \sum_{k=\lceil \frac{n}{2} \rceil}^n \binom{n}{k} \frac{1}{4^k} \left(\frac{3}{4}\right)^{n-k} = \frac{1}{4^n} \sum_{k=\lceil \frac{n}{2} \rceil}^n \binom{n}{k} 3^{n-k} \quad (15)$$

Which, as a numerical result, for $n = 18$, $\mathbb{P}(X_{18} \geq 9) \approx 0.0193 > 0.01$, while for $n = 19$, $\mathbb{P}(X_{19} \geq 10) \approx 0.0089 < 0.01$. So, $n = 19$ is the smallest number of questions, so that with 4 choices for each question, the probability of passing by random guessing is less than 0.01 (or 1%).

- c) Here, can suppose all 10 students' score are independent of each other (since their guess theoretically shouldn't be affecting each other). Hence, the probability of all 10 students not passing, is the same as product of probability that each student doesn't pass.

Using information in part a), we know that given 10 questions and each with 4 choices, the probability that a student passes by randomly guessing is $p \approx 0.0781$, so the probability of not passing when just randomly guessing is $q = 1 - p \approx 0.9219$. Hence, the probability that all 10 students don't pass is given by $q^{10} \approx 0.4434$.

□

7

Question 7. 6.3.9:

A right triangle has a hypotenuse of length 9. If the probability density function of one side's length is given by:

$$f(x) = \begin{cases} x/6 & \text{if } 2 < x < 4 \\ 0 & \text{otherwise} \end{cases}$$

what is the expected value of the length of the other side?

Proof. (**Rmk:** Since side length is only valid for positive number, we'll assume $x > 0$ to be the case).

Given it's a right triangle with hypotenuse of length 9. If one side length is given by variable $x \in (0, \infty)$, then the other side is given by $y = \sqrt{81 - x^2}$. Which, y is only well-defined if $81 - x^2 \geq 0$, or $x \in [-9, 9]$. Let Y denotes the random variable calculating the length of the other side y .

Then, based on the probability density function, one can conclude the following expected value $\mathbb{E}[Y]$ (which, for y, x to both be valid, one must have $0 < x \leq 9$):

$$\mathbb{E}[Y] = \int_0^9 y f(x) dx = \int_0^2 y \cdot 0 dx + \int_2^4 y \cdot \frac{x}{6} dx + \int_4^9 y \cdot 0 dx = \int_2^4 \sqrt{81 - x^2} \cdot \frac{x}{6} dx \quad (16)$$

Performing substitution $u = 81 - x^2$, then $du = -2x dx$; at $x = 2$, $u = 81 - 4 = 77$, and at $x = 4$, $u = 81 - 16 = 65$. So, the above integral becomes:

$$\mathbb{E}[Y] = \int_2^4 \sqrt{81 - x^2} \cdot \frac{x}{6} dx = - \int_{77}^{65} \sqrt{u} \cdot \frac{1}{12} \cdot (-2x) dx = - \int_{77}^{65} \sqrt{u} \cdot \frac{1}{12} du \quad (17)$$

$$= \frac{1}{12} \int_{65}^{77} \sqrt{u} du = \frac{1}{12} \cdot \frac{2}{3} u^{\frac{3}{2}} \Big|_{65}^{77} = \frac{1}{18} \left((77)^{\frac{3}{2}} - (65)^{\frac{3}{2}} \right) \approx 8.4236 \quad (18)$$

So, the expected value of the other side is $y \approx 8.4236$.

□

Question 8. 6.3.10:

Let X be a random variable with probability density function:

$$f(x) = \frac{1}{2}e^{-|x|}, \quad x \in \mathbb{R}$$

Calculate $\text{Var}(X)$.

Proof. We'll first rewrite the function $f(x)$ as follow:

$$f(x) = \begin{cases} \frac{1}{2}e^{-x} & x \geq 0 \\ \frac{1}{2}e^x & x < 0 \end{cases} \quad (19)$$

Hence, with $\text{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2$, it suffices to calculate the moments. The two moments involved are given as follow (Note: All the below integral converges since it's polynomial growth · exponential decay when $x \rightarrow \pm\infty$, so it's valid to perform a substitution $u = -x$, $du = -dx$ for one of the parts):

$$\mathbb{E}[X] = \int_{-\infty}^{\infty} xf(x)dx = \int_{-\infty}^0 x \cdot \frac{1}{2}e^x dx + \int_0^{\infty} x \cdot \frac{1}{2}e^{-x} dx \quad (20)$$

$$= \int_{-\infty}^0 \frac{1}{2}xe^x dx + \int_0^{-\infty} \frac{1}{2}(-u)e^u \cdot (-1)du \quad (21)$$

$$= \int_{-\infty}^0 \frac{1}{2}xe^x dx + \int_0^{-\infty} \frac{1}{2}ue^u du \quad (22)$$

$$= \int_{-\infty}^0 \frac{1}{2}xe^x dx - \int_{-\infty}^0 \frac{1}{2}xe^x dx = 0 \quad (23)$$

$$\mathbb{E}[X^2] = \int_{-\infty}^{\infty} x^2 f(x)dx = \int_{-\infty}^0 x^2 \cdot \frac{1}{2}e^x dx + \int_0^{\infty} x^2 \cdot \frac{1}{2}e^{-x} dx \quad (24)$$

$$= \int_{-\infty}^0 x^2 \cdot \frac{1}{2}e^x dx + \int_0^{-\infty} u^2 \cdot \frac{1}{2}e^u \cdot (-1)du \quad (25)$$

$$= \int_{-\infty}^0 x^2 \cdot \frac{1}{2}e^x dx + \int_{-\infty}^0 u^2 \cdot \frac{1}{2}e^u du \quad (26)$$

$$= \int_{-\infty}^0 x^2 e^x dx = x^2 e^x - 2xe^x + 2e^x \Big|_{-\infty}^0 = 2 \quad (27)$$

Hence, the variance $\text{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2 = 2 - 0^2 = 2$. □

Question 9. 6.3.16:

Let X be a continuous random variable with probability density function $f(x)$. Determine the value of y for which $\mathbb{E}[|X - y|]$ is minimum.

Proof. Let $y \in \mathbb{R}$ be arbitrary. Then, the function $|x - y|$ is expressed as follow:

$$|x - y| = \begin{cases} x - y & x \geq y \\ y - x & x < y \end{cases} \quad (28)$$

Hence, when calculating $\mathbb{E}[|X - y|]$, it is given as follow:

$$G(y) := \mathbb{E}[|X - y|] = \int_{-\infty}^{\infty} |x - y|f(x)dx = \int_{-\infty}^y (y - x)f(x)dx + \int_y^{\infty} (x - y)f(x)dx \quad (29)$$

$$= y \int_{-\infty}^y f(x)dx - \int_{-\infty}^y xf(x)dx + \int_y^{\infty} xf(x)dx - y \int_y^{\infty} f(x)dx \quad (30)$$

Notice that given $f(x)$ is continuous, the above integrals in terms of y are also continuous, hence $G(y)$ is a continuous function.

Then, the minimum of $G(y) = \mathbb{E}[|X - y|]$ must occure at a point where $G'(y) = 0$. Then, taking differentiation (using Fundamental Theorem of Calculus, Product Rule, etc.), we get the following for $G'(y)$:

$$G'(y) = \int_{-\infty}^y f(x)dx + yf(y) - yf(y) - yf(y) - \int_y^{\infty} f(x)dx - y(-f(y)) \quad (31)$$

$$= \int_{-\infty}^y f(x)dx - \int_y^{\infty} f(x)dx = \mathbb{P}(X \leq y) - \mathbb{P}(X > y) \quad (32)$$

Hence, if $G'(y) = 0$, one must have $\mathbb{P}(X \leq y) - \mathbb{P}(X > y) = 0$, or $\mathbb{P}(X \leq y) = \mathbb{P}(X > y)$. However, recall that $1 = \mathbb{P}(X \in \mathbb{R}) = \mathbb{P}(X \leq y \sqcup X > y) = \mathbb{P}(X \leq y) + \mathbb{P}(X > y)$. With the two being equal, one must have $\mathbb{P}(X \leq y) = \mathbb{P}(X > y) = \frac{1}{2}$. Hence, the only possible value where $\mathbb{E}[|X - y|]$ is minimum, is when $\mathbb{P}(X \leq y) = \frac{1}{2}$.

Finally, to check that it's actually a minimum, we'll consider the second derivative of G :

$$G''(y) = \frac{d}{dy} \left(\int_{-\infty}^y f(x)dx - \int_y^{\infty} f(x)dx \right) = f(y) - (-f(y)) = 2f(y) \geq 0 \quad (33)$$

Which, for $y \in \mathbb{R}$ to be a minimum of $G(y)$, one also needs $G''(y) > 0$ to conclude it. Hence, if $y \in \mathbb{R}$ satisfies $\mathbb{P}(X \leq y) = \frac{1}{2}$ and $f(y) > 0$, we can conclude that $\mathbb{E}[|X - y|]$ is at its minimum. $\square$

Question 10. 6.Rev.3:

Let X be a continuous random variable with density function:

$$f(x) = 6x(1 - x), \quad 0 < x < 1$$

What is the probability that X is within two standard deviations of the mean?

Proof. First, recall that $\text{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2$. Which, these two are given as follow respectively:

$$\mathbb{E}[X] = \int_0^1 x f(x) dx = \int_0^1 (6x^2 - 6x^3) dx = 2x^3 - \frac{3}{2}x^4 \Big|_0^1 = 2 - \frac{3}{2} = \frac{1}{2} \quad (34)$$

$$\mathbb{E}[X^2] = \int_0^1 x^2 f(x) dx = \int_0^1 (6x^3 - 6x^4) dx = \frac{3}{2}x^4 - \frac{6}{5}x^5 \Big|_0^1 = \frac{3}{2} - \frac{6}{5} = \frac{3}{10} \quad (35)$$

Hence, $\text{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2 = \frac{3}{10} - \frac{1}{4} = \frac{1}{20} = \frac{5}{100}$. Which, the standard deviation of X , $\sigma_X = \sqrt{\text{Var}(X)} = \frac{\sqrt{5}}{10}$.

Now, let $a = \mathbb{E}[X] - 2\sigma_X = \frac{1}{2} - \frac{\sqrt{5}}{5}$, and $b = \mathbb{E}[X] + 2\sigma_X = \frac{1}{2} + \frac{\sqrt{5}}{5}$ (which a is 2 standard deviation below the mean, while b is 2 standard deviation above the mean). As a side not, $a > 0$ and $b < 1$ (since $0 < \frac{\sqrt{5}}{5} = \frac{1}{\sqrt{5}} < \frac{1}{2}$). Hence, the probability that X is within two standard deviations of the mean, is given by $\mathbb{P}(a \leq X \leq b)$, which has the following form of the probability:

$$\mathbb{P}(a \leq X \leq b) = \int_a^b f(x) dx = \int_a^b (6x - 6x^2) dx = 3x^2 - 2x^3 \Big|_a^b = 3(b^2 - a^2) - 2(b^3 - a^3) \quad (36)$$

$$= 3(b - a)(b + a) - 2(b - a)(b^2 + a^2 + ba) = 3(b - a)(b + a) - 2(b - a)((a + b)^2 - ba) \quad (37)$$

Which, given the values of a, b as above, here are the corresponding terms' values:

$$b - a = \left(\frac{1}{2} + \frac{\sqrt{5}}{5} \right) - \left(\frac{1}{2} - \frac{\sqrt{5}}{5} \right) = \frac{2\sqrt{5}}{5} \quad (38)$$

$$b + a = \left(\frac{1}{2} + \frac{\sqrt{5}}{5} \right) + \left(\frac{1}{2} - \frac{\sqrt{5}}{5} \right) = 1 \quad (39)$$

$$ba = \left(\frac{1}{2} + \frac{\sqrt{5}}{5} \right) \left(\frac{1}{2} - \frac{\sqrt{5}}{5} \right) = \frac{1}{4} - \frac{5}{25} = \frac{1}{4} - \frac{1}{5} = \frac{1}{20} \quad (40)$$

Hence, the probability is given as:

$$\mathbb{P}(a \leq X \leq b) = 3 \cdot \frac{2\sqrt{5}}{5} \cdot 1 - 2 \cdot \frac{2\sqrt{5}}{5} \cdot \left(1^2 - \frac{1}{20} \right) = \frac{6\sqrt{5}}{5} - \frac{4\sqrt{5}}{5} \cdot \frac{19}{20} \quad (41)$$

$$= \frac{\sqrt{5}}{5} \left(6 - \frac{19}{5} \right) = \frac{\sqrt{5}}{5} \cdot \frac{11}{5} = \frac{11\sqrt{5}}{25} \quad (42)$$

So, the probability of being within 2 standard deviations of the mean, is $\frac{11\sqrt{5}}{25} \approx 0.9839$. $\square$