

# LIE ALGEBRA OF A LIE GROUP

Zih-Yu Hsieh      Mentor: Arthur Jiang

University of California Santa Barbara



## Tangent Vectors as Derivations

When embedding smooth manifolds into  $\mathbb{R}^n$ , tangent vectors are associated with directional derivatives. To generalize tangent vectors into abstract smooth manifold, we need an analogy:

### Definition

Any point  $u \in M$ , a **Derivation at  $u$** , is a linear map  $v_u : C^\infty(M) \rightarrow \mathbb{R}$ , that satisfies the product rule:

$$\forall f, g \in C^\infty(M), \quad v_u(fg) = f(u)(v_u g) + g(u)(v_u f)$$

The vector space of all derivations at  $u$ , or  $T_u(M)$ , is the **Tangent Space** of  $M$  at  $u$ , and each derivation  $v_u \in T_u(M)$  is a **Tangent Vector** of  $u$ .

## Vector Fields & Smooth Condition

### Definition

a vector field is a map  $X : M \rightarrow TM$  ( $TM$  denotes the **Tangent Bundle**), with  $X(u) = X_u \in T_u(M)$ .

Which,  $X$  is a **Smooth Vector Field**, if  $X : M \rightarrow TM$  is a smooth map.

A collection of smooth vector fields on  $M$  is  $\mathfrak{X}(M)$ , which is an  $\mathbb{R}$ -vector space.

An equivalent condition of saying a vector field  $X$  is smooth, is through smooth functions  $f \in C^\infty(M)$ : For all  $u \in M$ ,  $X(u) = X_u \in T_u(M)$  is a derivation at  $u$ , define  $Xf : M \rightarrow \mathbb{R}$  by  $Xf(u) = X_u(f)$ , then  $X$  is a smooth vector field iff  $Xf \in C^\infty(M)$ . Which, the **Derivation** is an equivalent condition for smooth vector field:

### Theorem

For all  $f, g \in C^\infty(M)$ , given vector field  $X$ ,  $X \in \mathfrak{X}(M)$  iff it satisfies product rule for all  $u \in M$ :

$$\begin{aligned} X(fg)(u) &= X_u(fg) = f(u)(X_u g) + g(u)(X_u f) = f(u)Xg(u) + g(u)Xf(u) \\ \implies X(fg) &= f(Xg) + g(Xf) \end{aligned}$$

## Vector Fields of Different Manifolds

Given  $M, N$  two smooth manifolds, and smooth map  $F : M \rightarrow N$ . Let  $X \in \mathfrak{X}(M)$ , an ideal situation is mapping  $X$  to a smooth vector field of  $N$  through  $F$ . Yet, this requires  $F$  to be bijective:




Figure 1: Example of Conflicting Tangent Vectors

So, we'll consider a weaker notion:

### Definition

Given  $X \in \mathfrak{X}(M)$  and  $Y \in \mathfrak{X}(N)$ , the two are **F-related**, if for all  $u \in M$ , the following is true:

$$dF_u(X_u) = Y_{F(u)}$$

Simply speaking,  $F$  maps the tangent vectors collected by  $X$ , to be compatible with tangent vectors collected by  $Y$ .




Figure 2: A demonstration of  $F$ -Relation

## Lie Bracket of Vector Fields

The initial motivation is to combine two vector fields  $X, Y \in \mathfrak{X}(M)$  to be another vector field. For all  $f \in C^\infty(M)$ , since  $Yf \in C^\infty(M)$ , then  $XYf := X(Yf) \in C^\infty(M)$ . But, in general  $XY$  is not a derivation, hence not a vector field:

### Example

Define vector fields  $X = \frac{\partial}{\partial x}$ ,  $Y = x\frac{\partial}{\partial y}$  on  $\mathbb{R}^2$ . Take smooth functions  $f(x, y) = x$  and  $g(x, y) = y$ , then we get the following:

$$XY(fg) = X\left(x\frac{\partial}{\partial y}(xy)\right) = \frac{\partial}{\partial x}(x^2) = 2x$$

But, product rule doesn't hold for this example:

$$f(XYg) + g(XYf) = x\left(X\left(x\frac{\partial}{\partial y}(y)\right)\right) + y\left(X\left(x\frac{\partial}{\partial y}(x)\right)\right) = x$$

So, we need to define a new operation on vector fields:

### Definition

The **Lie Bracket**  $[\cdot, \cdot] : \mathfrak{X}(M) \times \mathfrak{X}(M) \rightarrow \mathfrak{X}(M)$ , is defined as:

$$\forall X, Y \in \mathfrak{X}(M), \quad [X, Y] = XY - YX$$

Which, the output  $[X, Y] \in \mathfrak{X}(M)$ , since it satisfies product rule:

$$\begin{aligned} [X, Y](fg) &= X(Y(fg)) - Y(X(fg)) = X(f(Yg) + g(Yf)) - Y(f(Xg) + g(Xf)) \\ &= f(XYg) + (Yg)(Xf) + g(XYf) + (Yf)(Xg) - f(YXg) - (Xg)(Yf) - g(YXf) - (Xf)(Yg) \\ &= f(XYg - YXg) + g(XYf - YXf) = f[X, Y](g) + g[X, Y](f) \end{aligned}$$

Lie Bracket also satisfies these properties:

- **Bilinearity:**  $[aX + bY, Z] = a[X, Z] + b[Y, Z]$
- **Antisymmetry:**  $[X, Y] = -[Y, X]$
- **Jacobi's Identity:**  $[X, [Y, Z]] + [Y, [Z, X]] + [Z, [X, Y]] = 0$

Moreover, Lie Bracket inherits relation of smooth maps:

### Theorem

Given smooth map  $F : M \rightarrow N$ , if  $X_1, X_2 \in \mathfrak{X}(M)$  and  $Y_1, Y_2 \in \mathfrak{X}(N)$  are  $F$ -related respectively, then  $[X_1, X_2] \in \mathfrak{X}(M)$  and  $[Y_1, Y_2] \in \mathfrak{X}(N)$  are also  $F$ -related.

## Lie Groups & Left-Invariant Vector Fields

The initial motivation is to study group structures in some smooth manifolds.

### Definition

A **Lie Group**  $G$ , is a smooth manifold along with group structure, such that the group operation  $P : G \times G \rightarrow G$  by  $P(g, h) = gh$ , and the inversion map  $i : G \rightarrow G$  by  $i(g) = g^{-1}$  are both smooth maps between manifolds.

For all  $g \in G$ , denote the left multiplication  $L_g : G \rightarrow G$  by  $L_g(h) = gh$ , since  $L_g = P|_{\{g\} \times G}$ , it is a smooth map. Hence, there's a notion of  $X$  being  $L_g$ -related to itself:

### Definition

Given any  $X \in \mathfrak{X}(G)$  and all  $g \in G$ ,  $X$  is a **Left-Invariant Vector Field**, if for all  $g \in G$ ,  $X$  is  $L_g$ -related to itself. Which, for all  $g \in G$ :

$$d(L_g)_e(X_e) = X_{L_g(e)} = X_g$$

So,  $X$  is uniquely determined by its tangent vector at identity,  $X_e \in T_e(G)$ . In fact, each  $v_e \in T_e(G)$  also corresponds to a unique Left-Invariant vector field.

The collection of Left-Invariant vector fields  $\mathfrak{g} \subseteq \mathfrak{X}(G)$ , is itself a linear subspace, and  $\mathfrak{g} \cong T_e(G)$  as vector spaces, based on the above relation.

Recall that Lie Bracket of vector field preserves  $F$ -relation between manifolds, so:

### Theorem

For all  $X, Y \in \mathfrak{X}(G)$  that are left-invariant, since for all  $g \in G$ ,  $X$  and  $Y$  are  $L_g$  related to themselves, then the Lie Bracket  $[X, Y]$  is also  $L_g$ -related to  $[X, Y]$ . Hence,  $[X, Y]$  is also left-invariant, so  $\mathfrak{g}$  is closed under Lie Bracket's operation.

## Lie Algebra on a Lie Group

### Definition

Given a vector space  $\mathfrak{g}$  over  $\mathbb{R}$  or  $\mathbb{C}$ , with a binary operation  $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ , such that the following holds:

- **Bilinearity:**  $[aX + bY, Z] = a[X, Z] + b[Y, Z]$
- **Antisymmetry:**  $[X, Y] = -[Y, X]$
- **Jacobi's Identity:**  $[X, [Y, Z]] + [Y, [Z, X]] + [Z, [X, Y]] = 0$

Then, the pair  $(\mathfrak{g}, [\cdot, \cdot])$  is a **Lie Algebra**.

In general, Lie Algebra is non-associative, so Jacobi's Identity is an alternative condition. Finally, we can define **Lie Algebra of a Lie Group**:

### Definition

Given a lie group  $G$ , since the subset of left-invariant vector fields  $\mathfrak{g} \subseteq \mathfrak{X}(G)$  forms a linear subspace, while closed under Lie Bracket's operation, then the pair  $(\mathfrak{g}, [\cdot, \cdot])$  forms a **Lie Algebra** of  $G$ , denoted as  $Lie(G)$ .

Here's an example of Lie Algebra on a Lie Group:

### Example

**General Linear Group & its Lie Algebra:**

Given  $GL_n(\mathbb{R}) \subset M_n(\mathbb{R})$ , since  $M_n(\mathbb{R}) \cong \mathbb{R}^{n^2}$  and  $GL_n(\mathbb{R})$  is an open subset, it's a natural smooth manifold with dimension  $n^2$ . The product of matrices and inversion are smooth maps, so  $GL_n(\mathbb{R})$  is a Lie Group.

Now, consider  $\mathfrak{g} = Lie(GL_n(\mathbb{R}))$ : Each  $X \in \mathfrak{g}$  is uniquely characterized by  $X_{I_n} \in T_{I_n}(GL_n(\mathbb{R}))$ . And, as vector spaces,  $\mathfrak{g} \cong T_{I_n}(GL_n(\mathbb{R}))$ .

**Lie Algebra on  $M_n(\mathbb{R})$ :**

Given  $M_n(\mathbb{R})$  as  $\mathbb{R}$ -vector space and the commutator  $[A, B] = AB - BA$ , the pair  $(M_n(\mathbb{R}), [\cdot, \cdot])$  in fact forms a Lie Algebra, denoted as  $\mathfrak{gl}_n(\mathbb{R})$ .

**Lie Algebra Isomorphism between  $\mathfrak{g}$  and  $\mathfrak{gl}_n(\mathbb{R})$ :**

$GL_n(\mathbb{R})$  has a global coordinate provided by  $M_n(\mathbb{R})$ , denote as  $(X_j^i)_{1 \leq i, j \leq n}$ .

For each  $A \in \mathfrak{gl}_n(\mathbb{R})$ , it corresponds to a tangent vector in  $T_{I_n}(GL_n(\mathbb{R}))$ :

$$A = (A_j^i) \mapsto A_j^i \frac{\partial}{\partial X_j^i} \Big|_{I_n}$$

The above tangent vector defines a Left-Invariant vector field  $A^L \in \mathfrak{g}$ . For all  $X \in \mathfrak{g}$ , the left multiplication  $L_X$  is in fact a linear operator on  $M_n(\mathbb{R})$ , so its differential is identical to itself. Which, it provides the following relation:

$$A_X^L = d(L_X)_{I_n} \left( A_j^i \frac{\partial}{\partial X_j^i} \Big|_{I_n} \right) = X_j^i A_k^j \frac{\partial}{\partial X_k^i} \Big|_X, \quad A^L = X_j^i A_k^j \frac{\partial}{\partial X_k^i}$$

Which, for arbitrary  $A, B \in \mathfrak{gl}_n(\mathbb{R})$ , Lie Bracket of  $A^L, B^L \in \mathfrak{g}$  generates:

$$[A^L, B^L] = X_j^i A_k^j \frac{\partial}{\partial X_k^i} (X_q^p B_r^q) \frac{\partial}{\partial X_r^p} - X_q^p B_r^q \frac{\partial}{\partial X_r^p} (X_j^i A_k^j) \frac{\partial}{\partial X_k^i}$$

Which, each  $A_k^j, B_r^q$  are constants, while  $\frac{\partial}{\partial X_k^i} X_r^p = 1$  iff  $(i, k) = (p, r)$  and is 0 otherwise. Then, match  $j = q$  for the same intermediate index, we get:

$$[A^L, B^L] = X_j^i (A_k^j B_r^q - B_k^j A_r^q) \frac{\partial}{\partial X_r^p} = (AB - BA)^L = [A, B]^L$$

Hence, the map  $\mathfrak{gl}_n(\mathbb{R}) \rightarrow \mathfrak{g}$  by  $A \mapsto A^L$  is a Lie Algebra Isomorphism.

## Acknowledgements & Reference

I want to thank my mentor Arthur Jiang for the effort and insights on the materials and this project, as well as UCSB Math DRP for this opportunity. Also, check out the cool project of my peer **Siyou Chen's** poster about Lie Group and Lie Algebra's applications in physics!

**Reference:** Lee, J.M. *Introduction to Smooth Manifolds*; 2nd ed.; Springer: New York, 2012; 9781441999825