

# BRAID GROUPS, AND THEIR REPRESENTATIONS

Zih-Yu Hsieh, Benjamin Schoeb, Hongxu Chen, Elijah Ecret, Mentor: Choomno Moos

University of California Santa Barbara, College of Creative Studies



## Introduction


### Braid Groups & Mapping Class Groups

**Def:** Braid group of  $n$  strands  $B_n$  is generated by  $n - 1$  elements  $\{\sigma_1, \dots, \sigma_{n-1}\}$ , satisfying *Braid Relations*:

- $\sigma_i \sigma_j = \sigma_j \sigma_i$ , if  $|i - j| \geq 2$
- $\sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}$

**Def:** Let  $D_n$  be a disk with  $n$  holes. The *Mapping Class Group*  $\mathfrak{M}(D_n)$  collects classes isotopic self-homeomorphisms on  $D_n$  that fixes disk boundary  $\partial D$ , and sends punctures to punctures.

**Ex:** The  $i^{\text{th}}$  *Half Twist* is a Self-Homeomorphism of  $D_n$ , swapping the  $i^{\text{th}}$  and  $(i + 1)^{\text{th}}$  punctures, while fixing the remaining ones.



**Figure:** For  $n = 4$ , Half Twist  $\tau_2$  Swapping Punctures 2 and 3

**Property:** Collection of *Half Twists*  $\tau_1, \dots, \tau_{n-1}$  generates  $\mathfrak{M}(D_n)$  and satisfies *Braid Relations*; in fact,  $B_n \cong \mathfrak{M}(D_n)$ , by  $\sigma_i \mapsto \tau_i$ .

### Fundamental Group of $D_n$ & Braid Automorphism

For  $n$ -punctured disk  $D - Q_n$ , fixing a base point  $d \in \partial D$ , the fundamental group  $\pi_1(D - Q_n, d)$  is generated by the  $n$  nontrivial loops, each surrounding a specific puncture, which  $\pi_1(D - Q_n, d) = F_n(x_1, \dots, x_n)$  (the *Free Group* of  $n$  generators, each  $x_i$  is a loop around the  $i^{\text{th}}$  puncture).

Then, each self-homeomorphism in the mapping class group  $\mathfrak{M}(D - Q_n)$  generates a group automorphism on the fundamental group  $\pi_1(D - Q_n, d)$ , called *Braid Automorphism*. Specifically, each  $\tau_i$  acts on the fundamental group.

Which, the corresponding Braid Automorphism is:

$$(\tau_i)_* \in \text{Aut}(\pi_1(D - Q_n, d)), \quad (\tau_i)_*(x_j) = \begin{cases} x_i x_{i+1} x_i^{-1} & j = i \\ x_i & j = i + 1 \\ x_i & \text{Otherwise} \end{cases}$$


The collection of  $(\tau_i)_*$  again satisfies the Braid Relations, which generates a group homomorphism  $B_n \rightarrow \text{Aut}(\pi_1(D - Q_n, d))$  by  $\sigma_i \mapsto (\tau_i)_*$ ; this homomorphism is in fact injective, so the braid group of  $n$  strands  $B_n$  can be viewed as a subgroup of  $\text{Aut}(\pi_1(D - Q_n, d))$ .

## Burau Representation

**Definition - Reduced Burau Representation:**  $\psi_n^r : B_n \rightarrow \text{GL}_{n-1}(\mathbb{Z}[t^{\pm}])$  satisfies:


$$\psi_n^r(\sigma_1) = \begin{pmatrix} -t & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & I_{n-3} \end{pmatrix}, \quad \psi_n^r(\sigma_{n-1}) = \begin{pmatrix} I_{n-3} & 0 & 0 \\ 0 & 1 & t \\ 0 & 0 & -t \end{pmatrix}, \quad \psi_n^r(\sigma_i) = \begin{pmatrix} I_{i-2} & 0 & 0 & 0 \\ 0 & 1 & t & 0 \\ 0 & 0 & -t & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & I_{n-i-2} \end{pmatrix}$$

**EX / Homological Perspective:** A 4-punctured disk  $D_4$  can "continuously deform" into 4 circles joining at a single point ( $\bigvee_{i=1}^4 S^1$ ),  $\implies$  Same loop structures (Fundamental Group).



**Figure:** Deformation Retraction of  $D_4$  to  $\bigvee_{i=1}^4 S^1$

Let  $S^{(4)} := \bigvee_{i=1}^4 S^1$ , consider the following space  $\tilde{S}^{(4)}$ :



**Figure:** Infinite Cyclic Cover  $\tilde{S}^{(4)}$

Here,  $t$  is a right shift of  $\tilde{S}^{(n)}$  by degree 1:

- $t^k \cdot \tilde{d} = \text{degree } k \text{ right shift of } \tilde{d}$
- $t^k \cdot \hat{x}_i = \text{degree } k \text{ right shift of } \hat{x}_i$


There is a continuous map  $p : \tilde{S}^{(4)} \rightarrow S^{(4)}$ , each  $p(t^k \cdot \hat{x}_i) = x_i$ , and  $p(t^k \cdot \tilde{d}) = d$ . Define the "Base Loops"  $\ell_i := \hat{x}_{i+1} \cdot \hat{x}_i^{-1}$  (counterclockwise) for  $1 \leq i \leq 3$ :

- $-\ell_i = \text{counterclockwise version of } \ell_i$
- $t^k \cdot \ell_i = \text{degree } k \text{ right shift of } \ell_i$

Then, all "Integer Laurent Polynomial" combination of  $\ell_i$  forms  $H_1(\tilde{S}^{(4)})$  as a free  $\mathbb{Z}[t^{\pm}]$ -module with basis  $\ell_1, \ell_2, \ell_3$ .

**Recall:** braid automorphism  $(\tau_2)_*$  of  $\pi_1(D_4, d)$  satisfies  $(\tau_2)_*(x_2) = x_2 \cdot x_3 \cdot x_2^{-1}$ , and  $(\tau_2)_*(x_3) = x_2$ . Which, it uniquely lifts to an transformation on the  $\ell_i$  via  $p$ :

**EX:**  $\ell_2 = \hat{x}_3 \cdot \hat{x}_2^{-1} \mapsto \hat{x}_2 \cdot ((t \cdot \hat{x}_2) \cdot (t \cdot \hat{x}_3^{-1}) \cdot \hat{x}_2^{-1}) = -t \cdot \ell_2$ .



**Figure:**  $\ell_2$  (Counterclockwise) Maps to  $-t \cdot \ell_2$  (Right Shift by degree 1, Clockwise)

Doing this for each  $\ell_i$ , and write into matrix form with basis  $\ell_i$ , we recover the Representation.

## Conclusion & Future Directions

### Acknowledgement & Sources

We're genuinely thankful for the parent donors, Professor Cachadina and Professor Casteels who made this program possible. We also want to thank our mentor Choomno Moos for their great guidance.

- Braids, Links, Mapping Class Groups (Joan Birman)
- Briad Groups (Christian Kassel, Vladimir Turaev)
- Category Theory in Context (Emily Riehl)
- Algebra Chapter 0 (Paolo Aluffi)