

# BRAID GROUPS, AND THEIR REPRESENTATIONS

Zih-Yu Hsieh      Mentor: Choomno Moos

University of California Santa Barbara, College of Creative Studies

## Introduction

### Braid Groups & Mapping Class Groups


**Def:** Braid group of  $n$  strands  $B_n$  is generated by  $n - 1$  elements  $\{\sigma_1, \dots, \sigma_{n-1}\}$ , satisfying *Braid Relations*:

•  $\sigma_i \sigma_j = \sigma_j \sigma_i$ , if  $|i - j| \geq 2$

•  $\sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}$

**Def:** Let  $D_n$  be an  $n$ -punctured disk. The *Mapping Class Group*  $\mathcal{M}(D_n)$  collects classes of isotopic self-homeomorphisms on  $D_n$  that fixes disk boundary  $\partial D$ .

**Ex:** The  $i^{\text{th}}$  *Half Twist*  $\tau_i \in \mathcal{M}(D_n)$  swaps the  $i^{\text{th}}$  and  $(i + 1)^{\text{th}}$  punctures, while fixing the remaining ones.




**Figure:** For  $n = 4$ , Half Twist  $\tau_2$  Swapping Punctures 2 and 3

**Property:** *Half Twists*  $\tau_1, \dots, \tau_{n-1}$  generates  $\mathcal{M}(D_n)$  and satisfies *Braid Relations*; in fact,  $B_n \cong \mathcal{M}(D_n)$ , by  $\sigma_i \mapsto \tau_i$ .

### Fundamental Group of $D_n$ & Braid Automorphism

For  $n$ -punctured disk  $D_n$ , fix  $d \in \partial D$ , the fundamental group  $\pi_1(D_n, d)$  is generated by the  $n$  loops, each surrounding a puncture, which  $\pi_1(D_n, d) = F_n(x_1, \dots, x_n)$ , the *Degree- $n$  Free Group*.



**Figure:** Fundamental Group of  $D_4$

Then, each homeomorphism in  $\mathcal{M}(D_n)$  generates a group automorphism on  $\pi_1(D_n, d)$ , called *Braid Automorphism*.

**Ex:** Half Twist's action on  $\pi_1(D_n, d)$ :

$$(\tau_i)_* \in \text{Aut}(\pi_1(D_n, d)), \quad (\tau_i)_*(x_j) = \begin{cases} x_{i+1} & j = i \\ x_{i+1}x_i x_{i+1}^{-1} & j = i + 1 \\ x_i & \text{Otherwise} \end{cases}$$

**Figure:**  $\tau_2$  Action on Loops in  $D_4$


## Reduced Burau Representation

$\psi_n^r : B_n \rightarrow \text{GL}_{n-1}(\mathbb{Z}[t^{\pm}])$  satisfies:

$$\sigma_1 \mapsto \begin{pmatrix} -t & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & I_{n-3} \end{pmatrix}, \quad \sigma_{n-1} \mapsto \begin{pmatrix} I_{n-3} & 0 & 0 \\ 0 & 1 & t \\ 0 & 0 & -t \end{pmatrix}, \quad \sigma_i \mapsto \begin{pmatrix} I_{i-2} & 0 & 0 & 0 \\ 0 & 1 & t & 0 \\ 0 & 0 & -t & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & I_{n-i-2} \end{pmatrix}$$


### Ex: Homological Perspective on $D_4$

A 4-punctured disk  $D_4$  can "continuously deform" into 4 circles joining at one point ( $\bigvee_{i=1}^4 S^1$ ),  $\implies$  Same Fundamental Group.



**Figure:** Deformation Retraction of  $D_4$  to  $\bigvee_{i=1}^4 S^1$

Let  $S^{(4)} := \bigvee_{i=1}^4 S^1$ , consider the space  $\tilde{S}^{(4)}$  below, a *Covering Space* of  $S^{(4)}$ :



**Figure:** Infinite Cyclic Cover  $\tilde{S}^{(4)}$

Here,  $t$  is a right shift of  $\tilde{S}^{(n)}$  by degree 1:

- $t^k \cdot \tilde{d}$  = degree  $k$  right shift of  $\tilde{d}$
- $t^k \cdot \hat{x}_i$  = degree  $k$  right shift of  $\hat{x}_i$

There is a continuous covering map  $p : \tilde{S}^{(4)} \rightarrow S^{(4)}$ , each  $p(t^k \cdot \hat{x}_i) = x_i$ , and  $p(t^k \cdot \tilde{d}) = d$ .

Define the "Base Loops"  $\ell_i := \hat{x}_{i+1} \cdot \hat{x}_i^{-1}$  (counterclockwise) for  $1 \leq i \leq 3$ :


- $-\ell_i$  = clockwise version of  $\ell_i$
- $t^k \cdot \ell_i$  = degree  $k$  right shift of  $\ell_i$

Then, all "Integer Laurent Polynomial" combination of  $\ell_i$  forms  $H_1(\tilde{S}^{(4)})$  as a free  $\mathbb{Z}[t^{\pm}]$ -module with basis  $\ell_1, \ell_2, \ell_3$ .

## Braid Group's Action on Covering Space

**Recall:** braid automorphism  $(\tau_2)_*$  of  $\pi_1(D_4, d)$  satisfies  $(\tau_2)_*(x_2) = x_3$ , and  $(\tau_2)_*(x_3) = x_3 \cdot x_2 \cdot x_3^{-1}$ . Which, it uniquely lifts to an transformation on the  $\ell_i$  via  $p$ :

**EX:**  $\ell_2 = \hat{x}_3 \cdot \hat{x}_2^{-1} \mapsto \hat{x}_2 \cdot ((t \cdot \hat{x}_2) \cdot (t \cdot \hat{x}_3^{-1}) \cdot \hat{x}_2^{-1}) = -t \cdot \ell_2$ .



**Figure:**  $\ell_2$  (Counterclockwise) Maps to  $-t \cdot \ell_2$  (Right Shift by degree 1, Clockwise)

Doing this for each  $\ell_i$ , put into matrix form with basis  $\{\ell_i\}$ , we recover the Representation.

## Conclusion & Future Directions

### Acknowledgement & Sources

We're genuinely thankful for the parent donors, Professor Cachadina and Professor Casteels who made this program possible. We also want to thank our mentor Choomno Moos for their great guidance.

- Braids, Links, Mapping Class Groups (Joan Birman)
- Braid Groups (Christian Kassel, Vladimir Turaev)
- Category Theory in Context (Emily Riehl)
- Algebra Chapter 0 (Paolo Aluffi)